

## A Study to Evaluate the Efficacy of Platelet Count (Platelets) and its Indices in the Early Diagnosis of Sepsis

Manisha Kumari<sup>1</sup>, Ravindra Kumar<sup>2</sup>, Jayant Prakash<sup>3</sup>

<sup>1</sup>Senior Resident, Department of Pediatrics, IGIMS, Patna, Bihar, India

<sup>2</sup>Senior Resident, Department of Pediatrics, IGIMS, Patna, Bihar, India

<sup>3</sup>Professor and HOD, Department of Pediatrics, IGIMS, Patna, Bihar, India

Received: 04-01-2024 / Revised: 10-02-2024 / Accepted: 25-02-2024

Corresponding Author: Dr. Ravindra Kumar

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the study was to evaluate the efficacy of platelet count (Platelets) and its indices in the early diagnosis of sepsis.

**Methods:** This was a prospective study conducted in the Department of Pediatrics, IGIMS, Patna, Bihar, India for the period of two years. Sample size of 100 patients was recruited as per hospital neonatal intensive care unit work load during the period of study.

**Results:** The general characteristics of a study group showing 100 patients including 60 males and 40 females. On the basis of NNF guidelines, the patients were divided in three groups - 50% were clinically suspected sepsis (probable sepsis), 35% were with blood culture positive (culture positive sepsis), and remaining 15% in blood culture negative but CRP positive group (culture negative sepsis). The patients were divided into early and late onset. EOS was seen more in males (65%) as compared to females (35%). However, in LOS both male and females were equally affected.

**Conclusion:** Platelets can be considered as an early diagnostic tool for neonatal sepsis as it is cheap, rapid, and easily available and does not require additional equipment.

**Keywords:** Neonatal sepsis, Platelet indices, Thrombocytopenia.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Neonatal sepsis is major cause of neonatal morbidity and mortality worldwide contributing around 38% of all deaths in neonates. Situation is even more worsened in low income underdeveloped countries. As per the National Neonatal Perinatal Database 2002-2003, the incidence of neonatal sepsis is 30 per 1000 live births. [1] Globally, the Neonatal mortality rate has declined by 47% between 1990 and 2015 from 36 to 19 deaths per 1000 live births. [2] India contributes to one-fifth of global live births and more than a quarter of neonatal deaths. [3] Neonatal sepsis is one of the major causes of morbidity and mortality among newborns in the developing world and is more common when compared with developed countries. [4] In 2013, the global analysis revealed that the major contributing factors for mortality in neonates were infections and preterm with associated complications in developing countries. [3] Host defense against infection is based on two crucial mechanisms: the inflammatory response and the activation of coagulation. Platelets are involved in both hemostasis and immune response. These

mechanisms work together in a complex and synchronous manner making the contribution of platelets of major importance in sepsis and platelet count (Platelets) has been used as a marker of sepsis historically. National Neonatal Forum of India (NNF) has classified sepsis into probable, culture-negative, and culture-positive sepsis. [5]

Based on the onset of sepsis, neonatal sepsis is classified into early (<72 h) and late (more than 72 h) onset. In early-onset sepsis (EOS), infants acquire infection by a vertical transmission that is through the bacterial flora of the mother. In late-onset sepsis (LOS), the acquisition of infection is predominantly through the infant's environment. [6] Coagulase-negative Staphylococci (30.27%) were the most common organisms isolated followed by *Acinetobacter* sp (15.1%), *Klebsiella* sp (5.4%) *Staphylococcus aureus* (4.8%), and *Escherichia coli* (4.8%) in a study conducted by Asifa Nazir. [7] For a definite diagnosis of sepsis blood culture has always been the gold standard. [8] However, the blood culture has a huge disadvantage of delay in availability of the result along with other chances of false-negative results.

[9] Other tests such as Procalcitonin and IL-6 are very expensive and not easily affordable by most of the patients in developing countries. [10]

The aim of the study was to evaluate the efficacy of platelet count (Platelets) and its indices in the early diagnosis of sepsis.

### Materials and Methods

This was a prospective study conducted in the Department of Pediatrics, IGIMS, Patna, Bihar, India for the period of two years. Sample size of 100 patients was recruited as per hospital neonatal intensive care unit work load during the period of study.

Neonates delivered in IGIMS, Patna, Bihar, India and referred from outside with features of suspected septicemia or probable sepsis as per NNF criteria of poor feeding, irritability, excessive cry, lethargy poor cry and reflexes, fever, hypothermia, jaundice, vomiting, abdominal distension, tachypnea and grunting, convulsions, diarrhea, pustules, cyanosis, bulge fontanelle, disseminated intravascular coagulation, bleeding, poor perfusion, shock, and apnea were included in the study. Neonates who developed sepsis during the period of hospitalization for other reasons and proved to have sepsis by positive blood culture, cerebrospinal fluid (CSF), and urine culture were also included in the study while neonates with congenital anomalies of Gastrointestinal Tract system such as trachea-esophageal fistula, malrotation of gut, anomalies of

the cardiovascular system or respiratory system, central nervous system, and inborn error of metabolism were excluded from the study.

At admission, thorough history was taken and clinical examination was done. Sepsis profile was sent including hemoglobin, total leukocyte count, differential leukocyte count, Platelets and its indices-MPV, PDW and PCT, absolute neutrophil count (ANC), C-reactive protein (CRP), band cell ratio (BCR), and blood culture for all the patients. BECKMAN COULTER LS-750 Analyzer was used and as when required X-ray chest, Urine C/S, and CSF culture was done. The patients were categorized into three groups-Clinically suspected sepsis (probable sepsis), culture positive sepsis, and culture negative as per NNF criteria of sepsis screen comprising of total leukocyte count, I/T ratio (BCR), ANC, m-ESR e, and CRP. [11,12]

Statistical Package for the Social Sciences 20.0 version of statistical analysis was used. Sensitivity, Specificity, Positive predictive value (PPV) and Negative predictive value (NPV) of Platelets and its indices were evaluated and compared between probable sepsis with culture negative sepsis and probable sepsis with culture positive sepsis. Non-parametric tests like the Chi-square test were applied to see the association between the variables. The three groups were compared for Platelets and its indices with Analysis of variance (ANOVA) test.

### Results

**Table 1: Patient details**

Variables	N%
<b>Gender</b>	
Male	60 (60)
Female	40 (40)
<b>Types according to NNF</b>	
Suspected Sepsis	50 (50)
Culture positive sepsis	35 (35)
Culture Negative sepsis	15 (15)

The general characteristics of a study group showing 100 patients including 60 males and 40 females. On the basis of NNF guidelines, the patients were divided in three groups - 50% were clinically suspected sepsis (probable sepsis), 35% were with blood culture positive (culture positive sepsis), and remaining 15% in blood culture negative but CRP positive group (culture negative sepsis). The patients were divided into early and

late onset. Out of 40 preterm subjects, 25 (62.5%) were male and 15(37.5%) were female whereas out of 60 term subjects 36(60%) were male and 24(40%) were females. EOS was seen more in males (65%) as compared to females (35%). Probable sepsis was more among preterm babies whereas culture negative and culture positive were more in term babies.

**Table 2: Sensitivity and specificity of platelet count and its indices in probable sepsis and culture positive sepsis**

	Probable sepsis	Culture positive sepsis

Parameters	Platelet count	PDW	MPV	Plateletcrit	Platelet count	PDW	MPV	Plateletcrit
Specificity	68	62.5	62.5	6.2	68	62	62.5	6.25
Sensitivity	69	28	34	86	52.9	14	26.4	17
NPV	21.5	21.7	23.2	12.5	40.7	25.6	28	3.4
PPV	66.6	70	73.9	74.1	78.2	45	60	28.5
Accuracy	57.8	36	40.9	66.6	58	30	38	14

Sensitivity and accuracy for Platelets were 52.9% and 58% in the culture positive group, whereas sensitivity and accuracy for Platelets in the probable sepsis group was 69% and 57.5%. Specifically, NPV and PPV was 68%, 40.7%, and 78.2% in culture positive whereas it was 68%, 21.5%, and 66.6% in the probable sepsis group.

**Table 3: Comparison of normal with probable sepsis and culture positive**

Parameters	Normal	Probable sepsis (mean)	Culture + ve sepsis (mean)	Culture negative CRP + ve (mean)	p-value
Platelet count	150–400	219.2	174.8	205.78	0.030
PDW	8.3–56.6%	12.5	15.1	11.05	0.170
MPV	7.2–11.7fl	10.9	9.94	11.9	0.550
Plateletcrit	0.22–0.24%	2.5	1.46	3.2	0.60

Analysis on the basis of ANOVA for three groups—culture positive, culture negative, and probable sepsis was done for Platelets and its indices (MPV, PDW, and PCT). Platelets was found to be significant in all groups.

#### Discussion

Neonatal sepsis is major cause of neonatal morbidity and mortality worldwide contributing around 38% of all deaths in neonates. Situation is even more worsened in low income underdeveloped countries. As per the National Neonatal Perinatal Database 2002-2003, the incidence of neonatal sepsis is 30 per 1000 live births. [13]

The need for early recognition, diagnosis of neonatal septicemia, and prompt institution of treatment is paramount to prevent death (s) and complications associated with neonatal septicemia. The definitive diagnosis of sepsis is made by blood culture, which requires a minimum of 48–72 h, yields a positive result in 30–40% of cases. One of the most common hematological manifestations seen during early sepsis is thrombocytopenia. Platelet indices; PCT, MPV, and PDW are a group of derived platelet parameters obtained as a part of the automatic complete blood count. PI are biomarkers of platelet activation. Average mean cell volume is 7.2–11.7 fL Increased MPV indicates increased platelet diameter, which can be used as a marker of production rate and platelet activation. During activation, platelets' shapes change from biconcave discs to spherical, and a pronounced pseudopod formation occurs that leads to MPV increase during platelet activation. The PDW reported varies markedly, with reference intervals ranging from 8.3 to 56.6%. Under physiological conditions, there is a direct relationship between MPV and PDW; both usually

change in the same direction. PCT is the volume occupied by platelets in the blood as a percentage. The normal range for PCT is 0.22–0.24%.

Platelets play an important role in inflammation. Septic patients are observed to have low Platelets due to production of many cytokines, endothelial damage and bone marrow suppression. PI have been shown to have diagnostic value in certain inflammatory diseases, such as inflammatory bowel diseases, rheumatoid arthritis, ankylosing spondylitis, ulcerative colitis, and atherosclerosis. [14] In our study, sepsis was predominantly present in males (60%) as compared to females (40%) with early onset sepsis in 64% and 36% had late onset sepsis. In our study, there were 50% probable sepsis cases, 34% culture proven cases and 16% culture negative sepsis. In a similar study conducted by Salama et al. 39.8% of neonates had probable sepsis and were culture positive. [15]

Naik et al. reported thrombocytopenia in 66.25% of neonates with sepsis having 66.25% sensitivity and 87% specificity. [16] Culture positive patients in our study were found to have thrombocytopenia in 72% of neonates having 52.9% of sensitivity and 68% of specificity with NPV 40.7% and PPV 78.2%. Probable sepsis neonates also had decreased Platelets with sensitivity and specificity as 20% and 68%, respectively, with NPV of 21.5 and PPV of 6. In culture positive cases, in our study, PDW sensitivity is 14% with specificity of 62% and NPV of 25.6 with PPV of 45. While in probable sepsis there was sensitivity of 62% with specificity of 62.5% with NPV of 21.7 and PPV of 70. Sangsari et al. in their study on PI in different types of germs found that PDW was elevated in all culture positive infections specifically in Gram-negative sepsis. [17] PCT in culture proven cases had sensitivity of 17% and specificity of 6.25% with NPV of 3.4 and PPV of 28.5 while in probable sepsis PCT showed

sensitivity of 86% with specificity of 62% and NPV of 12.5 with PPV of 74.1 (Fig. 4). Very few studies show changes in PCT. Sandeep et al. reported high PDW, low PCT, and high but not significant MPV in preterm neonates which were compared with term neonate. [18]

The Platelets was significantly low (p value-0.030) in all three groups, probable sepsis, culture positive sepsis and culture negative sepsis; however, the PI such as PDW, MPV, and PCT did not show any significant difference in the various groups. Platelets may be considered an alternative, novel diagnostic tool for detection of early neonatal sepsis as it is cheap, rapid, and easily available and does not require additional equipment.

### Conclusion

Platelets can be considered as an early diagnostic tool for neonatal sepsis as it is cheap, rapid, and easily available and does not require additional equipment. High PDW, high MPV and low platelet count are more associated with neonatal sepsis.

### References

1. Shankar MJ, Aggarwal R, Deorari AK, Paul VK. Symposium on AIIMS protocols in neonatology III- Sepsis in new-born. *Indian J Pediatr.* 2008; 75(103):261-6.
2. Neonatal Mortality; 2021.
3. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, Cousens S, Mathers C, Black RE. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *The lancet.* 2015 Jan 31;385(9966):430-40.
4. GHO by Country India Statistics Summary (2002 Present); 2017.
5. Klingenberg C, Kornelisse RF, Buonocore G, Maier RF, Stocker M. Culture-negative early-onset neonatal sepsis—at the crossroad between efficient sepsis care and antimicrobial stewardship. *Frontiers in pediatrics.* 2018 Oct 9; 6:285.
6. Richard J, Fanaroff AA, Walsh M. Postnatal Bacterial infections. In: Ethan L, Dobbs K, editors. *Fanaroff and Martin's Neonatal Perinatal Medicine.* 10th ed. Philadelphia, PA: Elsevier Health Sciences; 2015. p. 734-50.
7. Ghotaslou R, Ghorashi Z, Nahaei MR. Klebsiella pneumoniae in neonatal sepsis: A 3-year-study in the pediatric hospital of Tabriz, Iran. *Jpn J Infect Dis* 2007;60:126-8.
8. Sonawane VB, Mehkarkar N, Gaikwad S, Kadam N. Comparison between sepsis markers and blood culture in diagnosis of neonatal sepsis: A prospective study. *Int J Res Med Sci* 2017; 5:1662-6.
9. Benitz WE. Adjunct laboratory tests in the diagnosis of early-onset neonatal sepsis. *Clin Perinatol* 2010; 37:421-38.
10. Ku LC, Boggess KA, Cohen-Wolkowicz M. Bacterial meningitis in infants. *Clin Perinatol* 2015; 42:29.
11. Polinski C. The value of the white blood cell count and differential in the prediction of neonatal sepsis. *Neonatal network: NN.* 1996 Oct 1;15(7):13-23.
12. Da Silva O, Ohlsson A, Kenyon C. Accuracy of leukocyte indices and C-reactive protein for diagnosis of neonatal sepsis: a critical review. *The Pediatric infectious disease journal.* 1995 May 1;14(5):362-6.
13. Shankar MJ, Aggarwal R, Deorari AK, Paul VK. Symposium on AIIMS protocols in neonatology III- Sepsis in new-born. *Indian J Pediatr.* 2008; 75(103):261-6.
14. Margetic S. Inflammation and haemostasis: *Biochem Med (Zagreb), v. 22.*
15. Salama K, Gad A, El Tatawy S. Sepsis profile and outcome of preterm neonates admitted to neonatal intensive care unit of Cairo University Hospital. *Egyptian Pediatric Association Gazette.* 2021 Dec; 69:1-9.
16. Bhat S, Naik S, Rafiq W, Tariq A S. Incidence of thrombocytopenia and changes in various platelet parameters, in blood culture positive neonatal sepsis. *International Journal of Pediatrics.* 2015 Jul 1;3(4.1):757-66.
17. Sangsari R, Kadivar M, Fardi R, Attary SJ, Jafari A. Platelets indices and different germs of neonatal sepsis. *Iranian Journal of Pediatrics.* 2019 Apr 30;29(2).
18. Sandeep M, Thammanna PS, Sridhar PV. Platelet indices in preterm neonates: a prospective study. *International journal of scientific study.* 2015;3(7):237-40.