

## A Study to Investigate the Correlation between Vitamin D and Cellular Senescence As Assessed by the Telomerase Enzyme in Individuals with Pre-HTN: A Retrospective Study

Manoj Kumar Singh<sup>1</sup>, Sanjeeb Sharan<sup>2</sup>, H. P. Dubey<sup>3</sup>

<sup>1</sup>Tutor, Department of Physiology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar, India

<sup>2</sup>Assistant Professor, Department of Physiology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar, India

<sup>3</sup>Professor and HOD, Department of Physiology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar, India

Received: 10-12-2023 / Revised: 17-01-2024 / Accepted: 21-02-2024

Corresponding Author: Dr. Sanjeeb Sharan

Conflict of interest: Nil

### Abstract

**Aim:** The aim of the present study was to explore the link between Vitamin D and cellular senescence measured with the enzyme telomerase in pre-HTN.

**Methods:** The present study was conducted from in the Department of Physiology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar, India for two years . Inclusion criteria for the pre- hypertensive group (pre-HTN) (n =50) were both genders between 18 and 25 years of age with SBP between 120 and 139 mmHg and DBP between 80 and 89 mmHg in apparently healthy individuals. The controls (n = 50) population were healthy individuals with 18– 25 years of age with SBP between 100 and 119 mmHg and DBP between 60 and 79 mmHg.

**Results:** The study population included 100 apparently healthy individuals. 50 were pre- hypertensive with the age of 22.58±1.56 and the age of controls was 18.82±1.22. Out of 100, 27 males, 23 females in pre- HTN group and 26 males, 24 females were in the control group. A significant difference was not found between-group differences in height and waist-hip ratio. However, pre- HTN group subject's BMI (P < 0.001) and weight (P < 0.001) was more compared to controls. In pre-HTN group, significantly higher HR (P < 0.001), SBP (P < 0.001), DBP (P < 0.001), MAP (P < 0.001), and RPP (P < 0.001) were seen when compared to controls. No significant difference was seen in PP but it was slightly high in pre-HTN group and negatively associated with Vitamin D. High telomerase levels have correlation with waist-hip ratio, SBP, DBP, MAP, and RPP but no significant correlation was seen with BMI, HR, and PP.

**Conclusion:** It can be concluded that reduced Vitamin D levels in pre-HTN may cause derangements of cardiovascular homeostatic mechanism, enhance the speed of cellular senescence measured by telomerase.

**Keywords:** Hypertension, Vitamin D, Cellular Senescence.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

### Introduction

Years of research have revealed the association of three conditions with the development of subclinical atherosclerosis and non-alcoholic fatty liver disease (NAFLD): obesity, vitamin D deficiency (VDD), and cellular senescence. Obesity is a serious worldwide chronic health problem for all ages. [1] Cellular senescence induces a cell-cycle arrest and a pro-inflammatory reaction, both of which promote aging and age-related diseases, and its exacerbation was seen with obesity in many studies. [2] VDD was reported to be induced by obesity among all age groups and in both genders. [1,3] Numerous genetic studies explored that excessive adiposity causes a reduction in

circulating 25-hydroxyvitamin D (25(OH)D), the main reflector of vitamin D status. [1,3]

VDD in obese people usually has no direct consequence but may affect many organs at the subclinical level and predispose them to a state of improper metabolism. [4] On the other hand, the effects of vitamin D supplementation on obese individuals do not largely extend beyond ameliorating the detrimental consequences of various obesity-induced cardiometabolic disorders, and no evidence proves a protective effect of vitamin D against obesity. [1,3] Obesity promotes cardiovascular diseases through vascular remodeling, which induces subclinical

atherosclerosis that finally ends in fatal cardiac events. [5]

High blood pressure (hypertension) is a serious risk factor for cardiovascular diseases, such as coronary artery disease, myocardial infarction, or stroke, if untreated. [6] Study results revealed that vitamin D deficiency ameliorates the development of hypertension (HT). [6,7] Vitamin D deficiency (25-OH-D < 30 ng/mL) is an independent risk factor for high blood pressure and is involved in the promotion of cardiovascular mortality. [8] Obesity is thought to affect cardiovascular events. Indeed, the concept of “metabolically healthy” obesity has been studied, and it has been shown that coronary heart disease may be increased in this obesity phenotype. [9] Low 25(OH)D3 concentrations increase the risk of hypertension, peripheral vascular disease, diabetes mellitus, myocardial infarction, heart failure, and cardiac mortality. [10] Moreover, it is reported that low 25(OH)D levels are associated with endothelial dysfunction, inflammation, increased vascular stiffness, and high coronary artery calcium scores. [11] Moreover, it is reported that shortened telomere as an early sign and characteristic feature of cellular senescence is a prognostic biomarker for the early identification of subjects at high risk of developing CVD before symptoms appear. [12]

The aim of the present study was to explore the link between Vitamin D and cellular senescence measured with the enzyme telomerase in pre-HTN.

#### Materials and methods

**Table 1: Comparison of anthropometric characteristics between pre-HTN and controls**

Parameters	Pre-HTN (n=50)	Controls (n=50)	P-value
Age	22.58±1.56	18.82±1.22	0.432
Gender (male/female)	27/23	26/24	1.346
Height (cm)	172.28±9.81	169.72±8.52	0.386
Weight (kg)	65.45±11.39	58.42±8.42	<0.001
BMI (k/m <sup>2</sup> )	23.17±4.82	21.69±4.90	<0.001
Waist to hip ratio	0.96±0.14	0.86±0.07	0.474

The study population included 100 apparently healthy individuals. 50 were pre- hypertensive with the age of 22.58±1.56 and the age of controls was 18.82±1.22. Out of 100, 27 males, 23 females in pre- HTN group and 26 males, 24 females were in

The present study was conducted the Department of Physiology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar, India for two years . Inclusion criteria for the pre- hypertensive group (pre-HTN) (n =50) were both genders between 18 and 25 years of age with SBP between 120 and 139 mmHg and DBP between 80 and 89 mmHg in apparently healthy individuals. The controls (n = 50) population were healthy individuals with 18– 25 years of age with SBP between 100 and 119 mmHg and DBP between 60 and 79 mmHg.

Individuals suffering from diabetes, hypertension, endocrine disorders, kidney diseases, and hypertensive patients already receiving medication were not considered to take part in this research. The volunteers were asked to not participate in heavy exercises, not drink alcohol and coffee 1 day before the data collection. Baseline, anthropometric parameters were recorded before recording of the BP by sphygmomanometer as per standard protocol. [13] Then, 5 ml of blood was collected, allowed to clot, and subjected to centrifugation to separate the serum. Serum was stored at –80°C for processing of Vit-D and telomerase levels as per the instructions provided in the commercially available kits.

Statistical analysis was done to analyze the data. To study the between-group differences, independent t-test, to assess the correlation of vitamin D with telomerase and other parameters, Pearson’s correlation coefficient analysis was applied.

#### Results

the control group. A significant difference was not found between-group differences in height and waist-hip ratio.

**Table 2: Comparison of cardiovascular parameters between pre-HTN and controls**

Parameters	Pre-HTN (n=50)	Controls (n=50)	P-value
HR (BPM)	87.43±5.75	82.48±5.25	<0.001
SBP (mmHg)	122.18±4.86	114.6±4.56	<0.001
DBP (mmHg)	82.68±4.06	76.34±3.96	<0.001
PP (mmHg)	41.73±5.90	39.01±5.55	0.175
MAP (mmHg)	94.56±2.82	87.73±3.72	<0.001
RPP	10950.80±704.76	9082.78±490.82	<0.001

However, pre- HTN group subject’s BMI (P < 0.001) and weight (P < 0.001) was more compared to controls. In pre-HTN group, significantly higher HR (P < 0.001), SBP (P < 0.001), DBP (P < 0.001), MAP (P < 0.001), and

RPP ( $P < 0.001$ ) were seen when compared to controls. No significant difference was seen in PP but it was slightly high in pre-HTN group and negatively associated with Vitamin D.

**Table 3: Comparison of Vitamin D and telomerase levels between pre-HTN and controls**

Parameters	Pre-HTN (n=50)	Controls (n=50)	P-value
Vitamin D (ng/ml)	19.21±4.36	23.04±6.24	0.044
Telomerase (IU/ml)	35.85±18.82	7.03±5.95	<0.001

High telomerase levels have correlation with waist-hip ratio, SBP, DBP, MAP, and RPP but no significant correlation was seen with BMI, HR, and PP.

### Discussion

Hypertension is a common health problem, one of the leading costs to the health care system, and a significant cause of mortality and morbidity worldwide. [14] Hypertension is also one of the most common and influential risk factors of cardiovascular disease including myocardial infarction, cerebral stroke, congestive heart failure, peripheral vascular disorders and kidney disease. [15] It has been estimated that eliminating high blood pressure would reduce the occurrence of stroke by 35% and heart attacks by 18%. [16,17] To reduce the burden of hypertension, a multicomponent lifestyle intervention that includes weight loss, increased physical activity, restricted sodium and alcohol consumption, and adherence to a Dietary Approach to Stop Hypertension like diet with plenty of fruits, vegetables, and low-fat dairy items and little saturated fat is needed. [18] Moreover, improved vitamin D status has been proposed as an easily modifiable risk factor. [19]

The study population included 100 apparently healthy individuals. 50 were pre-hypertensive with the age of 22.58±1.56 and the age of controls was 18.82±1.22. Out of 100, 27 males, 23 females in pre-HTN group and 26 males, 24 females were in the control group. A significant difference was not found between-group differences in height and waist-hip ratio. Vitamin D deficiency has recently emerged as a public health problem, affecting almost 50% of the population worldwide. [20] In addition to the reduced exposition to sunlight [21], also genetic and environmental factors have been suggested as a cause of this pandemic, such as pollution, diet, sedentary life style and stress. [22] Moreover, vitamin D is no longer considered as only a pivotal mediator of calcium metabolism and skeletal health, but it also regulates several cell functions, including differentiation and metabolism. This aspect may explain the reason why hypovitaminosis D has been proved to be an independent risk factor for overall mortality in various cohort analyses [23], whereas vitamin D supplementation significantly reduced mortality. [24] Zhao et al [25] in an ongoing report detailed a positive relationship between Vit-D and hypertension and pre-HTN. Forman et al [26]

reported a positive relationship between Vit-D and self-revealed occurrence hypertension among 38,388 men from the Health Professionals' follow-up study and 77,531 females from the Nurses' Health Study; a positive affiliation was likewise detailed between Vit-D and hypertension in a subsample of members. Further, a study concentrate from the second Nurses' health study detailed a positive relationship between serum Vit-D and hypertension among 1484 young females. In the NHANES, SBP was demonstrated to be conversely connected with Vit-D among 12,644 participants. [27]

However, pre-HTN group subject's BMI ( $P < 0.001$ ) and weight ( $P < 0.001$ ) was more compared to controls. In pre-HTN group, significantly higher HR ( $P < 0.001$ ), SBP ( $P < 0.001$ ), DBP ( $P < 0.001$ ), MAP ( $P < 0.001$ ), and RPP ( $P < 0.001$ ) were seen when compared to controls. No significant difference was seen in PP but it was slightly high in pre-HTN group and negatively associated with Vitamin D. High telomerase levels have correlation with waist-hip ratio, SBP, DBP, MAP, and RPP but no significant correlation was seen with BMI, HR, and PP. Earlier reports have shown that higher Vitamin D is related to longer telomere length, which underscores the conceivably advantageous impacts of this hormone on cell senescence and age-related conditions. [28] In this study, cellular senescence was assessed using telomerase. This enzyme attempts to inhibit the process of telomere shortening. [29] Since the cell telomere loss appears to result from cell division just to a fractional degree, different components, particularly oxidative stress, were attested to assume a job in the expanded rate for shortening of telomeres. [30] The exact mechanism by which lower Vit-D levels are associated with this cellular senescence is hypothesized dependent on the perceptions recommends that the degrees of the telomerase may really be related to oxidative stress, with higher oxidative stress prompting higher telomerase levels. Cells of nearly complex organism may not have an ability to divide. This marvel was depicted by Hayflick in 1961. [31]

### Conclusion

It can be concluded that reduced Vitamin D levels in pre-HTN may cause derangements of cardiovascular homeostatic mechanism, enhance the speed of cellular senescence measured by telomerase.

## References

1. Hyppönen E, Boucher BJ. Adiposity, vitamin D requirements, and clinical implications for obesity-related metabolic abnormalities. *Nutrition Reviews*. 2018 Sep 1;76(9):678-92.
2. Ogrodnik M, Zhu YI, Langhi LG, Tehkonia T, Krüger P, Fielder E, Victorelli S, Ruswhandi RA, Giorgadze N, Pirtskhalava T, Podgorni O. Obesity-induced cellular senescence drives anxiety and impairs neurogenesis. *Cell metabolism*. 2019 May 7;29(5):1061-77.
3. Vimalaswaran KS, Berry DJ, Lu C, Tikkanen E, Pilz S, Hiraki LT, Cooper JD, Dastani Z, Li R, Houston DK, Wood AR. Causal relationship between obesity and vitamin D status: bidirectional Mendelian randomization analysis of multiple cohorts. *PLoS medicine*. 2013 Feb 5;10(2): e1001383.
4. Vranić L, Mikolašević I, Milić S. Vitamin D deficiency: consequence or cause of obesity? *Medicina*. 2019 Aug 28;55(9):541.
5. Laing ST, Smulevitz B, Vatcheva KP, Rahbar MH, Reininger B, McPherson DD, McCormick JB, Fisher-Hoch SP. Subclinical atherosclerosis and obesity phenotypes among Mexican Americans. *Journal of the American Heart Association*. 2015 Mar 18;4(3):e001540.
6. Legarth C, Grimm D, Krueger M, Infanger M, Wehland M. Potential beneficial effects of vitamin d in coronary artery disease. *Nutrients*. 2019 Dec 30;12(1):99.
7. Wimalawansa SJ. Vitamin D and cardiovascular diseases: Causality. *The Journal of steroid biochemistry and molecular biology*. 2018 Jan 1; 175:29-43.
8. Rai V, Agrawal DK. Role of vitamin D in cardiovascular diseases. *Endocrinology and Metabolism Clinics*. 2017 Dec 1;46(4):1039-59.
9. Ortega FB, Lee DC, Katzmarzyk PT, Ruiz JR, Sui X, Church TS, Blair SN. The intriguing metabolically healthy but obese phenotype: cardiovascular prognosis and role of fitness. *European heart journal*. 2013 Feb 1;34(5):389-97.
10. Siasos G, Tousoulis D, Oikonomou E, Maniatis K, Kioufis S, Zaromitidou M, Kokkou E, Mazaris S, Konsola T, Stefanadis C. Vitamin D serum levels are associated with cardiovascular outcome in coronary artery disease. *European Heart Journal*. 2013 Aug 1;34(suppl\_1): P2486.
11. Lai H, Fishman EK, Gerstenblith G, Brinker JA, Tong W, Bhatia S, Detrick B, Lai S. Vitamin D deficiency is associated with significant coronary stenoses in asymptomatic African American chronic cocaine users. *International journal of cardiology*. 2012 Jul 12;158(2):211-6.
12. Fernández-Alvira JM, Fuster V, Dorado B, Soberón N, Flores I, Gallardo M, Pocock S, Blasco MA, Andrés V. Short telomere load, telomere length, and subclinical atherosclerosis: the PESA study. *Journal of the American College of Cardiology*. 2016 May 31;67(21):2467-76.
13. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo Jr JL, Jones DW, Materson BJ, Oparil S, Wright Jr JT, Roccella EJ. Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *hypertension*. 2003 Dec 1;42(6):1206-52.
14. Mittal BV, Singh AK. Hypertension in the developing world: challenges and opportunities. *American Journal of Kidney Diseases*. 2010 Mar 1;55(3):590-8.
15. Hollick MF. Vitamin D: important for prevention of osteoporosis, cardiovascular heart disease, type 1 diabetes, autoimmune diseases, and some cancers. *Southern medical journal*. 2005 Oct 1;98(10):1024-8.
16. Colley RC, Garriguet D, Janssen I, Craig CL, Clarke J, Tremblay MS. Physical activity of Canadian adults: accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health reports*. 2011 Mar 1;22(1):7.
17. Warburton DE, Charlesworth S, Ivey A, Nettlefold L, Bredin SS. A systematic review of the evidence for Canada's Physical Activity Guidelines for Adults. *International journal of behavioral nutrition and physical activity*. 2010 Dec;7(1):1-220.
18. Elmer PJ, Obarzanek E, Vollmer WM, Simons-Morton D, Stevens VJ, Young DR, Lin PH, Champagne C, Harsha DW, Svetkey LP, Ard J. Effects of comprehensive lifestyle modification on diet, weight, physical fitness, and blood pressure control: 18-month results of a randomized trial. *Annals of internal medicine*. 2006 Apr 4;144(7):485-95.
19. Hosseinpanah F, Yarjanli M, Sheikholeslami F, Heibatollahi M, Eskandary PS, Azizi F. Associations between vitamin D and cardiovascular outcomes; Tehran Lipid and Glucose Study. *Atherosclerosis*. 2011 Sep 1;218(1):238-42.
20. Vitamin D. deficiency. Holick MF. *N Engl J Med*. 2007; 357:266-81.
21. Lucas RM, Ponsonby AL, Dear K, Valery PC, Taylor B, Van Der Mei I, McMichael AJ, Pender MP, Chapman C, Coulthard A, Kilpatrick TJ. Vitamin D status: multifactorial contribution of environment, genes and other factors in healthy Australian adults across a latitude gradient. *The Journal of steroid biochemistry and molecular biology*. 2013 Jul 1;136: 300-8.
22. Holick MF. Environmental factors that influence the cutaneous production of vitamin D. *The American journal of clinical nutrition*. 1995 Mar 1;61(3): S638-45.

23. Pludowski P, Holick MF, Pilz S, Wagner CL, Hollis BW, Grant WB, Shoenfeld Y, Lerchbaum E, Llewellyn DJ, Kienreich K, Soni M. Vitamin D effects on musculoskeletal health, immunity, autoimmunity, cardiovascular disease, cancer, fertility, pregnancy, dementia and mortality—a review of recent evidence. *Autoimmunity reviews*. 2013 Aug 1;12(10):976-89.
24. Amer M, Qayyum R. Relationship between 25-hydroxyvitamin D and all-cause and cardiovascular disease mortality. *The American journal of medicine*. 2013 Jun 1;126(6):509-14.
25. Zhao G, Ford eS, li C, Kris-etherton PM, etherton td, balluz lS. Independent associations of serum concentrations of 25-hydroxyvitamin d and parathyroid hormone with blood pressure among uS adults. *J hypertens*. 2010; 28:1821-8.
26. Forman JP, Giovannucci E, Holmes MD, Bischoff-Ferrari HA, Tworoger SS, Willett WC, Curhan GC. Plasma 25-hydroxyvitamin D levels and risk of incident hypertension. *Hypertension*. 2007 May 1;49(5):1063-9.
27. Scragg R, Sowers M, Bell C. Serum 25-hydroxyvitamin D, ethnicity, and blood pressure in the Third National Health and Nutrition Examination Survey. *American journal of hypertension*. 2007 Jul 1;20(7):713-9.
28. Richards JB, Valdes AM, Gardner JP, Paximadas D, Kimura M, Nessa A, Lu X, Surdulescu GL, Swaminathan R, Spector TD, Aviv A. Higher serum vitamin D concentrations are associated with longer leukocyte telomere length in women. *The American journal of clinical nutrition*. 2007 Nov 1;86(5):1420-5.
29. von Zglinicki T. Telomeres and replicative senescence: is it only length that counts? *Cancer letters*. 2001 Jul 26;168(2):111-6.
30. Von Zglinicki T. Oxidative stress shortens telomeres. *Trends in biochemical sciences*. 2002 Jul 1;27(7):339-44.
31. Hayflick L. The limited in vitro lifetime of human diploid cell strains. *Experimental cell research*. 1965 Mar 1;37(3):614-36.