

Analysis of Diagnostic Accuracy of Computed Tomography Scan in Mediastinal Masses: A Cross-sectional Study

Anupriya Mishra¹, Priyanka Kumari², Ashok Kumar³, Madhukar Dayal⁴

¹Senior Resident, Department of Radiology, Sri Krishna Medical College & Hospital, Muzaffarpur, Bihar.

²Senior Resident, Department of Radiology, Nalanda Medical College & Hospital, Patna, Bihar.

³Associate Professor and HOD, Department of Radiology, Sri Krishna Medical College & Hospital, Muzaffarpur, Bihar.

⁴Associate Professor, Department of Radiology, Nalanda Medical College & Hospital, Patna, Bihar.

Received: 05-01-2026 / Revised: 15-01-2026 / Accepted: 25-01-2026

Corresponding author: Dr. Priyanka Kumari

Conflict of interest: Nil

Abstract

Background: From benign and malignant tumors, cysts, vascular abnormalities, and lymph node masses to diffuse lesions including pneumomediastinum, mediastinitis, mediastinal fibrosis, and encysted pleural effusion, mediastinal diseases encompass a broad spectrum of pathology. The basic aim of the study is to analyse the diagnostic accuracy of computed tomography scan in mediastinal masses assuming histopathological findings as gold standard.

Methods: This cross sectional study was conducted in Department of Radiology, Sri Krishna Medical College and Hospital, Muzaffarpur, Bihar from July 2025 to December 2025. After taking informed consent, computed Tomography was performed on CT scanner before and after intravenous contrast administration.

Results: Age range in this study was from 25-65 years with mean age of 48.57 ± 10.53 years. Majority of the patients 58.78% were between 46-65 years of age as shown in Table 1. Out of these 148 patients, 82 (55.41%) were males and 66 (44.59%) were females with ratio of 1.2:1. Mean duration of disease was 8.39 ± 4.78 months (Table 2). Mean size of mass was 5.21 ± 2.45 cm.

Conclusion: It is concluded that computed tomography is a highly sensitive and accurate non-invasive modality for differentiating malignant and benign mediastinal masses, and has not only dramatically improved our ability of accurate diagnosis of mediastinal masses but also improved patient care by timely and proper treatment.

Key words: Mediastinal, Histopathology, Vascular anomalies, Lymph node.

DOI: 10.25258/ijcpr.18.2.171

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Mediastinal lesions include a wide spectrum of pathology, ranging from tumors (benign & malignant), cysts, vascular anomalies, and lymph node masses to diffuse lesions such as pneumomediastinum, mediastinitis, mediastinal fibrosis and encysted pleural effusion.[1] These lesions are challenging problems frequently faced by the radiologist. Mediastinal masses span a wide histopathological and radiological spectrum.

The most frequent lesions encountered in the mediastinum are thymoma, neurogenic tumours and benign cysts, altogether representing 60% of patients with mediastinal masses.[2] Neurogenic tumours, germ cell neoplasms and foregut cysts represent 80% of childhood lesions, whereas primary thymic neoplasms, thyroid masses and lymphomas are the most common in adults.[3]. In a

study, malignant mediastinal masses were found to be 44.0%.[4]Chest radiograph is the imaging modality in the initial diagnosis of mediastinal masses, however, it may appear normal in the presence of mediastinal masses. Posteroanterior (PA) and lateral radiograph of the chest for an unrelated cause are the usual ways in which an asymptomatic mediastinal mass is identified.[5]

Chest radiography obviously is the first study that would be performed in an individual with symptoms referable to the thorax. The PA view allows for determination of bilaterality and superior or inferior location, while the lateral chest radiograph determines the specific compartment. Mediastinal sonography is an effective imaging technique and is characterized by low cost and multiplanar images. It is used as an adjunctive

technique to other investigations. Further evaluation is done with Computed Tomography (CT) and Magnetic Resonance Imaging (MRI). Percutaneous image-guided core biopsy of mediastinal lesions is also indicated to reach a conclusion.[5]

To develop an accurate and cost-effective imaging plan, there are 4 primary goals when imaging mediastinal masses: (1) identification and accurate compartmental localization; (2) detailed mass description; (3) provision of an accurate and succinct differential diagnosis; (4) recommendation of a cost-effective imaging and patient management plan.[6]

Material and Methods

This cross sectional study was conducted in Department of Radiology, Sri Krishna Medical College and Hospital, Muzaffarpur, Bihar from July 2025 to December 2025.

All patients with chest x-ray showing mediastinal widening by soft tissue mass of any size and >3 months duration, Patients 25-65 years of age and

both genders were included in this study. Patients already operated for mediastinal mass, having biopsy proven report and with chronic liver disease (assessed on history and medical record (s/creatinine >1.1 mg/dl) were excluded in this study.

Total number of 148 patients who were admitted in other departments of SKMCH and referred by clinician to the radiology department fulfilling the inclusion/exclusion criteria was selected.

After taking informed consent, computed Tomography was performed on CT scanner before and after intravenous contrast administration. Each CT scan findings was interpreted by one consultant radiologist (at least 5 years of experience) and was looked for benign or malignant mediastinal mass as per-operational definition.

Collected data was analyzed through computer software SPSS 20.0. Mean and standard deviation were calculated for quantitative variables i.e. age, duration of disease and size of mass.

Results

Table 1: Distribution of patients according to Age

Age (in years)	No. of Patients	Percentage
25-45	61	41.22%
46-65	87	58.78%
Total	148	100.0%

Mean±SD=48.57±10.53 years

Age range in this study was from 25-65 years with mean age of 48.57 ± 10.53 years. Majority of the patients 58.78% were between 46-65 years of age as shown in (Table 1).

Table 2: Distribution of patients according to duration of disease

Duration of disease	No. of Patients	Percentage
4-12months	93	62.84%
>12months	55	37.16%
Total	148	100.0%

Mean±SD=8.39±4.78 months

Out of these 148 patients, 82 (55.41%) were males and 66 (44.59%) were females with ratio of 1.2:1. Mean duration of disease was 8.39 ± 4.78 months (Table 2).

Table 3: Distribution of patients according to size of mass

Size of mass(cm)	No. of Patients	Percentage
≤5	101	68.24%
>5	47	31.76%
Total	148	100.0%

Mean±SD=5.21±2.45 cm

Mean size of mass was 5.21 ± 2.45 cm (Table 3).

Table 4: Computed tomography and histopathology findings

	Positive result on Histopathology	Negative result on Histopathology	P-value
Positive result on CT	79(TP)*	04(FP)***	0.0001
Negative result on CT	05(FN)**	60(TN)****	

*-TP=True positive**-FP=False positive***-FN=False negative, ****-TN=True negative

All the patients were subjected to CT scanning of the thorax and CT scan supported the diagnosis of malignant mediastinal mass in 83 (56.08%) patients and benign mediastinal mass in 65 (43.92%) patients (Table 4).

Discussion

Computed tomography is the most important imaging modality in the evaluation of mediastinal mass. Characterisation on CT scan is based on specific attenuation of air, fat, water and calcium. High resolution multiplanar reformation images display the detailed anatomical relationship of the tumour with the adjacent structures.

Magnetic Resonance imaging has high contrast resolution and multiplanar capability, thereby providing additional information as to location of tumour and extent of intraspinal involvement and is a preferred modality in evaluating neurogenic tumours.[7] I have conducted this study to determine the diagnostic accuracy of computed tomography scan in differentiating malignant and benign mediastinal masses, taking histopathology as gold standard.[8]

Age range in my study was from 25-65 years with mean age of 48.57 ± 10.53 years. Out of these 148 patients, 82 (55.41%) were males and 66 (44.59%) were females with ratio of 1.2:1. All the patients were subjected to CT scanning of the thorax and CT scan supported the diagnosis of malignant mediastinal mass in 83 (56.08%) patients and benign mediastinal mass in 65 (43.92%) patients.[9] Histopathology findings confirmed malignant mediastinal mass in 84 (56.76%) patients and benign mediastinal mass in 64 (43.24%) patients. In CT positive patients, 79 (True Positive) had malignant mediastinal mass and 04 (False Positive) had benign mediastinal mass on histopathology.[10] Among 65, CT negative patients, 05 (False Negative) had malignant mediastinal mass on histopathology whereas 60 (True Negative) had benign mediastinal mass on histopathology ($P=0.0001$)[11]

Conclusion

It is concluded that computed tomography is a highly sensitive and accurate non-invasive modality for differentiating malignant and benign mediastinal masses, and has not only dramatically improved our ability of accurate diagnosis of mediastinal masses but also improved patient care by timely and proper treatment.

References

1. Mushtaq N, Alam MM, Aslam S, Fadoo Z, Anwar-ul-Haq. Malignant mediastinal mass in children: a single institutional experience from a developing country. *J Pak Med Assoc.* 2014; 64: 386-389.
2. Yokoyama Y, Chen F, Date H. Surgical resection of a giant mediastinal teratoma occupying the entire left hemithorax. *Gen Thorac Cardiovasc Surg.* 2014; 62(4): 255-257.
3. Aroor AR, Prakasha SR, Seshadri SST, Raghuraj U. A study of clinical characteristics of mediastinal mass. *J Clin Diagn Res.* 2014; 8: 77-80.
4. Dutta P, Patel KKR, Choudhury PR. Mediastinal masses: a radiological study of 50 cases. *Intl J Biomed Res.* 2014; 5(7): 468-473.
5. Tomiyama N, Honda O, Tsubamoto M, Inoue A, Sumikawa H, Kuriyama K, et al. Anterior mediastinal tumors: diagnostic accuracy of CT and MRI. *Eur J Radiol.* 2009; 69: 280-288.
6. Thacker PG, Mahani MG, Heider WA, Lee EY. Imaging evaluation of mediastinal masses in children and adults practical diagnostic approach based on a new classification system. *J Thorac Imaging.* 2015; 30(4): 247-264.
7. Kalhan S, Sharma P, Sharma S, Dudani S, Ramakrishnan Ts, Chowdhry A. Evaluation of precision of guidance techniques in image guided fine needle aspiration cytology of thoracic mass lesions. *J Cytol.* 2012; 29: 6-10.
8. Kumar P, Kumar B, Thakur A. Echogenic mobile spherules in a cyst as an ultrasound finding in cystic mediastinal teratoma. *Indian J Surg.* 2013; 75 (Suppl 1): 385-387.
9. Li H, Wang DL, Liu XW, Geng ZJ, Xie CM. Computed tomography characterization of neuroendocrine tumors of the thymus can aid identification and treatment. *Acta Radiol.* 2013; 54(2): 175-180.
10. Kaur H, Tiwari P, Dugg P, Ghuman J, Shivhare P, Mehmi RS. Computed tomographic evaluation of mediastinal masses/lesions with contrast enhancement and correlation with pathological diagnosis-A study of 120 cases. *J Biomed Graphics Comput.* 2014; 4(3): 28-35.
11. Rashid N, Ahmad KK, Ali W. Diagnostic role of computed tomography in evaluation of mediastinal masses. *Ann King Edward Med Uni.* 2011; 17: 410-415.