

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

Shreya Verma¹, Vaishnavi Srivastava², Prashant Kumar Singh³, Ajay Kumar Gupta⁴,
Manishi Tripathi^{1*}

¹Department of Microbiology, School of Life Sciences and Biotechnology, Chhatrapati Shahu Ji Maharaj University, Kanpur, UP, India

²Department of Biotechnology, School of Life Sciences and Biotechnology, Chhatrapati Shahu Ji Maharaj University, Kanpur, UP, India

³Department of Pharmacology, Institute of Pharmaceutical Sciences, University of Lucknow, UP, India

⁴Department of Pharmacology, School of Pharmaceutical Sciences, Chhatrapati Shahu Ji Maharaj University, Kanpur, UP, India

*Corresponding Author: Dr. Manishi Tripathi, Department of Microbiology, School of Life Sciences and Biotechnology, Chhatrapati Shahu Ji Maharaj University, Kanpur, UP, India.

Email: manishitripathi@csjmu.ac.in

Abstract

Siderophores are metal-binding compounds that are of low molecular weight, high affinity, produced by living organisms to obtain iron under limiting environmental conditions. Iron plays a critical role in respiration, enzymatic processes, synthesis of DNA and redox balance but the bioavailability of iron is severely limited in aerobic conditions because it forms insoluble ferric complex. To mitigate this drawback, microorganisms, plants and even mammals synthesize siderophores which bind to the ferric ions and deliver them to the cell via special receptor-mediated processes. In addition to the iron-acquiring properties, siderophores are also involved in heavy metal detoxification, interspecies competition, pathogenesis, and various industrial and biomedical uses. Their spectacular structural diversity mostly the catecholates, hydroxamates, and carboxylates control their metal specificity, stability constants and transport route. The review describes the variety of siderophores in terms of chemistry and biological synthesis, the key approaches to their production and extraction, and their growing role in agriculture, the environment, medicine and biotechnology.

Keywords: Siderophores, Iron acquisition, Metal-binding compounds, Heavy metal detoxification, Ferric ion transport.

How to cite this article: Verma S, Srivastava V, Singh PK, Gupta AK, Tripathi M. Exploring Siderophore Diversity: Production, Extraction and Functional Applications. *Int J Drug Deliv Technol.* 2026;16(12s): 486-499. DOI: 10.25258/ijddt.16.12s.60

1. Introduction

Iron is one of the most indispensable micronutrients required for the survival and functioning of almost all living organism.¹ At the cellular level, iron serves as a critical cofactor in a wide range of biochemical and physiological processes, including cellular respiration, oxidative phosphorylation, DNA synthesis, nitrogen fixation, oxygen transport, and numerous enzymatic redox reactions. Iron-containing proteins such as cytochromes, ferredoxins, catalases, peroxidases, and hemoproteins are central to energy metabolism and maintenance of cellular homeostasis.² In microorganisms, iron plays an especially vital role in ATP generation through electron transport chains, regulation of gene expression, detoxification of reactive oxygen species, and biosynthesis of essential metabolites.³ Because of these diverse roles, iron availability directly influences microbial growth rate, metabolic efficiency, and ecological fitness.

Despite its abundance in the Earth's crust, iron presents a major bioavailability challenge in natural environments. Under aerobic conditions and neutral pH, ferrous iron (Fe²⁺) is rapidly oxidized to ferric iron (Fe³⁺), which forms highly insoluble ferric oxyhydroxides and polymeric complexes.⁴ These precipitated forms drastically reduce the concentration of soluble iron in soil, freshwater, marine systems, and host tissues to levels far below the physiological requirements of living cells. This contradiction where iron is plentiful but biologically inaccessible is often referred to as the "iron paradox".⁵ Consequently, microorganisms growing in such environments experience persistent iron limitation, which acts as a strong evolutionary pressure driving the development of sophisticated iron acquisition mechanisms.

To overcome iron scarcity, microorganisms have evolved multiple adaptive strategies, among which siderophore production represents the most efficient,

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

widespread, and evolutionarily conserved system. Siderophores are low molecular-weight (generally <10 kDa), high-affinity iron-chelating molecules synthesized intracellularly and secreted into the extracellular environment under iron-deficient conditions.⁶ These specialized metabolites exhibit extraordinary specificity for ferric ions and form thermodynamically stable Fe^{3+} -siderophore complexes through multidentate coordination. The stability constants of these complexes are among the highest known for natural metal chelators, allowing siderophores to effectively compete with environmental ligands and host iron-binding proteins.⁷ Once secreted, siderophores scavenge ferric iron from mineral surfaces, organic matter, and host iron-binding proteins such as transferrin, lactoferrin, ferritin, and heme-containing molecules. The ferric iron siderophore complex is then recognized by highly specific receptor proteins located on the microbial cell surface.⁸ Transport across the outer membrane occurs through energy-dependent systems such as TonB-dependent transporters in Gram-negative bacteria, followed by periplasmic shuttling and cytoplasmic import via ATP-binding cassette (ABC) transporters.⁹ After internalization, iron is released through reductive mechanisms or enzymatic degradation of the siderophore ligand, making it available for metabolic use.¹⁰ Importantly, siderophore biosynthesis and uptake are tightly regulated processes controlled by intracellular iron levels to prevent metal toxicity and maintain homeostasis.

A wide diversity of microorganisms are capable of siderophore production. Numerous bacterial genera including *Pseudomonas*, *Azotobacter*, *Bacillus*, *Enterobacter*, *Serratia*, *Azospirillum*, and *Rhizobium* are known to synthesize significant quantities of siderophores when exposed to iron-starved environments.¹¹ The genes encoding enzymes for siderophore biosynthesis, receptor proteins, and transport systems are frequently organized in coordinated gene clusters or operons, enabling synchronized regulation.¹² Once adequate intracellular iron levels are restored, regulatory proteins repress further siderophore synthesis to conserve metabolic energy. Interestingly, certain microorganisms possess the ability to utilize siderophores produced by other species, a phenomenon termed “xenosiderophore utilization,” which significantly influences microbial competition, cooperation, and ecological dynamics within complex communities.¹³

Beyond their central role in iron acquisition, siderophores exhibit remarkable versatility in metal

coordination chemistry. Many siderophores are capable of interacting with other biologically relevant and toxic metal ions, including zinc, copper, nickel, cobalt, cadmium, lead, and aluminum. This multi-metal binding capacity expands their functional significance beyond nutrition to include heavy metal detoxification, environmental metal cycling, and bioremediation of contaminated ecosystems.¹⁴ In soil environments, siderophores contribute to mineral weathering, nutrient mobilization, and rhizosphere interactions that support plant growth. In aquatic systems, they influence trace metal distribution and microbial productivity.¹⁵ Siderophore systems are also closely linked with microbial pathogenicity and host-pathogen interactions. In vertebrate hosts, free iron is extremely limited due to tight sequestration by high-affinity host proteins, creating an iron-restricted environment that suppresses microbial proliferation.¹⁶ Many pathogenic bacteria overcome this defense strategy by secreting siderophores that can extract iron directly from host proteins, thereby enhancing survival, colonization, and virulence. The ability to produce high-affinity siderophores is therefore considered a critical virulence determinant in several clinically important pathogens.

Extensive research over the past decades has revealed enormous structural diversity among siderophores. To date, nearly five hundred distinct siderophores have been identified across bacteria, fungi, cyanobacteria, plants, and even mammalian systems.¹⁷ Based on the functional groups responsible for metal chelation, siderophores are broadly classified into three principal categories: catecholates (phenolates), hydroxamates, and carboxylates. Differences in ligand chemistry, denticity, stereochemistry, and molecular architecture determine their metal-binding affinity, thermodynamic stability, transport behavior, and ecological specialization. The scientific importance of siderophores extends far beyond microbial physiology. Their unique chemical properties and biological functions have attracted considerable attention in agriculture, environmental biotechnology, industrial microbiology, medicine, and materials science. Applications range from plant growth promotion and biocontrol of phytopathogens to heavy metal remediation, radioactive waste management, iron chelation therapy, antimicrobial drug delivery, and development of bioengineered materials.¹⁸ As research advances, siderophores are increasingly recognized as multifunctional biomolecules with significant translational potential. Given their biochemical sophistication, ecological relevance, and expanding

technological applications, a comprehensive understanding of siderophore diversity is essential. Therefore, the present article aims to provide a comprehensive overview of siderophore diversity, encompassing their production and extraction strategies, structural and functional characteristics, and diverse applications in agriculture, environment, medicine, and biotechnology, with particular emphasis on their potential role in the management of iron deficiency conditions and related therapeutic innovations.

2. Methodology: Siderophores, with emphasis on their diversity, production, extraction methods, functional applications in pharmaceutical and biotechnological contexts along with other relevant keywords were identified using major scientific databases, including PubMed, Scopus, Web of Science, and Google Scholar, covering publications from both classical studies and recent advances. The selection criteria included peer-reviewed articles, patents, and book chapters that addressed siderophore chemistry, biosynthesis, structural diversity, and application in agriculture, environment, medicine, and industry. Studies focusing on pharmaceutical relevance, such as antimicrobial resistance, drug delivery, and therapeutic innovation, were prioritized. Data were extracted, categorized, and synthesized to highlight trends in siderophore production, extraction techniques, and functional roles across different organisms. The information was critically analysed to identify gaps in current knowledge, emerging applications, and potential directions for future research in biological sciences.

3. Siderophores and iron acquisition

Siderophores chelate iron through highly specific coordination chemistry that enables them to solubilize and transport ferric ions under iron-limiting conditions. These low molecular weight compounds possess functional groups such as catechol, hydroxamate, or carboxylate moieties, which form stable hexadentate complexes with Fe^{3+} . The strong affinity of siderophores for ferric ions outcompetes other ligands in the environment, thereby mobilizing otherwise insoluble iron. Once bound, the siderophore-iron complex is recognized by dedicated receptor proteins on microbial cell surfaces, facilitating active transport across membranes. Inside the cell, reductases or hydrolytic enzymes release Fe^{2+} for incorporation into essential metabolic pathways, including respiration, DNA synthesis, and enzymatic catalysis. This mechanism not only ensures iron homeostasis but also

contributes to microbial survival, pathogenicity, and ecological competitiveness.⁶⁵




Figure 1: Mechanism of iron chelation by Siderophore

4. Chemistry and Structural Diversity of Siderophores

The remarkable efficiency of siderophores in iron sequestration is primarily attributed to their highly specialized chemical architecture. These molecules are typically low-molecular-weight organic ligands designed to coordinate ferric iron through multiple electron-donating functional groups arranged in precise three-dimensional configurations. Iron chelation occurs via coordinate covalent bonding, where lone pairs of electrons from oxygen and nitrogen atoms bind to the empty d-orbitals of Fe^{3+} .²⁰ Most siderophores function as **hexadentate ligands**, meaning six donor atoms coordinate a single ferric ion to form a highly stable octahedral complex. This multidentate binding greatly enhances thermodynamic stability, prevents iron precipitation, and minimizes displacement by competing environmental ligands.

The strength of metal binding depends on several structural parameters, including the nature of functional groups, denticity, stereochemistry, ligand flexibility, molecular backbone composition, and electronic properties. The stability constants of Fe^{3+} -siderophore complexes are among the highest observed in biological systems and often exceed those of host iron-binding proteins.²¹ Such extraordinary affinity enables microorganisms to successfully compete for iron even in extremely iron-limited environments such as marine ecosystems, alkaline soils, and vertebrate host tissues. Beyond ferric iron, many siderophores can coordinate other transition metals, reflecting their adaptable ligand chemistry and broader environmental relevance.

4.1 Functional Groups Involved in Metal Chelation

Siderophore chelation chemistry is governed by oxygen- and nitrogen-containing functional groups capable of donating electron pairs to metal centres. The principal chelating moieties include **catecholate hydroxyls**, **hydroxamate groups**, and **carboxylate/hydroxycarboxylate ligands**. These groups differ in electron density, acidity, coordination

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

geometry, and metal selectivity, which ultimately determine siderophore classification and function.

Catecholate groups contain adjacent phenolic hydroxyl substituents on an aromatic benzene ring. Upon deprotonation, these oxygen atoms form strong bidentate coordination bonds with Fe^{3+} . The aromatic system provides resonance stabilization and electron delocalization, significantly strengthening complex formation.²¹

Hydroxamate groups consist of the functional unit $C(=O)N-OH$, where both the carbonyl oxygen and hydroxylamine oxygen participate in metal coordination.²² These groups typically form stable five-membered chelate rings with ferric ions and are widely distributed in microbial siderophores.

Carboxylate siderophores utilize carboxyl and hydroxycarboxyl functional groups that coordinate iron via oxygen atoms. These ligands are often arranged within amino polycarboxylic acid frameworks that provide flexible coordination environments.²³

4.2 Major Structural Classes of Siderophores

Based on dominant chelating groups, siderophores are broadly categorized into **catecholate**, **hydroxamate**, and **carboxylate** types.²⁴ This classification reflects differences in coordination chemistry, biosynthetic origin, ecological adaptation, and taxonomic distribution.

Catecholate (Phenolate) Siderophores

Catecholate siderophores are characterized by ortho-dihydroxybenzene rings that act as powerful oxygen donors for ferric iron coordination.²⁵ Each catechol unit functions as a bidentate ligand; thus, three catechol groups typically form a hexadentate octahedral complex with one Fe^{3+} ion. These siderophores exhibit exceptionally high stability constants and are regarded as the strongest natural iron chelators. Structurally, catecholate siderophores may exist in cyclic macrostructures or linear forms depending on their peptide or ester backbone organization. Their rigid aromatic framework enhances binding strength and resistance to hydrolytic degradation.²⁶

Hydroxamate Siderophores

Hydroxamate siderophores are among the most widely distributed types in bacteria and fungi. Their defining feature is the hydroxamate functional group, where adjacent oxygen atoms coordinate ferric iron. Three hydroxamate units combine to form stable hexadentate octahedral complexes. Compared with catecholates, hydroxamates show slightly lower—yet still very high—binding affinities.²⁶ Their structures often include cyclic depsipeptides or linear peptide chains

that provide conformational flexibility, supporting roles in extracellular iron scavenging, intracellular transport, and iron storage.

Carboxylate (Complexone) Siderophores

Carboxylate siderophores employ carboxyl and hydroxycarboxyl groups for iron coordination.²⁷ These molecules usually possess amino polycarboxylic acid skeletons that create adaptable metal-binding environments. Unlike catecholates and hydroxamates, carboxylate siderophores frequently form mixed ligand systems and may bind multiple metals with moderate affinity. Their structural plasticity allows interaction with a wider range of metal ions, contributing to mineral weathering, nutrient mobilization, and soil metal dynamics.

4.3 Mixed-Type Siderophores

Many siderophores do not belong exclusively to a single structural class. Several organisms synthesize **mixed-type siderophores** that incorporate combinations of catecholate, hydroxamate, and carboxylate groups within the same molecule.²⁸ This hybrid architecture enhances metal-binding versatility and allows coordination under diverse environmental conditions. Mixed-type siderophores often display intermediate stability constants and improved adaptability to fluctuations in pH, redox potential, and metal availability.²⁶ Their structural complexity reflects evolutionary optimization for survival in competitive ecological niches.

Table 1. Major Classes of Siderophores Based on Chemical Structure

Class	Functional Group	Coordination Mode	Key Features	Typical Producers
Catecholates (Phenolates)	Ortho-dihydroxybenzene groups	Hexadentate via oxygen donors	Extremely high Fe^{3+} affinity, often cyclic/linear peptides	Mostly Gram-negative bacteria
Hydroxamates	$C(=O)N-(OH)R$ group	Hexadentate via oxygen	Chemically stable, strong	Bacteria and fungi

		n donors	g visible absorbance, widespread distribution	
Carboxylates	Carboxyl & hydroxycarboxyl groups	Mixed ligand coordination	Broad metal binding spectrum, high structural diversity	Soil microbes, plants
Mixed-type	Combination of above groups	Multi-mode chelation	Enhanced flexibility and metal specificity	Diverse microorganisms

4.4 Structure–Function Relationship

The biological performance of siderophores is directly governed by their molecular architecture. Ligand denticity determines the number of coordination bonds and strongly influences thermodynamic stability. Molecular rigidity enhances binding strength, whereas flexibility facilitates receptor recognition and membrane transport.²⁹ Stereochemical configuration affects receptor specificity and enzymatic processing. Hydrophilicity, molecular size, and charge distribution influence diffusion through extracellular matrices and transport across membranes. Structural diversity also dictates ecological specialization: high-affinity catecholates dominate iron-deficient niches, hydroxamates support balanced acquisition and storage, and carboxylates contribute to broader metal mobilization in soil ecosystems.³⁰ Mixed-type siderophores provide adaptive advantages in fluctuating environments.

5. Classification Based on Producing Organisms

Siderophore diversity is not only reflected in chemical structure but also in the wide range of biological systems that synthesize them. Different groups of organisms have evolved distinct siderophore systems tailored to their ecological niches, physiological requirements, and environmental iron availability. These organism-specific variations influence siderophore chemistry, transport mechanisms, regulatory pathways, and functional roles in survival, competition, and symbiosis.³¹ Based on biological origin, siderophores are broadly classified into bacterial, fungal, cyanobacterial, mammalian, and plant siderophores.

5.1 Bacterial Siderophores

Bacteria represent the most extensively studied and diverse producers of siderophores. Both Gram-negative and Gram-positive bacteria synthesize extracellular siderophores to overcome iron limitation in soil, aquatic systems, and host-associated environments. Bacterial siderophores are secreted into the surroundings where they chelate ferric ions and form soluble Fe³⁺-siderophore complexes that are transported back into the cell via highly specific receptor-mediated systems.³² Gram-negative bacteria typically employ TonB-dependent outer membrane receptors coupled with periplasmic transport proteins and ATP-binding cassette (ABC) transporters for iron internalization.³⁵ Gram-positive bacteria, lacking an outer membrane, utilize membrane-anchored binding proteins and permeases for siderophore uptake. Many bacterial species possess multiple siderophore systems with different iron affinities, enabling flexible adaptation to fluctuating iron conditions.^{21;33}

Several free-living soil and rhizosphere bacteria produce siderophores that enhance ecological competitiveness. In contrast, pathogenic bacteria rely heavily on high-affinity siderophores to extract iron from host proteins such as transferrin, lactoferrin, ferritin, and haemoglobin.³⁴ This iron-scavenging ability is considered a major virulence determinant and contributes significantly to infection establishment and persistence.

5.2 Fungal Siderophores

Fungi synthesize a wide array of siderophores primarily belonging to hydroxamate and carboxylate classes. These molecules support both extracellular iron acquisition and intracellular iron storage, ensuring tight regulation of iron homeostasis across different developmental stages.⁶² Fungal siderophores are often categorized into extracellular siderophores, which capture iron from the environment, and intracellular

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

siderophores, which store and distribute iron within hyphae and spores. Saprophytic fungi play an essential ecological role by recycling nutrients in soil and decomposing organic matter, processes that depend heavily on siderophore-mediated iron mobilization.¹ Pathogenic fungi, on the other hand, utilize siderophores to survive within iron-restricted host environments and to resist oxidative stress imposed by immune defenses. The coordinated regulation of siderophore biosynthesis and transport enables fungi to balance iron acquisition with protection against iron-induced toxicity.⁶⁴

5.3 Cyanobacterial Siderophores

Cyanobacteria, the photosynthetic prokaryotes inhabiting marine and freshwater ecosystems, produce specialized siderophores adapted to aquatic environments where dissolved iron concentrations are extremely low.⁶¹ These siderophores are predominantly hydroxamate-type and are optimized for efficient iron solubilization and uptake in oxic water systems.

Iron acquisition in cyanobacteria is mediated by sophisticated ATP-binding cassette (ABC) transporter systems involving periplasmic iron-binding proteins, membrane permeases, and ATPase components.⁵⁹ These transporters enable high-affinity uptake of ferric iron complexes under nutrient-limited conditions. In addition to iron, cyanobacterial siderophores can chelate multiple heavy metals, contributing to metal detoxification and biogeochemical cycling. In agricultural ecosystems, particularly paddy fields and freshwater bodies, cyanobacterial siderophores assist in reducing heavy metal toxicity and improving nutrient availability, thereby supporting crop productivity and environmental sustainability.⁶⁰

5.4 Mammalian Siderophores

Recent discoveries have revealed the presence of endogenous siderophore-like molecules in mammalian systems, challenging the long-held belief that siderophores are exclusively microbial products. Mammalian siderophores participate in intracellular iron trafficking, mitochondrial iron delivery, and regulation of iron homeostasis.⁵⁸ These molecules function in coordination with iron-binding proteins and transport pathways to ensure controlled distribution of iron while preventing oxidative damage caused by free iron. Disruption of mammalian siderophore pathways has been associated with iron overload disorders, oxidative stress, mitochondrial dysfunction, and metabolic abnormalities.⁵⁷ Their identification has opened new perspectives in understanding iron metabolism and related disease mechanisms.

5.5 Plant Siderophores (Phytosiderophores)

Graminaceous plants synthesize and secrete phytosiderophores into the rhizosphere to mobilize iron from insoluble soil minerals. These compounds belong primarily to the mugineic acid family and form highly stable hexadentate complexes with ferric ions. The Fe-phytosiderophore complexes are subsequently transported into root cells via specialized membrane transporters.

Phytosiderophore secretion is strongly induced under iron deficiency and is particularly crucial in calcareous and alkaline soils where iron solubility is extremely low.⁵⁶ By enhancing iron acquisition efficiency, phytosiderophores support plant growth, chlorophyll synthesis, and overall crop productivity. In addition to iron mobilization, they influence rhizosphere microbial interactions and nutrient dynamics.

Table 2. Siderophore Classification Based on Producing Organisms

Producing Group	Major Siderophore Types	Primary Function	Ecological/Clinical Significance
Bacteria	Catecholates, Hydroxamates, Mixed	Extracellular iron scavenging	Survival in iron-limited niches, virulence in pathogens
Fungi	Hydroxamates, Carboxylates	Iron uptake and intracellular storage	Nutrient recycling, host colonization
Cyanobacteria	Predominantly Hydroxamates	Aquatic iron acquisition	Metal detoxification, nutrient cycling
Mammals	Endogenous siderophore-like molecules	Intracellular iron trafficking	Iron homeostasis, metabolic regulation
Plants (Grasses)	Phytosiderophores (mugineic acid family)	Rhizosphere iron solubilization	Essential for growth in alkaline soils

6. Production and Extraction of Siderophores

Siderophore production is a tightly regulated physiological response triggered primarily by iron-deficient conditions. When intracellular iron levels fall below a critical threshold, microorganisms activate specific gene clusters responsible for siderophore

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

biosynthesis, secretion, and uptake.³⁶ This regulation is commonly mediated by iron-responsive transcriptional regulators that repress siderophore-related genes under iron-sufficient conditions and derepress them during iron starvation. The efficiency of siderophore production therefore depends not only on the genetic potential of the organism but also on environmental and nutritional factors that influence metabolic activity.³⁷

Several physicochemical parameters significantly affect siderophore yield. Among these, iron concentration is the most critical factor, as even trace amounts of iron in culture media can suppress biosynthesis. Optimal production is typically achieved in iron-limited synthetic or semi-synthetic media. Carbon sources such as glucose, sucrose, glycerol, and organic acids influence metabolic flux and precursor availability for siderophore assembly.³⁸ Nitrogen sources, including ammonium salts, nitrates, amino acids, and peptones, also modulate production by affecting cellular growth rate and enzyme synthesis. Environmental conditions such as pH, temperature, aeration, agitation speed, and incubation time further determine productivity. Most bacterial siderophores are optimally produced under neutral to slightly alkaline pH and mesophilic temperature ranges, although extremophiles exhibit distinct preferences. Adequate aeration enhances oxidative metabolism and supports synthesis of iron-chelating ligands, particularly in aerobic microorganisms.³⁹ Incubation time is equally important, as siderophore secretion often correlates with the late exponential or early stationary growth phase when nutrient limitation intensifies.

Table 3. Factors Affecting Siderophore Production

Factor	Influence on Production
Iron concentration	Low iron strongly induces biosynthesis
Carbon source	Alters metabolic flux and precursor supply
Nitrogen source	Affects growth rate and enzyme synthesis
pH	Influences solubility and metabolic activity
Temperature	Controls enzymatic reaction rates

Aeration & agitation	Enhances aerobic metabolism and secretion
Incubation time	Peak production near stationary phase
Microbial strain	Genetic capacity determines yield

Strain selection plays a decisive role in maximizing siderophore yield. Wild-type strains isolated from iron-poor environments such as rhizosphere soils, marine habitats, and mining sites often exhibit superior production capacity. Advances in microbial biotechnology have enabled strain improvement through mutagenesis, adaptive evolution, and recombinant DNA approaches to enhance biosynthetic efficiency.⁴⁰ Fermentation strategies—including batch, fed-batch, and continuous culture systems—are employed at laboratory and industrial scales to optimize production kinetics and metabolite recovery. Following production, siderophores must be efficiently extracted and purified from culture media for characterization and application. The choice of extraction method depends on the chemical nature, polarity, stability, and molecular weight of the siderophore molecule. Initial recovery typically involves removal of microbial biomass by centrifugation or filtration to obtain cell-free supernatant containing extracellular siderophores.




Figure 2. Genetic regulation of siderophore biosynthesis under iron starvation

Solvent extraction is widely used for preliminary isolation. Organic solvents such as ethyl acetate, chloroform, methanol, and butanol are employed based on siderophore polarity and solubility characteristics. Liquid-liquid extraction enables partitioning of siderophores into solvent phases, followed by solvent evaporation to obtain crude extracts.⁴¹ This method is particularly useful for nonpolar or moderately polar siderophores. pH-mediated precipitation techniques exploit the acid-base properties of siderophores. Adjusting the pH of culture supernatant alters

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

ionization states, leading to selective precipitation of target compounds. Similarly, salt-induced precipitation using ammonium sulfate or other salts reduces solubility through ionic strength manipulation, facilitating recovery of siderophore fractions.

Chromatographic purification provides high-resolution separation and is essential for analytical and pharmaceutical applications. Ion-exchange chromatography separates siderophores based on charge differences, while gel filtration chromatography enables size-based fractionation. High-performance liquid chromatography (HPLC) is the most widely used technique for purification and quantitative analysis due to its high sensitivity, reproducibility, and compatibility with diverse detection systems.⁴² Reverse-phase HPLC is particularly effective for separating structurally related siderophores.

Advanced analytical tools such as mass spectrometry, nuclear magnetic resonance spectroscopy, and UV-visible spectrophotometry are subsequently employed for structural characterization and purity assessment.⁴³

The choice of purification workflow depends on downstream applications, required purity level, cost considerations, and scalability. Efficient production and extraction strategies are fundamental for translating siderophore research into agricultural, environmental, and biomedical applications. Optimization of fermentation conditions combined with robust purification protocols ensures consistent yield, structural integrity, and functional performance of siderophore molecules.




Figure 3: Major biosynthetic pathways involved in siderophore synthesis

Table 4. Major Extraction and Purification Techniques

Method	Principle	Advantages	Limitations
Solvent extraction	Polarity-based partitioning	Simple, cost-effective	Lower purity

pH precipitation	Acid–base solubility shift	Easy recovery	Not universal
Salt precipitation	Ionic strength manipulation	Good for bulk recovery	Co-precipitation impurities
Ion-exchange chromatography	Charge-based separation	High selectivity	Requires optimization
Gel filtration chromatography	Size-based separation	Gentle on molecules	Lower resolution
HPLC (Reverse phase)	Hydrophobic interaction	High purity & quantification	Expensive instrumentation

7. Functional Applications of Siderophores

The exceptional metal-chelating capacity and biological versatility of siderophores have enabled their application across multiple scientific and technological domains.⁴⁴ Originally studied for their role in microbial iron acquisition, siderophores are now recognized as multifunctional biomolecules with significant relevance in agriculture, environmental management, medicine, and industrial biotechnology.⁴⁵ Their high affinity for ferric iron, ability to interact with diverse metals, ecological compatibility, and biodegradability make them valuable tools in sustainable and translational research.

7.1 Agricultural Applications

Siderophore-producing microorganisms play a pivotal role in sustainable agriculture by improving nutrient availability and promoting plant growth. In iron-deficient soils, particularly calcareous and alkaline soils, ferric iron remains insoluble and inaccessible to plant roots. Microbial siderophores chelate ferric ions from soil minerals and organic complexes, converting them into soluble forms that can be absorbed by plants either directly or through microbe–plant interactions in the rhizosphere.⁴⁶ Plant growth–promoting rhizobacteria (PGPR) that produce siderophores enhance root development, chlorophyll synthesis, photosynthetic efficiency, and overall biomass accumulation. In addition to iron mobilization, siderophore-producing microbes often exhibit complementary traits such as phosphate solubilization,

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

nitrogen fixation, and phytohormone production, collectively improving soil fertility and crop productivity.⁴⁷ Siderophores also contribute to soil detoxification by immobilizing toxic heavy metals, thereby reducing metal uptake by plants and preventing phytotoxic effects. Their integration into biofertilizer formulations supports eco-friendly farming practices and reduces dependency on synthetic agrochemicals.

7.2 Biocontrol Activity

Siderophores are critical mediators of biological control in agricultural ecosystems. Beneficial rhizobacteria suppress phytopathogenic microorganisms through competitive iron sequestration.⁴⁸ Because iron is essential for pathogen survival and virulence, siderophore-mediated iron deprivation creates nutrient-limited conditions that inhibit pathogen growth, spore germination, and infection capability. This mechanism of competition-based antagonism reduces the incidence of soil-borne diseases and enhances crop resilience. Experimental evidence demonstrates that microbial mutants deficient in siderophore production exhibit significantly reduced biocontrol efficiency, underscoring the ecological importance of siderophore systems.⁴⁹ As a result, siderophore-producing microbes are widely explored as biopesticide agents and components of integrated disease management strategies.

7.3 Environmental Applications

The strong metal-binding affinity of siderophores extends beyond iron to numerous environmentally relevant heavy metals. Siderophores can chelate toxic elements such as cadmium, lead, mercury, chromium, nickel, copper, and arsenic, forming stable complexes that alter metal mobility and bioavailability. This property enables their application in bioremediation of contaminated soils and aquatic systems. Siderophore-producing microorganisms enhance removal or stabilization of heavy metals, contributing to detoxification of industrial waste sites, mining areas, and polluted farmlands.⁵⁰ In wastewater treatment systems, siderophores assist in metal recovery and purification processes. Their metal-chelating capacity also supports radioactive waste management and regulation of trace metal distribution in ecosystems. By influencing metal speciation and geochemical cycling, siderophores play a significant role in maintaining environmental health and sustainability.

7.4 Medical Applications

Siderophores have attracted considerable interest in biomedical research due to their high specificity for iron and compatibility with physiological systems. Iron overload disorders—commonly arising from repeated

blood transfusions, genetic conditions, or metabolic dysfunction—lead to accumulation of excess iron in vital organs such as the liver, heart, and endocrine tissues.⁵¹ Siderophore-based iron chelators are used therapeutically to bind excess iron and facilitate its excretion, thereby preventing tissue damage and organ failure.

Certain siderophores also demonstrate affinity for aluminum and other metals, enabling their use in diagnosis and treatment of metal toxicity. Their biocompatibility and strong metal-binding properties make them promising candidates for development of targeted chelation therapies. In addition, siderophores are being explored as drug delivery vehicles in antimicrobial therapy. Conjugation of antibiotics with siderophores enables “Trojan horse” strategies where pathogens actively transport drug-siderophore complexes into their cells, enhancing therapeutic efficiency and overcoming drug resistance mechanisms.⁵⁴

7.5 Role in Infection and Pathogenesis

Iron acquisition is a decisive factor in microbial pathogenicity. Within vertebrate hosts, free iron is tightly sequestered by high-affinity proteins such as transferrin, lactoferrin, ferritin, and hemoglobin, creating an iron-restricted environment that limits microbial proliferation.⁵² Pathogenic microorganisms counteract this defense by synthesizing high-affinity siderophores capable of extracting iron directly from host proteins. Siderophore-mediated iron uptake enhances pathogen survival, colonization efficiency, biofilm formation, and virulence. The presence of multiple siderophore systems in pathogens further increases their adaptability within host tissues.⁵³ In response, host organisms produce siderophore-binding proteins that neutralize microbial siderophores and restrict iron availability, forming a dynamic host-pathogen iron competition.⁵⁵ Understanding siderophore-mediated host-pathogen interactions is crucial for development of novel antimicrobial strategies, vaccines, and siderophore-targeted therapeutics.

Table 5. Functional Applications of Siderophores

Application Area	Functional Role	Practical Outcomes
Agriculture	Iron mobilization, PGPR support	Improved plant growth & yield
Biocontrol	Pathogen iron deprivation	Reduced crop diseases
Environmental remediation	Heavy metal chelation	Soil & water detoxification

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

Waste management	Metal recovery & stabilization	Safer waste processing
Medicine	Iron chelation therapy	Treatment of iron overload
Drug delivery	Trojan-horse antibiotic transport	Enhanced antimicrobial efficacy
Pathogenesis research	Host-microbe iron competition	Novel therapeutic targets

8. Conclusion

Siderophores represent one of the most elegant and efficient biochemical solutions evolved by living systems to overcome iron limitation. Their extraordinary affinity for ferric iron, remarkable structural diversity, and tightly regulated biosynthetic pathways highlight the sophistication of microbial survival strategies. Across terrestrial, aquatic, plant-associated, and host environments, siderophores ensure iron acquisition, maintain metal homeostasis, and support essential metabolic functions that sustain life. The diversity of siderophores spanning catecholate, hydroxamate, carboxylate, and mixed-type molecules reflects adaptation to distinct ecological niches and physicochemical constraints. Variations in ligand chemistry, denticity, molecular architecture, and transport systems allow organisms to fine-tune iron-scavenging efficiency under fluctuating environmental conditions. This chemical versatility also underpins their broader metal-binding capabilities, extending functional roles beyond nutrition to detoxification and geochemical cycling. From a biological perspective, siderophore systems are deeply integrated with microbial ecology, interspecies competition, symbiosis, and pathogenesis. Beneficial microorganisms employ siderophores to enhance plant nutrition and soil fertility, whereas pathogenic microbes utilize them as virulence determinants to circumvent host iron restriction. The ongoing molecular arms race between host defense mechanisms and microbial iron acquisition systems underscores the central importance of siderophore biology in infection dynamics.

Technologically, siderophores have emerged as powerful tools in sustainable agriculture, environmental remediation, and clinical therapeutics. Their application as biofertilizers and biocontrol agents supports eco-friendly crop production and reduces reliance on chemical inputs. In environmental systems, their metal-chelating capacity enables remediation of heavy metal pollution, wastewater treatment, and

regulation of trace metal mobility. In medicine, siderophore-based chelation therapies, diagnostic tools, and targeted drug delivery platforms are opening new frontiers in treatment of iron overload disorders, metal toxicity, and antimicrobial resistance. Advances in genomics, metabolomics, and synthetic biology are accelerating discovery of novel siderophores and enabling metabolic engineering for enhanced production. Structural modification and nano-biotechnological integration further expand their translational potential. However, challenges remain in large-scale production, cost-effective purification, stability optimization, and regulatory approval for field and clinical applications.

Future research should prioritize integrative approaches combining molecular genetics, systems biology, environmental chemistry, and translational biotechnology. Exploration of unexplored ecological niches, development of engineered microbial consortia, and design of synthetic siderophore analogues will broaden application horizons. A deeper understanding of siderophore-mediated metal interactions will also contribute to climate-resilient agriculture, circular bioeconomy strategies, and precision medicine. In summary, siderophores are multifunctional biomolecules of immense scientific and practical significance. Continued interdisciplinary research will unlock their full potential as sustainable solutions for global challenges spanning food security, environmental protection, and human health.

9. Acknowledgement:

The author(s) sincerely express their gratitude to the Hon'ble Vice Chancellor of Chhatrapati Shahu Ji Maharaj University, Kanpur, for fostering a research-oriented environment and for the continuous encouragement, as well as the Director of the School of Life Sciences and Biotechnology (SLSBT) for providing institutional support. Their support and vision have been instrumental in shaping the perspectives presented in this paper.

REFERENCES

- 1) Kumar, M., Kumar, D., Sharma, A., Bhadauria, S., Thakur, A., & Bhatia, A. (2024). Micronutrients throughout the life cycle: needs and functions in health and disease. *Current Nutrition & Food Science*, 20(1), 62-84.
- 2) Dutt, S., Hamza, I., & Bartnikas, T. B. (2022). Molecular mechanisms of iron and heme metabolism. *Annual review of nutrition*, 42, 311-335.

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

- 3) Zandi, P., & Schnug, E. (2022). Reactive oxygen species, antioxidant responses and implications from a microbial modulation perspective. *Biology*, *11*(2), 155.
- 4) Deb, A., Rumky, J., & Sillanpää, M. (2023). Fenton, photo-Fenton, and electro-Fenton systems for micropollutant treatment processes. In *Advanced oxidation processes for micropollutant remediation* (pp. 157-185). CRC press.
- 5) Mahaseth, T. (2015). *The iron-dependent cyanide and hydrogen peroxide co-toxicity in Escherichia coli and its catastrophic consequences for the chromosome* (Doctoral dissertation, University of Illinois at Urbana-Champaign).
- 6) Dhuldhaj, U., & Pandya, U. (2021). Diversity, function, and application of fungal iron chelators (Siderophores) for integrated disease management. In *Role of microbial communities for sustainability* (pp. 259-288). Singapore: Springer Singapore.
- 7) Raymond, K. N., Allred, B. E., & Sia, A. K. (2015). Coordination chemistry of microbial iron transport. *Accounts of chemical research*, *48*(9), 2496-2505.
- 8) Pecoraro, L., Wang, X., Shah, D., Song, X., Kumar, V., Shakoor, A., ... & Rani, R. (2021). Biosynthesis pathways, transport mechanisms and biotechnological applications of fungal siderophores. *Journal of Fungi*, *8*(1), 21.
- 9) Silale, A., & van den Berg, B. (2023). TonB-dependent transport across the bacterial outer membrane. *Annual review of microbiology*, *77*(1), 67-88.
- 10) Rodríguez, D., & González-Bello, C. (2023). Siderophores: chemical tools for precise antibiotic delivery. *Bioorganic & Medicinal Chemistry Letters*, *87*, 129282.
- 11) Ajinde, A. O., Dayo-Olagbende, O. G., & Akpor, O. B. (2024, April). Direct and indirect mechanisms of growth promotion by plant growth promoting rhizobacteria. In *2024 International conference on science, engineering and business for driving sustainable development goals (SEB4SDG)* (pp. 1-16). IEEE.
- 12) Satyal, S. S., Nath, M., Bhatt, M. D., Gohil, T., & Bhatt, D. (2021). Molecular mechanisms deciphering cross-talk between quorum sensing genes and major iron regulons in rhizospheric communities. In *Microbial Metatranscriptomics Belowground* (pp. 615-630). Singapore: Springer Singapore.
- 13) Jiang, K., Pang, X., Li, W., Xu, X., Yang, Y., Shang, C., & Gao, X. (2025). Interbacterial warfare in the human gut: insights from Bacteroidales' perspective. *Gut Microbes*, *17*(1), 2473522.
- 14) Maret, W. (2024). The extracellular metallometabolome: metallophores, metal ionophores, and other chelating agents as natural products. *Natural Product Communications*, *19*(8), 1934578X241271701.
- 15) Pradhan, N., Singh, S., Saxena, G., Pradhan, N., Koul, M., Kharkwal, A. C., & Sayyed, R. (2025). A review on microbe–mineral transformations and their impact on plant growth. *Frontiers in Microbiology*, *16*, 1549022.
- 16) Seyoum, Y., Baye, K., & Humblot, C. (2021). Iron homeostasis in host and gut bacteria—a complex interrelationship. *Gut Microbes*, *13*(1), 1874855.
- 17) Schalk, I. J. (2025). Bacterial siderophores: diversity, uptake pathways and applications. *Nature Reviews Microbiology*, *23*(1), 24-40.
- 18) Asad, M., Waqas, M., Ahmad, I., Ahmad, I., Khan, M. A. A., Sahar, P., ... & Gul, K. (2025). APPLICATION OF RHIZOSPHERIC BACTERIA FOR SUSTAINABLE AGRICULTURE PRACTICES, BIOREMEDIATION OF HEAVY METALS AND PRODUCTION OF BIOACTIVE COMPOUNDS. *Kashf Journal of Multidisciplinary Research*, *2*(05), 30-50.
- 19) Nawaz, T., Gu, L., Hu, Z., Fahad, S., Saud, S., & Zhou, R. (2024). Advancements in synthetic biology for enhancing cyanobacterial capabilities in sustainable plastic production: A green horizon perspective. *Fuels*, *5*(3), 394-438.
- 20) Mohsin, N., Khalid, S., Rasool, N., Aman, L., Kanwal, A., & Imran, M. (2025). Metallo-organic complexes containing transition metals; synthetic approaches and pharmaceutical aspects. *ChemPlusChem*, *90*(5), e202400748.
- 21) Kumar, R., Singh, A., & Srivastava, A. (2025). Xenosiderophores: bridging the gap in microbial iron acquisition strategies. *World Journal of Microbiology and Biotechnology*, *41*(2), 69.
- 22) Tsai, Y. H., Campalani, C., Toupay, T., Jacquemin, I., & Monbaliu, J. C. M. (2025). Catalytic Generation of Nitrosocarbonyls Under Mild Conditions: Advances, Challenges, and Opportunities. *ChemSusChem*, e202501565.
- 23) Zhang, S., Fan, S., He, H., Zhu, J., Murray, L., Liang, G., ... & Zhang, Y. (2025). Cyclic natural

- product oligomers: diversity and (bio) synthesis of macrocycles. *Chemical Society Reviews*, 54(1), 396-464.
- 24) Cai, J., Fang, Y., Liu, X., Adjei, M. O., & Fan, B. (2026). Revealing Microbial Siderophores: From Genes to Applications. *Microorganisms*, 14(2), 393.
 - 25) Dembitsky, V. M., Terent'ev, A. O., & Baranin, S. V. (2026). Life with Boron: Microbial Boron-Binding Siderophores, Adaptation, and Function. *Microbiology Research*, 17(3), 57.
 - 26) Chukwuma, C. E., Duckworth, O. W., & Baars, O. (2025). Degradation of catecholate, hydroxamate, and carboxylate model siderophores by extracellular enzymes. *Plos one*, 20(8), e0330432.
 - 27) Bhattacharjee, R., & Parli, B. V. (2025). Marine siderophores and its implications in changing polar ecosystem: a review. *Polar Biology*, 48(1), 37.
 - 28) Sharma, R., Kumar, D., Mir, N. R., Sharma, M., Singh, A., Mir, B. A., ... & Kapoor, N. (2025). Siderophores and their action against metalloids toxicity. In *Metalloids in Biology* (pp. 246-269). CRC Press.
 - 29) Hamzeh, M., Mirahsani, S., Ahmadpoor, F., & Farokhirad, S. (2025). From Single Ligand-Receptor Bond Strength to Collective Avidity: Mechanics-Guided Superselective Nanoparticle Adhesion to Biological Membranes. *Langmuir*, 41(51), 34210-34226.
 - 30) Li, Y., Yu, H., Li, L., Liu, Y., Kang, G., Liang, X., & Cao, Y. (2025). Advanced lithium extraction nanofiltration membrane with fast transport channels via competitive diffusion and reaction of rigid electropositive phenylbiguanide molecules. *Journal of Membrane Science*, 713, 123362.
 - 31) Gu, S., Shao, Z., Qu, Z., Zhu, S., Shao, Y., Zhang, D., ... & Li, Z. (2025). Siderophore synthetase-receptor gene coevolution reveals habitat-and pathogen-specific bacterial iron interaction networks. *Science Advances*, 11(3), eadq5038.
 - 32) Pantopoulos, K. (2025). BioIron: Origin, Chemical Properties, and Biological Functions. In *Iron Metabolism in Human Health and Disease* (pp. 1-15). Cham: Springer Nature Switzerland.
 - 33) Wilks, A., & Egoshi, R. (2025). Heme Trafficking and the Importance of Handling Nature's Most Versatile Cofactor. *Chemical Reviews*, 125(23), 11358-11378.
 - 34) Hamad, M. (2025). The Fight for Iron: A Central Theme in Host-Pathogen Interactions. In *Metabolic Dynamics in Host-Microbe Interaction* (pp. 249-289). Singapore: Springer Nature Singapore.
 - 35) Śmiga, M., & Olczak, T. (2025). Exploring heme and iron acquisition strategies of *Porphyrromonas gingivalis*—current facts and hypotheses. *FEMS Microbiology Reviews*, 49, fuaf019.
 - 36) Rocha, B. M., Pinto, E., Sousa, E., & Resende, D. I. (2025). Targeting siderophore biosynthesis to thwart microbial growth. *International Journal of Molecular Sciences*, 26(8), 3611.
 - 37) Shafi, Z., Shahid, M., Ilyas, T., & Singh, K. (2025). Next-generation perspectives on microbially synthesized siderophores: molecular engineering, multi-omics insights, and applications for smart climate-resilient crops. *World Journal of Microbiology and Biotechnology*, 41(10), 393.
 - 38) Musialowski, M., Kowalewska, Ł., Stasiuk, R., Krucoń, T., & Debiec-Andrzejewska, K. (2023). Metabolically versatile psychrotolerant Antarctic bacterium *Pseudomonas* sp. ANT_H12B is an efficient producer of siderophores and accompanying metabolites (SAM) useful for agricultural purposes. *Microbial Cell Factories*, 22(1), 85.
 - 39) Bryukhanov, A. L., Klimko, A. I., & Netrusov, A. I. (2022). Antioxidant properties of lactic acid bacteria. *Microbiology*, 91(5), 463-478.
 - 40) Zimmermann, A., Prieto-Vivas, J. E., Voordeckers, K., Bi, C., & Verstrepen, K. J. (2024). Mutagenesis techniques for evolutionary engineering of microbes—exploiting CRISPR-Cas, oligonucleotides, recombinases, and polymerases. *Trends in Microbiology*, 32(9), 884-901.
 - 41) Gomes, A. F., Sousa, E., & Resende, D. I. (2024). A practical toolkit for the detection, isolation, quantification, and characterization of siderophores and metallophores in microorganisms. *ACS omega*, 9(25), 26863-26877.
 - 42) Dhull, P., Dunuweera, S., Bietsch, J., Bandu, R., Wannere, C., Achanta, S., ... & Senanayake, C. (2025). Recent advances and application of liquid chromatography in pharmaceutical industry. *Journal of Liquid Chromatography & Related Technologies*, 48(6-10), 168-187.
 - 43) Prasad, R. D., Sarvalkar, P. D., Prasad, N., Prasad, S. R., Prasad, R. S., Prasad, R. B., ... & Ghosh, J.

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

- (2024). A review on spectroscopic techniques for analysis of nanomaterials and biomaterials. *ES Energy and Environment*, 27, 1264.
- 44) Oubohssaine, M., Sbabou, L., & Aurag, J. (2025). Enhancing ecosystem restoration and soil productivity through PGPR: a sustainable approach to bioremediation and biofertilization. *Discover Applied Sciences*, 7(12), 1379.
- 45) Shafi, Z., Shahid, M., Ilyas, T., & Singh, K. (2025). Next-generation perspectives on microbially synthesized siderophores: molecular engineering, multi-omics insights, and applications for smart climate-resilient crops. *World Journal of Microbiology and Biotechnology*, 41(10), 393.
- 46) Pradhan, N., Singh, S., Saxena, G., Pradhan, N., Koul, M., Kharkwal, A. C., & Sayyed, R. (2025). A review on microbe–mineral transformations and their impact on plant growth. *Frontiers in Microbiology*, 16, 1549022.
- 47) Chen, Q., Song, Y., An, Y., Lu, Y., & Zhong, G. (2024). Mechanisms and impact of rhizosphere microbial metabolites on crop health, traits, functional components: A comprehensive review. *Molecules*, 29(24), 5922.
- 48) Kumar, R., Singh, A., Shukla, E., Singh, P., Khan, A., Singh, N. K., & Srivastava, A. (2024). Siderophore of plant growth promoting rhizobacterium origin reduces reactive oxygen species mediated injury in *Solanum* spp. caused by fungal pathogens. *Journal of Applied Microbiology*, 135(2), 1xae036.
- 49) Shao, Z., Gu, S., Zhang, X., Xue, J., Yan, T., Guo, S., ... & Wei, Z. (2024). Siderophore interactions drive the ability of *Pseudomonas* spp. consortia to protect tomato against *Ralstonia solanacearum*. *Horticulture Research*, 11(9), uhae186.
- 50) Wang, Y., Huang, W., Ali, S. W., Li, Y., Yu, F., & Deng, H. (2022). Isolation, identification, and characterization of an efficient siderophore producing bacterium from heavy metal contaminated soil. *Current microbiology*, 79(8), 227.
- 51) Tayal, A., Kaur, J., Sadeghi, P., & Maitta, R. W. (2025). Molecular Mechanisms of Iron Metabolism and Overload. *Biomedicines*, 13(9), 2067.
- 52) Rose, A. E., Fansler, R. T., & Zhu, W. (2025). Commensal resilience: ancient ecological lessons for the modern microbiota. *Infection and Immunity*, 93(6), e00502-24.
- 53) Takemura, K., Kolasinski, V., Del Poeta, M., Vieira de Sa, N. F., Garg, A., Ojima, I., ... & Pereira de Sa, N. (2025). Iron acquisition strategies in pathogenic fungi. *MBio*, 16(6), e01211-25.
- 54) Xiao, J., Bu, Y., Tao, Y., Zhou, Z., Zou, J., Cui, W., & Yang, Z. (2025). The Role of Bacterial Siderophores in Infection Therapy: From Anti-Infective Mechanisms to Therapeutic Advances. *International Journal of Nanomedicine*, 15951-15978.
- 55) Lan, P., Lu, Y., Fu, Y., Yu, Y., & Zhou, J. (2025). Siderophores and beyond: A comprehensive review of iron acquisition in *Klebsiella pneumoniae*. *Virulence*, 16(1), 2550621.
- 56) Prasad, R., Shivay, Y. S., & Mandi, S. (2022). Phytosiderophores and absorption of iron and other cations by plants. In *Cation transporters in plants* (pp. 385-399). Academic Press.
- 57) Liu, X., Luo, Q., Zhao, Y., Ren, P., Jin, Y., & Zhou, J. (2025). The ferroptosis–mitochondrial axis in depression: unraveling the feedforward loop of oxidative stress, metabolic homeostasis dysregulation, and neuroinflammation. *Antioxidants*, 14(5), 613.
- 58) Pijuan, J., Moreno, D. F., Yahya, G., Moisa, M., Ul Haq, I., Krukiewicz, K., ... & Cavalu, S. (2023). Regulatory and pathogenic mechanisms in response to iron deficiency and excess in fungi. *Microbial Biotechnology*, 16(11), 2053-2071.
- 59) Yong, C. W., Deng, B., Liu, L. M., Wang, X. W., & Jiang, H. B. (2023). Diversity and evolution of iron uptake pathways in marine cyanobacteria from the perspective of the coastal strain *Synechococcus* sp. strain PCC 7002. *Applied and Environmental Microbiology*, 89(1), e01732-22.
- 60) Nawaz, T., Gu, L., Fahad, S., Saud, S., Bleakley, B., & Zhou, R. (2024). Exploring sustainable agriculture with nitrogen-fixing cyanobacteria and nanotechnology. *Molecules*, 29(11), 2534.
- 61) Qiu, G. W., Koedooder, C., Qiu, B. S., Shaked, Y., & Keren, N. (2022). Iron transport in cyanobacteria—from molecules to communities. *Trends in Microbiology*, 30(3), 229-240.
- 62) Kermeur, N., Pédrot, M., & Cabello-Hurtado, F. (2023). Iron availability and homeostasis in plants: a review of responses, adaptive mechanisms, and signaling. *Plant abiotic stress signaling*, 49-81.

Exploring Siderophore Diversity: Production, Extraction And Functional Applications

- 63) Kumar, R., Jindal, K., Singh, J. P., & Singh, S. P. (2024). Unravelling the Dynamic Role of Beneficial Microbes in Regenerative Agriculture. In *Regenerative Agriculture* (pp. 13-22). CRC Press.
- 64) Anas, M., Falak, A., Hassan, S., Khattak, W. A., Saleem, M. H., Khan, K. A., ... & Fahad, S. (2025). Microbial interactions and bacterial responses to metal stress in plants: mechanisms, adaptations, and applications for sustainable agriculture. *Journal of Crop Health*, 77(1), 36.
- 65) Gräff, Á. T., & Barry, S. M. (2024). Siderophores as tools and treatments. *npj Antimicrobials & Resistance*, 2(1), 1–12.