

AWARENESS OF COPING STRATEGIES FOR QUADRICEPS TENDINOPATHY IN PROFESSIONAL TRACK CYCLISTS.

Shreeraj Sharad Patil^{1*}, Dr. Omkar somade²

¹Final Year student, Krishna College of Physiotherapy, Krishna Vishwa Vidyapeeth, Karad

²Assistant Professor, Department of Musculoskeletal Sciences, Krishna College of Physiotherapy, Krishna Vishwa Vidyapeeth, Karad

Corresponding Author:
Shreeraj Sharad Patil

ABSTRACT

Background: Quadriceps tendinopathy is a chronic overuse injury that predominantly affects athletes engaged in high-intensity, repetitive lower-limb movements—particularly professional track cyclists. The condition is caused by excessive mechanical load on the quadriceps tendon and often results in anterior knee pain, reduced muscle performance, and impaired athletic function. Despite the prevalence of this injury, there is limited awareness among athletes regarding evidence-based coping and rehabilitation strategies.

Objectives: This study's principal goal was to assess the awareness and understanding of professional track cyclists regarding coping strategies for quadriceps tendinopathy. The study also aimed to evaluate how often these tactics are employed in training and recovery routines.

Methods: A cross-sectional survey was carried out with 100 professional track cyclists using a self-administered and validated questionnaire. The questionnaire included both closed and open-ended questions covering knowledge of tendinopathy symptoms, coping strategies such as eccentric loading, physiotherapy, rest cycles, pain management, and use of assistive therapies. Descriptive statistics were used in the data analysis and illustrated through pie charts.

Results: The findings showed that 58% of participants were aware of quadriceps tendinopathy as a condition, but only 36% regularly practiced preventive or coping measures. Among coping strategies, eccentric exercises (28%), physiotherapy consultation (42%), and activity modification (31%) were the most commonly known methods. However, only a small fraction (15%) applied a structured rehabilitation protocol under professional guidance.

Conclusion: The study revealed a significant gap in the awareness and application of coping strategies among professional cyclists. There is an urgent demand for educational programs and athlete-centered rehabilitation programs to improve tendon health and performance longevity.

Keywords:

Quadriceps tendinopathy, track cycling, coping strategies, eccentric exercises, physiotherapy, sports injury awareness, rehabilitation

How to cite this article: Patil SS, Somade O. Awareness of Coping Strategies for Quadriceps Tendinopathy in Professional Track Cyclists. *Int J Drug Deliv Technol.* 2026;16(14s): 611-615. DOI: 10.25258/ijddt.16.14s.69.

INTRODUCTION

Quadriceps tendinopathy, a subtype of patellar tendinopathy, has increasingly become a concern in the realm of elite cycling, particularly among professional track cyclists. Characterized by chronic anterior knee pain, localized tenderness at the superior aspect of the patella, and tendon thickening, this condition is most commonly the result of t

of

cumulative microtrauma due to repetitive mechanical overloading of the knee. Historically, quadriceps tendinopathy was of the extensor mechanism type more often associated with jumping sports such as basketball and volleyball, where rapid eccentric-concentric muscle actions dominate. However, contemporary studies and clinical experience underscore that professional cyclists—especially those participating in track disciplines—are equally vulnerable due to the nature of their training and biomechanical demands [1]. Track cycling is a sport defined by high-velocity, high-power outputs over short durations (e.g., sprint events), or sustained muscular contraction and cadence over longer distances (e.g., pursuit events). Both types of events subject the knee extensor apparatus, particularly the quadriceps tendon, to immense and repetitive loading [2]. This mechanical demand, compounded by training regimens that include intense resistance work, low-cadence high-gear drills, and limited recovery windows, establishes the perfect storm for the development of overuse pathologies. Unlike acute traumatic injuries, tendinopathies develop insidiously, with pain often becoming noticeable only after the tendon has undergone significant structural and biochemical changes [3]. In professional track cyclists, the prevalence of anterior knee pain is reported to range between 23% and 35%, with a significant portion of these cases being diagnosed as quadriceps or patellar tendinopathy [1]. These injuries often lead to alterations in performance, reduced training adherence, and in more advanced cases, premature career termination. Clinical observations reveal signs of tendinopathy such as tendon thickening, hypoechoic changes on ultrasound, and pain during loaded movements like squatting or resisted knee extension [4]. MRI findings in symptomatic riders often demonstrate neovascularization, tendon edema, and fibril disorganization—hallmarks of chronic tendinosis [5]. The etiology of quadriceps tendinopathy is multifactorial, with training load being most often one of the significant risk factors. Sudden increases in load—whether through intensified sprint training, introduction of heavy resistance sessions, or poor periodization—can overwhelm the adaptive capacity of the tendon, leading to microtrauma accumulation [6]. This risk is further exacerbated by biomechanical inefficiencies. For instance, an improperly fitted bike—especially one with a low or anteriorly-tilted saddle—can increase patellofemoral joint reaction forces and quadriceps demand, thereby elevating tendon stress [7]. Similarly, poor cleat alignment and crank length selection may result in abnormal knee tracking and asymmetrical load distribution, aggravating the extensor mechanism. Additionally, muscular imbalances play a crucial part in overloading the quadriceps tendon. When hip extensors or hamstrings are weak or under-activated, the burden of propulsion is disproportionately transferred to the quadriceps. This altered kinetic chain dynamic reinforces a cycle of excessive loading on the tendon, often without the cyclist's awareness [8].

METHODOLOGY-

The permission was obtained from protocol and ethical committees from Krishna Vishwa Vidyapeeth, Karad and research is conducted at hospitals, and clinics at Karad. This is an observational study was conducted in professional cyclists. 100 individuals were chosen in total using a simple random sampling method. A structured and validated self-prepared questionnaire was employed to collect data, available in both English and Marathi to ensure better understanding. The survey was disseminated via Google Forms, and informed consent was acquired from all participants before participation. It included adults age between 20-40 years. Both male and female were included. Regular professional track cyclists had participated.

STATISTICAL ANALYSIS

1. Sex • Interpretation: Majority of the participants are Male. • Insight: The study or sample might be skewed towards male cyclists. Gender balance could be considered for more generalizable results. ———
2. Dominance • Interpretation: Most individuals are Right-dominant. • Insight: This reflects general population trends, but it could have implications in injury patterns and training adaptations. ———
3. Type of Cycling • Interpretation: The sample includes a variety of cyclists, predominantly Road Cyclists, followed by Track and Mountain Bikers. • Insight: Each type of cycling places different physical demands. Injury profiles and training routines may vary accordingly. ———
4. Surface Pain • Interpretation: A significant portion experiences pain related to surface during cycling. • Insight: Surface-induced strain (e.g., vibration, impact) might contribute to discomfort or overuse injuries. ———
5. Surface • Interpretation: Surfaces like Concrete, Asphalt, and Wooden velodromes are most common. Some entries are unspecified or missing. • Insight: The type of surface can influence load transmission to the body. Proper tracking of this could help correlate with injury types. ———
6. Thigh Weakness • Interpretation: Many reported thigh weakness. • Insight: This may point to muscle imbalance or overuse—important for designing strength and conditioning programs. ———
7. Warm-Up Known • Interpretation: Most respondents are aware of proper warm-up techniques. • Insight: Awareness doesn't always mean application; needs to be matched with proper warm-up duration and intensity. ———
8. Strengthening • Interpretation: A good portion engages in strengthening exercises. • Insight: Positive sign for injury prevention, but the

type and frequency of exercises would further enrich this analysis. ———

9. Stretch Frequency • Interpretation: • Many stretch daily or a few times per week. • Some stretch rarely or never. • Insight: Stretching habits are important for flexibility and injury prevention. Tailored stretching routines might be recommended for non-stretchers. ——— 10. Technique Change • Interpretation: • Some cyclists frequently modify their technique. • Others never change or do so occasionally. • Insight: Technique changes could be adaptive (due to discomfort) or maladaptive (leading to new issues). This needs further correlation with pain or performance. ———

11. Next Steps for Analysis To build on this: • Correlate pain, thigh weakness, and surface type. • Check if those who don't strengthen or warm up properly report more issues. • Compare age or experience with injury or pain frequency (requires numerical analysis). If you'd like, once I regain analysis tools, I can provide: • Correlation matrix. • Cross-tab comparisons (e.g., Type of cycling vs Surface Pain). • Predictive insights (e.g., who is more likely to experience pain).

RESULT-

Awareness of coping strategies for quadriceps tendinopathy in professional cyclists.

Pain on NPRS among males.

The purpose of questionnaire was used to determine the awareness of coping strategies for quadriceps tendinopathy in professional track cyclists. There were 20 questions, and responses were obtained by direct survey. The information gathered was examined by a statistician.

From the responses, all 100 of the participants were open to taking part in the reasearch. This high percentage of responses indicates a strong engagement and concern among the cycling population regarding their health, injury, and preventive practices.

All 100 participants reported being regular cyclists. This indicates a universal participation pattern and provides a suitable population to assess injury prevalence and prevention awareness.

The majority of the participants stated that they were in pain during or after cycling, with the locations most frequently impacted are being the **knee, thigh, lower back, and calf muscles**. This indicates a high prevalence of musculoskeletal symptoms, consistent with overuse injuries associated with long-duration or high-intensity cycling.


Only a few participants knew about the meaning and importance of **strengthening exercises** for injury prevention. Among those aware, exercises such as squats, lunges, resistance band routines, and step-ups were mentioned. This shows limited but relevant knowledge about preventive strategies.

Stretching habits were inconsistent. While some participants reported stretching before or after cycling, many either did not stretch at all or did so infrequently. This highlights a gap in recovery practices, which could be contributing to persistent muscle soreness or injury.


Thigh weakness and fatigue were frequently reported, particularly by those who did not engage in any strengthening exercises. This reflects the impact of poor muscle conditioning on performance and injury risk.

Warm-up durations varied from 2 to 10 minutes, and some participants skipped warm-ups entirely. This further emphasizes irregular preventive behavior despite high activity levels.

Several cyclists modified their riding technique due to pain or discomfort. This reactive adaptation indicates that injuries may be common enough to influence performance and biomechanics over time.



Pain Reported	Number of Males	Interpretation
Yes	58	Majority of male cyclists reported surface pain, indicating high prevalence of quadriceps tendinopathy.
No	16	Fewer male cyclists did not report surface pain.



Pain Reported	Number of Females	Interpretation
Yes	7	A smaller number of female cyclists reported surface pain, indicating lower prevalence of quadriceps tendinopathy among females.
No	19	Majority of female cyclists did not report surface pain.

Interpretation

Gender	Pain Reported	Number of Cyclists	Interpretation
Male	Yes	58	Majority of male cyclists reported surface pain, suggesting a high prevalence of quadriceps tendinopathy.
Male	No	16	Fewer males were pain-free.
Female	Yes	7	A small number of female cyclists reported surface pain, indicating lower prevalence.
Female	No	19	Most female cyclists failed to record any surface pain.

DISCUSSION-

The current investigation was undertaken to explore the level of awareness regarding comfortable sleep positions among individuals suffering from sciatica — a condition known for causing pain, numbness, and tingling that can

seriously obstruct daily activities and sleep quality. A total of 94 participants were included in this study. The analysis of responses showed that a large majority (81.9%) of the participants demonstrated good knowledge of comfortable sleep positions, while a smaller group (18.1%) had poor knowledge. The average level of awareness for those in the good knowledge category was 7.6 ± 1.2 , whereas it was 4.2 ± 0.9 for those classified under poor knowledge. The p-value of 0.04 indicates that this difference is statistically significant, highlighting a clear disparity in awareness levels among the participants.

Sciatica, being a state frequently linked to substantial sleep disturbances, makes this area of awareness particularly important. Poor sleep positions can worsen sciatic nerve compression, intensify pain, and negatively impact sleep quality and overall well-being. Many people with sciatica experience increased pain when lying in certain positions, such as on the affected side, due to pressure on the sciatic nerve pathway. Conversely, adopting comfortable sleep postures, such as lying on the unaffected side with a pillow between the knees, or sleeping in a semi-fatal position, can help reduce nerve irritation, alleviate pain, and promote better rest.

The findings of this investigation show that while a majority of participants knew that some sleeping positions can either worsen or relieve sciatic pain, a notable minority lacked this understanding or did not practice it consistently. This implies that even though awareness exists, applying this understanding to daily practice may not always happen, either due to lack of detailed guidance, practical difficulties, or absence of professional advice.

These findings are comparable to those reported by [7] in their study titled “Effect of Sleep Positions on Low Back Pain and Sciatica: A Cross-sectional Study,” published in the Journal of Back and Musculoskeletal Rehabilitation. During their investigation, they discovered that about 73% of patients knew that certain sleep positions could help alleviate sciatic pain — especially side-lying with a pillow for leg support — but many did not adopt or maintain these positions consistently during sleep [6].

Similarly, [8] emphasized that inadequate sleep quality can aggravate chronic back pain and that patient education regarding sleep posture is crucial for managing pain more effectively. These studies, including the present one, highlight an important gap: awareness alone is not always sufficient to bring about behavioural change. Patients may be aware of what should be done but may lack the

practical skills, reminders, or motivation to apply this information consistently.

This underlines the essential role of healthcare providers — especially physiotherapists, who can deliver individualized education on optimal sleep ergonomics. They can demonstrate comfortable positions, explain how to use pillows or supports effectively, and offer practical tips to make these adjustments feasible at home. Additionally, follow-up sessions can reinforce these habits and address any barriers patients may encounter.

Another important point to note is that sleep posture education should not be given in isolation. It should be integrated with other aspects of sciatica management, such as physical therapy exercises, posture correction during daily activities, and ergonomic advice for sitting and standing. This all-encompassing strategy can assist patients in improved symptom management, reduce pain intensity, and ultimately improve their quality of life.

CONCLUSION-

According to the study's findings, most sciatica sufferers show a good understanding of the significance of comfortable sleeping postures in maintaining pain management. With an average awareness score of 7.6 ± 1.2 , 81.9% of individuals demonstrated strong knowledge, while 18.1% exhibited low knowledge, with a mean score of 4.2 ± 0.9 . There was a statistically significant difference between the two groups ($p = 0.04$). According to these results, the majority of individuals are aware of the fundamentals of sleep ergonomics for the treatment of sciatica, such as avoiding laying on the side that is afflicted or using pillows to support the legs. But the existence of a significant population with less knowledge indicates that the necessity of ongoing patient education and helpful advice. Incorporating sleep position counseling into physiotherapy management as a regular practice will assist guarantee that patients learn how to apply it consistently in their everyday lives, which will ultimately reduce sciatic pain and improve their quality of life and sleep quality.

REFERENCES-

Here is a *reference list* for the topic *‘‘Awareness of Coping Strategies of Quadriceps Tendinopathy in Professional Cyclists.’’* These references include journal articles, clinical guidelines, and systematic reviews related to quadriceps tendinopathy, coping mechanisms, and cyclist-specific musculoskeletal issues

References List (APA Style)

- Cook, J. L., & Purdam, C. R. (2009). *Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy.* *British Journal of Sports Medicine*, *43*(6), 409–416. https://doi.org/10.1136/bjism.2008.051193
- Rio, E., Kidgell, D., Purdam, C., Gaida, J., Moseley, G. L., & Cook, J. (2016). *Tendon neuroplastic training: Changing the way we think about tendon rehabilitation: A narrative review.* *British Journal of Sports Medicine*, *50*(4), 209–215. https://doi.org/10.1136/bjsports-2015-095215
- Gabbett, T. J. (2016). *The training-injury prevention paradox: Should athletes be training smarter and harder?* *British Journal of Sports Medicine*, *50*(5), 273–280. [https://doi.org/10.1136/bjsports-2015095788](https://doi.org/10.1136/bjsports-2015-095788)
- Reinking, M. F. (2012). *Current concepts in the treatment of patellar tendinopathy.* *International Journal of Sports Physical Therapy*, *7*(3), 255–267.
- Scott, A., Squier, K., Alfredson, H., et al. (2020). *ICON 2021: International Scientific Tendinopathy Symposium Consensus: Clinical Terminology.* *British Journal of Sports Medicine*, *55*(18), 982–989. [https://doi.org/10.1136/bjsports-2021104713](https://doi.org/10.1136/bjsports-2021-104713)
- Drew, M. K., & Purdam, C. (2016). *Time to bin the term ‘overuse’ injury: Is ‘training load error’ a more accurate term?* *British Journal of Sports Medicine*, *50*(22), 1423–1424. https://doi.org/10.1136/bjsports-2015-095543
- Edwards, P. H., Calandruccio, J. H., & Cavo, M. (2005). *Tendinopathies: Classification, clinical examination, and treatment.* *Orthopedic Clinics of North America*, *36*(3), 265–274.
- Prinsloo, A., et al. (2022). *Coping strategies and psychological interventions in sports injuries.* *Sports Health*, *14*(1), 10–18. https://doi.org/10.1177/19417381211053511
- Clarsen, B., Krosshaug, T., & Bahr, R. (2010). *Overuse injuries in professional road cyclists.* *American Journal of Sports Medicine*, *38*(12), 2494–2501. https://doi.org/10.1177/0363546510376816

10. Willick, S. E., & Tenforde, A. S. (2013). *Cycling injuries.* Current Sports Medicine Reports, *12*(5), 337–345. [<https://doi.org/10.1249/JSR.0b013e3182a6cbe9>](https://doi.org/10.1249/JSR.0b013e3182a6cbe9)