

Foreign Body Aspiration in Children – A Case Series

Kulkarni TP^{1*}, Harwalkar VS², Kokatanur S³^{1,2,3}SBMPMC&RC, BLDE (DU), Vijayapur¹Email: trimal.kulkarni@gmail.com²Email: veena.harwalkar@bldedu.ac.in³Email: kokatnurrshridhar@gmail.com

ABSTRACT

Introduction: Foreign body aspiration (FBA) in children typically presents with persistent or recurrent cough. This issue is seen mostly among the toddlers and preschoolers due to active pincer grasp and mouthing reflex. Incidence is more in males probably due to the nature to explore the surroundings. Other common symptoms associated with FBA are breathing difficulty. The problem is making an early diagnosis wherein the sharp clinical acumen, high index of suspicion is needed by the treating pediatrician because history may not be elicited in many situations.^{1,2} Some of the clues can be absence of fever in an acute accidental event, and sudden onset of symptoms like cough and breathing difficulty.^{2,3} Other signs developed could be due to the site of impaction in the respiratory tract like stridor. An accurate examination picking up the decreased or absent air entry on the affected side can help to arrive at a quick diagnosis.

Diagnosis is aided by typical features of obstructive emphysema, collapse or mediastinal shift seen on a chest radiograph.^{2,3} In some cases clinical features will not be peculiar as seen above. Or else in some instances the duration of symptoms may not be acute. Also in some cases of delayed presentations or small radiolucent foreign bodies chest x-ray may show subtle signs rather than those mentioned above.^{4,5} In such cases many patients are diagnosed late or undergo many higher investigations like chest Tomography (CT) ending up bearing radiation exposure and also the cost. A pediatrician can sometimes be misled by such delayed or complicated presentations of foreign body aspiration.⁶ Here we discuss some of these cases which can help in timely diagnosing FBA in children.

Keywords: foreign body aspiration, chest x-ray, flexible bronchoscopy, rigid bronchoscopy.

How to cite this article: Kulkarni TP, Harwalkar VS, Kokatanur S. Foreign Body Aspiration in Children – A Case Series. *Int J Drug Deliv Technol.* 2026;16(18s): 714-719. DOI: 10.25258/ijddt.16.18s.76

Case 1: Persistent cough

8 years male child was brought by mother with complaints of cough on and off since Four weeks. This child appeared to be well nourished and there was no history of any Coughing patient in the family. There was no history of any environmental fumes exposure, Recurrent fever, or weight loss in the recent past. No evidence of recurrent wheezing or Asthma in the family. But the mother was very particular about the coughing since four Weeks. This child also received symptomatic treatment from three general practitioners in During this period without any improvement in the cough.

On examination this child had Heart rate (HR)-88/min, respiratory rate (RR)-26/min Blood pressure (BP)-110/72 mm of Hg and saturation of 94%. Rest of the general Examination was normal and the respiratory system examination revealed normal inspection And palpatory findings apart from a mild decrease in air entry in the right hemi thorax

Which was probably not conspicuous so missed by earlier practitioner.

In the absence of any clinical clues we decided to investigate this child in our tertiary care Center with complete blood picture and chest radiograph posterior-anterior (PA) view. The CBP was in normal limits (HB-116G/L, TC-5×10⁹/L, DC – N-68%, L-30%, PCV-0.34, RBC-4.89 ×10¹²/L PLT-240×10⁹/L MCH 31 Pg MCV-72fL, MCHC-316g/L) and x-ray as Shown below in figure 1.

Initially we labelled this child may have habit cough and counselled about the benign nature Of the entity to the mother. But the child was noticed to cough even in the night.

Hence we decide to do flexible bronchoscopy in this child which revealed a small broken Piece of groundnut in the right lung. This was later on removed by rigid bronchoscopy on the Same day. Child was shifted to ward and required nebulization with salbutamol and Parenteral antibiotic for a week before we could discharge him. This child was followed up in

Foreign Body Aspiration in Children – A Case Series

Outpatient department after 2 weeks and 6 weeks and found to be asymptomatic.




Figure 1 mild haze (short arrow) and hyperinflation (long arrow) in right lung.

Case 2. Pyrexia of unknown origin

7 years male child presented to our outpatient department with complaints of fever on and off Since two months. This child was treated by many general practioners in rural areas who Treated him with multiple courses of oral antibiotics with symptom free intervals of two to Three days in between but undocumented. The fever was mild to moderate in severity, Intermittent subsided with paracetamol, not associated with chills, rigors or rash. There was No history of cough or cold, vomiting, loose stools. No history of headache, dyspnea, Dysuria, joint pains. There was no significant past or family history of any illness. On

Page 4

Examination he appeared to be normal built child with good nutrition and his vitals were Temperature- 101 degrees of Fahrenheit, HR- 130/min RR-32/min BP-98/74mm of hg SPO2-94%. Systemic examination was normal apart from mild decrease in air entry to right Side. In the absence of any clue preliminary Investigations in the form of complete Blood Picture, urine routine, and chest x-ray A-P view, CRP, ESR and blood culture were done.




Figure: 2 mild haziness in right lung in the middle and lower lobe

CBP (HB-116 G/L, TC- $12 \times 10^9/L$, DC – N-68%, L-30%, PCV-0.34, RBC- $4.99 \times 10^{12}/L$

PLT- $250 \times 10^9/L$ MCH 30 Pg MCV-70fL, MCHC-306g/L). The child was admitted for

Observation. Blood culture did not show any growth. CRP was 8 mg/L, Dengue serology was Negative. The child continued to spike fever even after normal initial Investigations.

After admission Parenteral empirical antibiotics were started and 2-D

Echocardiography was done to rule out any vegetation over valves or coronary dilatation, Even if inflammatory markers like CRP and ESR were normal to rule out Kawasaki or FIRES which were seen commonly in the Post-covid pandemic. Even after giving parenteral Antibiotics to this child for 72 hours fever Continued to spike till 101 degree Fahrenheit.

There was no focus from preliminary investigations and previous chest x-ray was normal. We Did one more that revealed mild haziness in right lung in the middle and lower lobe.

During this stay the child was visited by many relatives and other family Members of the House. Coincidentally we obtained a history of choking episode when this Child was eating

Foreign Body Aspiration in Children – A Case Series

a tamarind seed 2 months before, from his grandmother who had come to see her grandson.

This was a clue for cause of fever and we decided to go for flexible bronchoscopy

Of this child which revealed a small broken piece of tamarind seed with debris around it in

The lower bronchus which needed clearing before retrieval with rigid Bronchoscopy.

This child was afebrile after 48 more hours of retrieval of foreign body.

He was given antibiotics for three more days and discharged. Later he was followed up in the Outpatient department and was asymptomatic.

Case 3: Air leaks

An 8-month-old male infant presented with cough and noisy breathing on and off for 2

Weeks. These symptoms partially responded to treatment with antibiotics and cough syrups

Which were prescribed by general practitioners in rural areas. On admission infant had

Breathlessness and cough and needed admission to the pediatric intensive care unit.

On examination child was afebrile, HR-170/min, RR-66/min, BP-76/38 mm of Hg.

Spo2-90% requiring nasal prongs. Noisy breathing was heard. General physical examination

Was showing swelling in neck and axilla. There was crepitus over these swollen areas.

Systemic examination showed normal inspection and palpation but on auscultation the air

Entry was reduced on the Right side. CBP was (HB116G/L, TC- 5×10^9 /L, DC: N68%, L30%,

PCV-0.34, RBC- 4.89×10^{12} /L PLT- 240×10^9 /L MCH 31 Pg MCV-72fL, MCHC-316g/L)

Normal and the initial chest radiograph showed mediastinal widening and Subcutaneous




Figure 3a pneumomediastinum and bilateral subcutaneous emphysema

Emphysema with an artefact seen over the spine.

There was no apparent cause for

Mediastinal widening and noisy breathing (which was stridor) continued so the child was

Stabilized in picu and one more radiograph was done. In the Subsequent radiograph the

Artefact appeared to be a pin-like foreign body. So this child after initial

Stabilization was posted for immediate fiberoptic bronchoscopy which revealed the pin

Obstructed in the right main bronchus. This was removed by direct laryngoscopy soon after

Fiberoptic bronchoscopy. Subcutaneous emphysema was managed conservatively over

Following 4 days of observation. The child was treated with prophylactic antibiotics and

Nebulization with bronchodilators until fifth day when he was asymptomatic and accepting

Feeds hence discharged. Subcutaneous emphysema and pneumomediastinum with the pin at

Carina (could be missed as artefact due to superimposition on the thoracic spine.)

But the subsequent image figure 3b which was taken after stabilization was showing the pin

To slide towards the right main bronchus and become identifiable as pin.

Foreign Body Aspiration in Children – A Case Series




Figure 3b pin shown by white arrow.



Figure 3c

Figure 3c shows the Pin after it was removed

Discussion:

Foreign body aspiration in children is a frequently handled problem by all the pediatricians.

Sometimes it can lead to emergency or life threatening situation especially when it occurs in infants and preschoolers and is unnoticed by parents.

Some instances on the other hand can present as subacute or chronic illness particularly if they produce subtle pointers in initial investigations like complete blood picture and chest radiograph⁴. Also it is more frequently seen in males³

Organic material like peanuts, tamarind seeds are commonly seen instead of inorganic²

Material like pins, ball- pen caps. Organic foreign bodies are prone to produce inflammation

And fever like symptoms particularly if retained for longer duration.⁵

Such foreign bodies may pose difficulty in diagnosis and hence delay in treatment.

Other problems are many a times these foreign bodies could be fragmented and at many

Locations.^{6, 7} hence it is rewarding and crucial to repeat x-ray particularly in case of FBA.

In some scenarios the examination may produce very minimal or no signs and diagnosis

Cannot be made particularly if history is not obtained from all the members involved in the care of the child or if aspiration or choking event was unnoticed by any adult.^{2,5}

Hence at times pediatricians end up advising chest tomography in such children.

CT scan though non-invasive but exposes the child to ionizing radiations and can be negative

In cases of radiolucent foreign bodies. Although low dose CT scans are considered sensitive

And specific for detecting FBA in children yet the risk of exposure precludes its widespread

Use in doubtful cases.⁸

In many situations there is temporal discrepancy between clinical features and initial

Chest x-ray, which could delay the diagnosis until the treating pediatrician has a high index

Of suspicion and diagnostic bronchoscopy is done. Here ultra-low- dose CT scan can be

Considered⁹. But some studies have shown that a normal CT chest does not rule out FBA¹⁰

Flexible bronchoscopy is gold standard normally and more rewarding where history of FBA

Is not obtained or accidental aspiration¹¹. It would be worthwhile to have flexible

Bronchoscopy as an office Procedure. There has been growing evidence towards use of

Flexible Bronchoscopy Not Only for diagnosis but also for therapeutic use over rigid

Bronchoscopy^{12, 13}

Although Many Factors like the location, size, duration of symptoms and local

Pathology and Otorhinologist Preference may play role in the choice of the bronchoscopy

Used for the removal of foreign Body.^{14, 15}

Abbreviations: foreign body aspiration- FBA,
Chest Tomography- CT Chest
Chest radiograph- chest x-ray.

Acknowledgments: NIL

Conflict of interest: NIL

References:

1. Kliegman RM, St. Geme JW, Blum NJ, Tasker RC, Wilson KM, Schuh AM, et al. Nelson Textbook of Pediatrics. 22nd ed.

Foreign Body Aspiration in Children – A Case Series

- International edition. Philadelphia: Elsevier; 2024.
2. Parameswaran N, Das S, Biswal N. Respiratory Morbidity Following Foreign Body Aspiration among South Indian Children: A Descriptive Study. *Cureus*. 2018 Nov 23;10(11):e3629. DOI: 10.7759/cureus.3629.
 3. Sai Akhil R, Priya TG, Behera BK, Biswal B, Swain SK, Rath D, Mohanty MD, Choudhury J. Clinico-Radiological Profile and Outcome of Airway Foreign Body Aspiration in Children: A Single-Center Experience From a Tertiary Care Center in Eastern India. *Cureus*. 2022 Feb 13;14(2):e22163. DOI: 10.7759/cureus.22163.
 4. Hegde SV, Hui PK, Lee EY. Tracheobronchial foreign bodies in children: imaging assessment. *Semin Ultrasound CT MR*. 2015 Feb;36(1):8-20. DOI: 10.1053/j.sult.2014.10.001.
 5. Goyal, S., Jain, S., Rai, G. *et al*. Clinical variables responsible for early and late diagnosis of foreign body aspiration in pediatrics age group. *J Cardiothorac Surg* **15**, 271 (2020). <https://doi.org/10.1186/s13019-020-01314-9>
 6. Tokar B, Ozkan R, Ilhan H. Tracheobronchial foreign bodies in children: importance of accurate history and plain chest radiography in delayed presentation. *Clin Radiol*. 2004 Jul; 59(7):609-15. DOI: 10.1016/j.crad.2004.01.006.
 7. Liu B, Ding, F, An, Y, Li Y, Pan Z, Wang G, *et al*. Occult foreign body aspirations in pediatric patients: 20-years of experience. *BMC Pulm Med* **20**, 320 (2020). <https://doi.org/10.1186/s12890-020-01356-8>
 8. Çelikkaya ME, Atıcı A, Korkmaz İ, Karadağ M, El Ç, Akçora B. Negative Bronchoscopy or Computed Tomography Radiation in Children with Suspected Foreign Body Aspiration? Pros and Cons. *Tomography*. 2025 Feb 14; 11(2):17. DOI: 10.3390/tomography11020017.
 9. Jones H, Alshammasi R, Ryan S, Bowden L, Moran A, Smyth AE, Heffernan CB. Ultra-low dose CT for suspected paediatric foreign body aspiration: comparison with conventional radiograph. *J Laryngol Otol*. 2025 Dec; 139(12):1190-1195. DOI: 10.1017/S0022215125103356.
 10. Qiu W, Wu L, Chen Z. Foreign body aspiration in children with negative multi-detector Computed Tomography results: Own experience during 2011-2018. *Int J Pediatr Otorhinolaryngol*. 2019 Sep; 124:90-93. DOI: 10.1016/j.ijporl.2019.05.031.
 11. Sautin A, Marakhouski K, Pataleta A, Sanfirau K. Flexible bronchoscopy for foreign body aspiration in children: A single-centre experience. *World J Clin Pediatr*. 2024 Jun 9;13(2):91275. DOI: <https://dx.doi.org/10.5409/wjcp.v13.i2.91275>
 12. Lima E, Espindola BF, Morais IO, Scordamaglio PR, Rodrigues AJ. Flexible bronchoscopy: the first-choice method of removing foreign bodies from the airways of children. *J Bras Pneumol*. 2022 Mar 14; 48(1):e20210387. DOI: 10.36416/1806-3756/e20210387.
 13. Ünal G, Yılmaz Aİ, Tok T, Pekcan S. Use of Flexible Bronchoscopy in Foreign Body Aspiration. *Turk Arch Otorhinolaryngol*. 2022 Jun; 60(2):88-94. DOI: 10.4274/tao.2022.2022-6-6.
 14. Li D, Yang N, Dong J, Wang J, Cui X, Yu H, Liu Y. Comparison of rigid bronchoscopy and flexible bronchoscopy for the management of foreign body aspiration in children. *Am J Otolaryngol*. 2024 Mar-Apr; 45(2):104092. DOI: 10.1016/j.amjoto.2023.104092.
 15. Safia A, Abd Elhadi U, Bader R, Khater A, Karam M, Bishara T, et al. Flexible versus Rigid Bronchoscopy for Tracheobronchial Foreign Body Removal in Children. A Comparative Systematic Review and Meta-Analysis. *J Clin Med* 2024 Sep 23;

Foreign Body Aspiration in Children – A Case Series

13(18):5652.

DOI:

<https://doi.org/10.3390/jcm13185652>