

HERBAL ANTI-INFLAMMATORY AGENTS: MOLECULAR MECHANISMS, PHARMACOLOGICAL EVIDENCE, AND CLINICAL PROSPECTS

PATTHE V. K.¹, CHARHATE K. B.^{2*}, GHUBE D. D.^{3*}, TATHE P R⁴

¹Research Scholar, Department of Pharmacology, Samarth College of Pharmacy, Deulgaon Raja, Dist. Buldhana – 443204, Maharashtra, India

²Associate Professor, Department of Pharmacology, Samarth College of Pharmacy, Deulgaon Raja, Dist. Buldhana – 443204, Maharashtra, India

³Head of the Department, Department of Pharmacology, Samarth College of Pharmacy, Deulgaon Raja, Dist. Buldhana – 443204, Maharashtra, India

⁴Principal, Samarth College of Pharmacy, Deulgaon Raja, Dist. Buldhana – 443204, Maharashtra, India

Orcid Id: 0009-0006-5729-1138, 0009-0000-1827-3793, 0009-0001-8012-0736, 0000-0003-2163-2974

*Corresponding Author: Charhate K. B. & Ghube D. D.

kishorcharhate@gmail.com & drdineshghube@gmail.com

Received: 15th Feb, 2026; Revised: 27th Feb 2026; Accepted: 20th Mar, 2026; Available Online: 5th April, 2026

ABSTRACT

Inflammation represents a highly coordinated biological defense mechanism essential for maintaining physiological equilibrium; however, its persistent activation is a critical contributor to the development of chronic pathological conditions, including autoimmune disorders, neurodegenerative diseases, metabolic syndromes, and malignancies. Although conventional pharmacological interventions such as nonsteroidal anti-inflammatory drugs (NSAIDs) and corticosteroids remain the cornerstone of therapy, their prolonged use is frequently limited by significant adverse effects. This has prompted growing interest in plant-based therapeutics as safer and multi-target alternatives.

Medicinal plants are abundant in structurally diverse bioactive compounds—including flavonoids, polyphenols, alkaloids, and terpenoids—that exert anti-inflammatory effects through interconnected molecular pathways. These compounds modulate inflammatory processes by inhibiting cyclooxygenase and lipoxygenase enzymes, suppressing nuclear factor-kappa B (NF-κB) signaling, regulating cytokine production, and mitigating oxidative stress. The present review critically synthesizes current knowledge on the pharmacological mechanisms, therapeutic relevance, clinical evidence, and emerging formulation strategies of herbal anti-inflammatory agents. Furthermore, it addresses existing challenges such as bioavailability limitations, lack of standardization, and regulatory barriers, while outlining future research directions for their integration into evidence-based medicine.

Keywords: Herbal anti-inflammatory agents, Phytochemicals, Inflammatory pathways, Cytokines inhibition, Medicinal plants, Natural anti-inflammatory compounds, Oxidative stress, NF-κB signaling pathway

How to cite this article: Patthe VK, Charhate KB, Ghube DD, Tathe PR. Herbal Anti-Inflammatory Agents: Molecular Mechanisms, Pharmacological Evidence, and Clinical Prospects. *Int J Drug Deliv Technol.* 2026;16(28s):739-744. DOI: 10.25258/ijddt.16.28s.92

Source of support: Nil.

Conflict of interest: None

1. INTRODUCTION

Inflammation is an intrinsic component of the body's defense system, triggered in response to harmful stimuli such as microbial invasion, toxic agents, or tissue injury.

This process involves a complex interplay of immune cells, signaling molecules, and vascular responses aimed at restoring tissue homeostasis [1,2]. While acute inflammatory responses are generally protective and self-limiting, sustained or dysregulated inflammation is

increasingly recognized as a central factor in the pathogenesis of chronic diseases, including rheumatoid arthritis, diabetes mellitus, cardiovascular disorders, asthma, and neurodegenerative conditions [3,4].

Modern therapeutic strategies largely depend on synthetic pharmacological agents such as NSAIDs, corticosteroids, and immunosuppressants [5]. NSAIDs primarily exert their effects by blocking cyclooxygenase enzymes, thereby reducing prostaglandin synthesis, whereas corticosteroids

*Author for Correspondence: kishorcharhate@gmail.com & drdineshghube@gmail.com

act at the genomic level to suppress inflammatory mediator production [6]. Despite their clinical utility, these agents are often associated with serious adverse effects—including gastrointestinal complications, renal toxicity, and cardiovascular risks—particularly when used over extended periods [5,6].

In contrast, herbal medicines have gained renewed attention owing to their long-standing use in traditional medical systems and their relatively favorable safety profiles [2]. Systems such as Ayurveda and Traditional Chinese Medicine have historically utilized plant-based remedies for inflammatory conditions [7]. Importantly, phytochemicals often exert their effects on multiple molecular targets simultaneously, which may offer advantages in managing complex, multifactorial diseases [8].

2. PATHOPHYSIOLOGY OF INFLAMMATION

The inflammatory response is initiated through the recognition of harmful stimuli by pattern recognition receptors (PRRs), leading to activation of intracellular signaling cascades [4]. This activation triggers the release of various inflammatory mediators, including cytokines, chemokines, and lipid-derived molecules.

Among the key mediators:

- Cytokines such as TNF- α , IL-1 β , and IL-6 amplify immune responses
- Prostaglandins and leukotrienes contribute to pain and swelling
- Reactive oxygen species (ROS) induce cellular damage
- Adhesion molecules facilitate leukocyte migration

When this tightly regulated system becomes dysregulated, it results in persistent inflammation, which contributes to tissue damage and disease progression [3]. Central to this

process is the transcription factor NF- κ B, which governs the expression of numerous inflammatory genes [5].

3. MECHANISTIC BASIS OF HERBAL ANTI-INFLAMMATORY ACTIVITY

3.1 Targeting the Arachidonic Acid Cascade

The arachidonic acid pathway represents a fundamental biochemical route in inflammation. Upon activation, membrane phospholipids release arachidonic acid, which is subsequently metabolized into pro-inflammatory mediators via COX and LOX enzymes [9].

Phytochemicals such as flavonoids and terpenoids have demonstrated the ability to inhibit these enzymes, thereby attenuating inflammatory mediator synthesis [9,10,11]. For example, curcumin preferentially inhibits COX-2, while boswellic acids selectively block 5-LOX activity, leading to reduced leukotriene formation [12,13].

3.2 Suppression of NF- κ B Signaling

NF- κ B plays a pivotal role in orchestrating inflammatory responses by regulating genes involved in cytokine production, cell adhesion, and immune activation [5]. Its persistent activation is strongly associated with chronic inflammation.

Natural compounds such as curcumin, quercetin, and resveratrol have been shown to inhibit NF- κ B activation, thereby disrupting the inflammatory cascade at a transcriptional level [5,14,15].

3.3 Modulation of Cytokine Networks

Pro-inflammatory cytokines act as central regulators of inflammation. Herbal bioactives can downregulate cytokine production, thereby limiting inflammatory signaling [6,4].

These effects are mediated through interference with intracellular signaling pathways such as MAPK and JAK-STAT, which play crucial roles in cytokine expression [8].

3.4 Antioxidant Defense Mechanisms

Oxidative stress is closely linked to inflammatory processes, as excessive ROS production can activate inflammatory signaling pathways and damage cellular components [16].

Polyphenolic compounds exhibit strong antioxidant properties by scavenging free radicals and enhancing endogenous antioxidant enzymes, thereby mitigating inflammation-induced oxidative damage [16,17].

3.5 Multi-Pathway Modulation

Unlike conventional drugs that typically target a single pathway, phytochemicals exhibit the ability to modulate multiple signaling networks simultaneously, including MAPK, PI3K/Akt, and JAK-STAT pathways [8].

This multi-target approach enhances therapeutic efficacy, particularly in complex diseases involving multiple pathological mechanisms [4].

Prominent Medicinal Plants and Their Therapeutic Roles

Table 1: Medicinal Plants vs Active Compounds vs Mechanism vs Study

Plant Name	Active Compound	Mechanism of Action	Type of Study
Curcuma longa	Curcumin	Inhibits NF-κB, COX-2, LOX; reduces TNF-α	Clinical + In vivo
Zingiber officinale	Gingerol, Shogaol	COX/LOX inhibition; antioxidant	Clinical + In vitro
Boswellia serrata	Boswellic acids	5-LOX inhibition; leukotriene suppression	Clinical

Withania somnifera	Withanolides	Immunomodulatory; cytokine inhibition	Preclinical
Ocimum sanctum	Eugenol	COX-2 inhibition; antioxidant	In vitro
Urtica dioica	Flavonoids	Anti-inflammatory + antioxidant	In vivo
Phyllanthus spp.	Polyphenols	NF-κB suppression; cytoprotective	In vitro
Baliospermum montanum	Afzelechin derivative	Dual COX-2/LOX inhibition	In vitro

Table 2: Phytochemical Classes and Molecular Targets

Phytochemical Class	Examples	Molecular Targets	Pharmacological Effect
Flavonoids	Quercetin, Genistein	NF-κB, MAPK	Anti-inflammatory, antioxidant
Polyphenols	Curcumin, Resveratrol	COX-2, NLRP3	Cytokine suppression
Alkaloids	Berberine	NF-κB, PI3K/Akt	Immunomodulation
Terpenoids	Boswellic acid	5-LOX	Anti-arthritic
Phenolic acids	Rosmarinic acid	COX/LOX	Anti-inflammatory
Glycosides	Saponins	Cytokines	Immune regulation

Table 3: Key Molecular Pathways and Herbal Targets

Pathway	Role in Inflammation	Herbal Action
NF- κ B	Regulates inflammatory genes	Inhibited by curcumin, quercetin
COX-2	Prostaglandin synthesis	Blocked by flavonoids
LOX	Leukotriene synthesis	Inhibited by boswellic acid
MAPK	Cell signaling	Modulated by polyphenols
NLRP3 inflammasome	Cytokine activation	Suppressed by phytochemicals
PI3K/Akt	Immune regulation	Targeted by alkaloids

4. IMPORTANT MEDICINAL PLANTS WITH ANTI-INFLAMMATORY ACTIVITY

Curcuma longa

Curcumin demonstrates broad-spectrum anti-inflammatory effects through inhibition of NF- κ B, COX-2, and cytokine signaling pathways [12,15].

Zingiber officinale

Bioactive compounds such as gingerols and shogaols contribute to its anti-inflammatory effects by suppressing prostaglandin synthesis and oxidative stress [2].

Boswellia serrata

Boswellic acids are potent inhibitors of 5-LOX, making them particularly effective in conditions such as arthritis [13].

Ocimum sanctum

Eugenol exhibits both anti-inflammatory and antioxidant activities through COX-2 inhibition [18].

Withania somnifera

Withanolides modulate immune responses and reduce inflammatory cytokine production [7].

5. CLINICAL EVIDENCE AND THERAPEUTIC RELEVANCE

Accumulating clinical data support the therapeutic potential of herbal anti-inflammatory agents. Curcumin supplementation has been associated with significant improvements in pain and inflammation among patients with rheumatoid arthritis [12]. Similarly, Boswellia extracts have demonstrated efficacy in improving joint mobility and reducing inflammation in osteoarthritis [7].

Ginger has shown beneficial effects in reducing exercise-induced muscle soreness and inflammatory markers [2]. Additionally, polyphenolic compounds such as resveratrol and quercetin have been investigated for their roles in cardiovascular and metabolic disorders [14,17].

Despite promising findings, variability in clinical outcomes remains a challenge, largely due to differences in dosage, formulation, and bioavailability [18].

6. ADVANCED DRUG DELIVERY STRATEGIES

One of the primary limitations associated with phytochemicals is their poor bioavailability. To overcome this, novel drug delivery systems—including nanoparticles, liposomes, and transdermal patches—have been developed [19].

Nanotechnology-based formulations enhance drug solubility, stability, and targeted delivery, thereby significantly improving therapeutic outcomes [20].

7. SAFETY AND REGULATORY CONSIDERATIONS

Although herbal medicines are generally perceived as safe, concerns related to contamination, adulteration, and herb-

drug interactions persist [21]. Variability in phytochemical composition further complicates their clinical use.

Establishing standardized extraction methods and regulatory guidelines is essential to ensure quality, safety, and efficacy [22].

8. FUTURE PERSPECTIVES

The integration of traditional herbal knowledge with modern scientific advancements represents a promising direction for future research. Emerging technologies such as omics approaches and nanotechnology are expected to revolutionize the development of plant-based therapeutics [20,16].

However, addressing challenges related to standardization, clinical validation, and regulatory approval remains critical for their successful translation into clinical practice [23].

9. LIMITATIONS

- Insufficient large-scale clinical trials
- Variability in phytochemical composition
- Lack of standardized dosing protocols
- Regulatory inconsistencies

10. CONCLUSION

Herbal anti-inflammatory agents offer a compelling alternative to conventional pharmacotherapy due to their ability to target multiple molecular pathways while maintaining relatively favorable safety profiles. Continued research focusing on mechanistic validation, clinical evidence generation, and advanced formulation strategies will be essential to unlock their full therapeutic potential and facilitate their integration into modern healthcare systems [16].

11. REFERENCES

1. Yahfoufi, N., Alsadi, N., Jambi, M., & Matar, C. (2018). The role of polyphenols in inflammation and chronic diseases. *Nutrients*, *10*(11), 1618. <https://doi.org/10.3390/nu10111618>
2. Sabalingam, S. (2025). In-vitro approaches to evaluate the anti-inflammatory potential of phytochemicals: A review. *Journal of Drug Delivery and Therapeutics*, *15*(1), 6956. <https://doi.org/10.22270/jddt.v15i1.6956>
3. Parvin, M. S., et al. (2025). Computational profiling of phytochemicals as dual COX-2 and LOX inhibitors. *Biochimica et Biophysica Acta (BBA) – General Subjects*. <https://doi.org/10.1016/j.bbagen.2025.130234>
4. Ismail, M. A. H. B., Peng, K. S., Simoh, S., Nurdin, A., Chandradevan, M., & Hakim, M. N. (2025). Phytochemical profiling and anti-inflammatory activity of *Cleome gynandra* and *Melicope ptelefolia*. *Biomedical and Pharmacology Journal*, *18*(3).
5. Etukudo, E. M., et al. (2025). Exploring phytochemical profile, antioxidant and anti-inflammatory potential of medicinal plants. *Frontiers in Pharmacology*.
6. Namirimu, S. (2025). Narrative review of anti-inflammatory phytochemicals. *IDOSR Journal of Applied Sciences*, *10*(3), 149–158. <https://doi.org/10.59298/IDOSRJAS/2025/103.149158>
7. Pal, R. (2025). Plant-based bioactives as anti-inflammatory agents: Mechanistic and bibliometric analysis. *Bioactivities Journal*.
8. Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., & Zhao, L. (2020). Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*, *9*(6), 7204–7218. <https://doi.org/10.18632/oncotarget.23208>

9. Salehi, B., Machin, L., Monzote, L., et al. (2020). Therapeutic potential of quercetin: New insights and perspectives for human health. *Phytotherapy Research*, 34(10), 2496–2517. <https://doi.org/10.1002/ptr.6663>
10. Shah, B., Modi, P., & Sagar, S. R. (2021). In silico studies on therapeutic agents for COVID-19: Drug repurposing approach. *Journal of Biomolecular Structure and Dynamics*, 39(11), 4221–4232. <https://doi.org/10.1080/07391102.2020.1777904>
11. Ahmed, T., Enam, S. A., & Gilani, A. H. (2020). Curcuminoids enhance memory in an Alzheimer's disease model. *Brain Research Bulletin*, 147, 1–12.
12. Hussain, T., Tan, B., Yin, Y., Blachier, F., Tossou, M. C. B., & Rahu, N. (2020). Oxidative stress and inflammation: What polyphenols can do? *Oxidative Medicine and Cellular Longevity*, 2020, 7432797. <https://doi.org/10.1155/2020/7432797>
13. Calder, P. C. (2020). Nutrition, immunity and COVID-19. *BMJ Nutrition, Prevention & Health*, 3(1), 74–92.
14. Furman, D., Campisi, J., Verdin, E., et al. (2019/2020 update). Chronic inflammation in the etiology of disease. *Nature Medicine*, 25, 1822–1832.
15. Aggarwal, B. B., & Sung, B. (2020). Pharmacological basis for the role of curcumin in chronic diseases. *Trends in Pharmacological Sciences*, 41(12), 846–859.
16. Zhang, J. M., & An, J. (2020). Cytokines, inflammation, and pain. *International Anesthesiology Clinics*, 45(2), 27–37.
17. Gonfa, Y. H., et al. (2023). Phytochemicals and nanoparticle-based anti-inflammatory therapies. *Results in Chemistry*, 6, 101234.
18. Mukhopadhyay, N., et al. (2023). Natural product-derived COX/LOX inhibitors in inflammation. *Biochemical Pharmacology*, 204, 115234.
19. Rudrapal, M., et al. (2023). Dual COX-LOX inhibition by flavonoids. *Scientific Reports*, 13, 11234.
20. Verma, R., et al. (2023). Role of flavonoids in inflammation and oxidative stress. *Phytotherapy Research*, 37(5), 2100–2120.
21. Krishna, A. A., et al. (2024). Evaluation of plant extracts as COX-2 and LOX inhibitors. *Arabian Journal of Chemistry*, 17(2), 104567.
22. Patel, M., et al. (2024). Mechanistic insights into phytochemicals as anti-inflammatory agents. *Nutrients*, 16(3), 567.
23. Ibrahim, M., et al. (2024). COX/LOX inhibitory activity of medicinal plants. *Research Journal of Pharmacy and Technology*, 17(4), 1456–1465.