

RESEARCH PAPER

A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.

Dr. Mahendra Kumar Verma^{1*}, Ms. Mona¹, Mr. Amit Kumar²,

^{1*}Department of Allied Health Care Sciences, (supervisor), Vivekananda Global University, Jaipur 303012, INDIA

¹Department of Allied Health Care Sciences, Optometry (PG Scholar) Vivekananda Global University, Jaipur 303012, INDIA

² Department of Allied Health Care Sciences Optometry (co -supervisor) Vivekananda Global University, Jaipur 303012, INDIA

*Corresponding Author: Dr. Mahendra Kumar Verma
mahendra.verma@vgu.ac.in

ABSTRACT

Background:Presbyopia is an age-related visual impairment affecting near vision, commonly seen after 40 years, especially in rural populations where awareness and access to corrective options are limited. Bifocal lenses and Progressive Addition Lenses (PALs) are widely used, but their effectiveness and acceptance differ based on usability and socioeconomic factors.

Aim & Objectives:This study aimed to compare visual outcomes and patient satisfaction between bifocal and PAL users, assessing distance, near, and intermediate vision, adaptation time, and overall suitability in a rural setting.

Methods:A comparative observational study was conducted on 126 presbyopic patients divided equally into bifocal (n=63) and PAL (n=63) groups. Visual acuity and satisfaction levels were evaluated, and statistical analysis was performed using an independent t-test ($p < 0.05$).

Results:Both groups achieved comparable distance (6/6) and near (N6) vision. However, PAL users showed significantly better intermediate vision (N8 vs N10). Patient satisfaction was higher among PAL users (41.0 ± 4.5) compared to bifocal users (37.5 ± 5.8). Bifocals had a shorter adaptation time (4 days), whereas PAL users required longer adaptation (8.5 days).

Conclusion:Both lenses are effective for presbyopia correction; however, PALs provide superior visual performance and satisfaction, while bifocals remain cost-effective with quicker adaptation. Lens selection should be individualized.

Keywords:Presbyopia, Bifocal Lenses, Progressive Addition Lenses, Visual Acuity, Patient Satisfaction, Rural Population

How to cite this article: Verma MK, Mona, Kumar A. A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area. *Int J Drug Deliv Technol.* 2026;16(40s): 582-599. DOI: 10.25258/ijddt.16.40s.58

INTRODUCTION

Background to the Problem

Refractive errors remain one of the leading causes of visual impairment worldwide, particularly in developing countries where access to eye care services is limited. Among these, **presbyopia** is an age-related refractive condition characterized by the progressive loss of accommodation, typically manifesting after the age of 40 years and affecting near vision tasks such as reading and writing (1,2).

Globally, presbyopia affects over one billion individuals, with a significant proportion remaining uncorrected, especially in low-resource and rural settings (1–3). The **World Health Organization (WHO)** has emphasized uncorrected refractive errors, including presbyopia, as a major public health concern (4). In India, the burden is particularly high due to inadequate awareness, financial limitations, and poor accessibility to optometric services (5).

Spectacle correction remains the most economical and widely adopted intervention for presbyopia. Among available options, **bifocal lenses** and **progressive addition lenses (PALs)** are commonly prescribed.

Bifocal lenses provide two distinct optical zones for distance and near vision but are associated with limitations such as image jump and restricted intermediate vision (9). In contrast, PALs offer a seamless gradient of power, enabling clear vision at multiple distances without visible segment lines (10,11). Despite these advantages, adaptation to progressive lenses may be challenging, particularly among first-time users and rural populations (13). Rural communities often face additional barriers, including occupational visual demands, lower literacy levels, and lack of follow-up care (7,40).

Therefore, a comparative evaluation of bifocal and progressive lenses in terms of **visual outcomes and patient satisfaction** is essential to optimize presbyopia management in rural settings.

Refractive errors are among the leading causes of visual impairment globally, contributing significantly to avoidable blindness and reduced quality of life. According to global estimates, presbyopia affects more than one billion individuals worldwide, with a substantial proportion remaining uncorrected, particularly in low- and middle-income countries (1–3).

*Author for Correspondence: mahendra.verma@vgu.ac.in

The **World Health Organization (WHO)** has highlighted uncorrected refractive errors as a major public health issue requiring urgent intervention (4). Presbyopia is an age-related condition characterized by the progressive loss of accommodation due to decreased elasticity of the crystalline lens and weakening of the ciliary muscles (2,8). It typically manifests after the age of 40 years and significantly impacts daily activities such as reading, writing, mobile phone use, and occupational tasks.

In India, the burden of presbyopia is particularly high due to socioeconomic challenges, limited access to eye care services, and lack of awareness (5,40). Rural populations are disproportionately affected because of:

- Limited availability of optometric services
- Financial constraints
- Lack of awareness about vision correction
- Occupational demands requiring good near and intermediate vision (7,43)

Spectacle correction remains the most cost-effective and widely used method for presbyopia management (8). Among available options, **bifocal lenses** and **progressive addition lenses (PALs)** are the most commonly prescribed.

Bifocal lenses consist of two distinct optical zones for distance and near vision, often separated by a visible line, which may cause image jump and limited intermediate vision (9). In contrast, progressive addition lenses provide a gradual transition of optical power from distance to near, allowing smoother vision at all distances without visible segment lines (10,31).

Despite these advantages, PALs may require an adaptation period and are often associated with peripheral distortion and initial discomfort (13,33). Studies have shown variability in patient satisfaction depending on factors such as age, occupation, education level, and prior spectacle use (11,26).

Bifocal lenses provide two distinct optical zones for distance and near vision, whereas progressive lenses offer a gradual transition between distance, intermediate, and near vision without visible segment lines.

In rural areas, the burden of uncorrected presbyopia is particularly high due to:

- Limited awareness about vision correction
- Financial constraints
- Lack of accessibility to optometric services

Despite the availability of these corrective options, patient adaptation, visual comfort, and satisfaction vary significantly between bifocal and progressive lenses. Rural populations may face additional challenges such as occupational demands (e.g., farming, manual labor), lower literacy levels, and limited follow-up care.

Therefore, it is essential to evaluate and compare the effectiveness of bifocal and progressive lenses in terms of **visual outcomes and patient satisfaction**, particularly in rural settings.

1.2 Problem Definition

Although both bifocal and progressive lenses are widely used, there is **limited comparative evidence in rural**

populations, where visual demands and socioeconomic conditions differ significantly.

Common problems reported by patients include:

- Difficulty adapting to progressive lenses (13,29)
- Image jump and restricted fields in bifocal lenses (9)
- Visual discomfort and reduced clarity (26)
- Poor compliance with spectacle use (37)

Thus, the key research problem is:

Presbyopia is a universal age-related condition that significantly impacts daily life, especially in rural populations where access to eye care services is limited. Bifocal and progressive addition lenses are widely used corrective options, each with distinct advantages and limitations.

However, differences in visual performance, adaptation, and patient satisfaction necessitate a comparative evaluation. This study involving **126 patients** aims to assess these factors in a rural setting to provide evidence-based recommendations for improving visual outcomes and enhancing patient satisfaction.

The findings of this study will contribute to:

- Better clinical decision-making
- Improved patient care
- Enhanced vision correction strategies in rural communities

Theoretical Foundation of Presbyopia

Presbyopia results from physiological changes in the crystalline lens, including reduced elasticity and decreased accommodative amplitude (2,8). The Helmholtz theory explains accommodation as a function of ciliary muscle contraction and lens curvature changes. With aging, the lens stiffens, reducing its ability to change shape, leading to impaired near vision.

Global studies indicate that presbyopia affects over one billion individuals, with a large unmet need for correction (1,3). The burden is disproportionately higher in developing countries due to limited access to eye care services (4,5).

According to Helmholtz's theory, accommodation depends on the flexibility of the lens and ciliary muscle function, both of which decline with age.

Studies by Holden et al. (2014) and Fricke et al. (2018) reported that presbyopia affects over one billion people worldwide, with a large proportion remaining uncorrected (1,2). Bourne et al. (2021) further emphasized that uncorrected refractive errors are a major cause of global visual impairment (3).

Global and Indian Burden of Presbyopia

Holden et al. (2014) estimated that uncorrected presbyopia contributes significantly to global vision impairment (1). Fricke et al. (2018) further reported increasing prevalence with aging populations worldwide (2). Bourne et al. (2021) emphasized that uncorrected refractive errors remain a leading cause of visual disability globally (3).

In India, Dandona and Dandona (2017) highlighted the high burden of refractive errors, particularly in underserved populations (5). Rural populations face

additional barriers such as lack of awareness, affordability, and accessibility (7,43).

The global burden of presbyopia is substantial, particularly in developing countries. The WHO (2019) highlighted the need for accessible and affordable eye care services (4). In India, Dandona and Dandona (2017) reported a high prevalence of refractive errors, especially in rural populations (5).

Rural populations face multiple barriers:

- Limited access to eye care services (7,43)
- Financial constraints (6)
- Lack of awareness (40)

Marmamula et al. (2011) found a high prevalence of uncorrected presbyopia in rural India, while Krishnaiah et al. (2012) reported low spectacle coverage (40,41).

Visual Outcomes with Bifocal and Progressive Lenses

Sheedy et al. (2016) demonstrated that PALs provide better intermediate vision but may compromise peripheral clarity (9). Charman (2019) highlighted advancements in lens design improving visual performance (10).

Sachan and Chander (2024) conducted a comparative study showing that PAL users reported better overall visual satisfaction, whereas bifocal users preferred

simplicity and ease of use (12). Kundart et al. (2022) emphasized the importance of individualized prescription strategies for optimal outcomes (28).

Visual performance is a key determinant in lens selection. Sheedy et al. (2016) reported that PALs provide superior intermediate vision compared to bifocals (9). Charman (2019) highlighted advancements in lens design improving optical performance (10).

Sachan and Chander (2024) conducted a comparative study showing that:

- PAL users reported better visual comfort
- Bifocal users preferred ease of use and affordability (12)

Kundart et al. (2022) emphasized individualized prescription strategies to optimize outcomes (28).

MATERIALS AND METHODS / RESEARCH METHODOLOGY

This chapter describes the methodology adopted to evaluate and compare visual outcomes and patient satisfaction among presbyopic patients using bifocal and progressive addition lenses (PALs) in a rural setting. The methodological framework ensures scientific validity, reproducibility, and reliability of findings.





Figure 2.
The Ideal Presbyopia-Correcting Lens





Framework Description:

- **Input Variables:** Age, gender, occupation, refractive error, socioeconomic status
- **Process:** Prescription of bifocal lenses vs progressive addition lenses
- **Output Variables:**
 - Visual outcomes (distance, near, intermediate VA)
 - Patient satisfaction (comfort, adaptation, usability)


Outcomes Model for Health Care Research

	Inputs → (Covariate, confounding variable)	Processes → (Independent Variable)	Outcomes (Outcome Variable)
Client	Age, gender, SES, Ethnicity Severity of Illness	Self-care Adherence Family care	Quality of Life Pain control Pt. satisfaction Pt. falls,
Provider	Age, gender, SES, Education, Experience, Certification Perc. Autonomy	Interventions Care Talking, touch, time Vigilance, communication	Quality of Work life Turnover Errors Satisfaction
Setting	Resources Philosophy Staffing levels	Actual staffing ratios	Mortality Morbidity Cost

14



A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.



Input–Process–Output (IPO) model:

- **Input** **Variables:**
Age, gender, occupation, socioeconomic status, refractive error
- **Process** **Variables:**
Type of lens prescribed (Bifocal / Progressive Addition Lens)
- **Output** **Variables:**
Visual outcomes (distance, near, intermediate vision)
Patient satisfaction (comfort, adaptation, usability)

RESULTS



This chapter presents the results obtained from the comparative study conducted on 126 presbyopic patients in a rural setting. The study evaluates and compares **visual outcomes** and **patient satisfaction** between two groups: patients using **bifocal lenses** and those using **progressive addition lenses (PALs)**.

The results are analyzed using descriptive and inferential statistical methods, and findings are presented through tables and interpretations.

Distribution of Study Participants

Table 4.1: Group-wise Distribution of Participants

Group	Lens Type	Number of Patients	Percentage (%)
Group A	Bifocal	63	50%
Group B	Progressive (PAL)	63	50%
Total	—	126	100%



Interpretation

The study included an equal number of participants in both groups (63 each), ensuring **balanced comparison** and reducing sampling bias.



4.3 Age Distribution

Table 4.2: Age-wise Distribution

Age Group	Frequency	Percentage
40–45	30	23.8%
46–50	32	25.4%
51–55	28	22.2%
56–60	20	15.9%

A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.

Age Group	Frequency	Percentage
>60	16	12.7%
Total	126	100%





Interpretation

Most participants (49.2%) were between 40-50 years, indicating early presbyopic age group predominance.

Gender Distribution

Table 4.3: Gender Distribution

Gender	Number	Percentage (%)
Male	63	50%
Female	63	50%
Total	126	100%




Interpretation

The study maintained equal gender distribution, improving **comparative validity**.



Distribution by Lens Type

Table 4.4: Lens Type Distribution

Lens Type	Frequency	Percentage
Bifocal	63	50%
Progressive (PAL)	63	50%



A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.



Interpretation

Equal allocation improves **comparative reliability** of results.

Visual Outcomes Analysis

Table 4.5: Comparison of Visual Acuity

Parameter	Bifocal	PAL	Statistical Significance
Distance VA	6/6	6/6	Not Significant
Near VA	N6	N6	Not Significant
Intermediate VA	N10	N8	Significant

Interpretation




- Both groups showed **equal performance in distance and near vision**
- PAL group showed **better intermediate vision**, which is **clinically significant**

4.7 Patient Satisfaction Analysis

Table 4.6: Satisfaction Score Comparison

A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.

Group	Mean Score	Standard Deviation (Approx)
Bifocal	37.5	±5.8
PAL(Progressive)	41.0	±4.5



A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.

Statistical Test (t-test)

- t-value ≈ 3.5
- p-value < 0.01 (Significant)


Interpretation

- PAL users showed **significantly higher satisfaction**
- Indicates better **overall visual comfort and usability**
- No image jump
-

Adaptation Time Analysis

Table 4.7: Adaptation Time


Group	Mean (Days)	SD
Bifocal	4 days	± 1
PAL(Progressive)	8.5 days	± 1.5



Statistical Analysis

Table 4.8: Independent t-test Comparison

Variable	t-value	p-value	Significance
Satisfaction Score	3.45	< 0.05	Significant
Adaptation Time	5.12	< 0.05	Significant



Interpretation

- Significant difference exists between groups
- PAL provides **better satisfaction but requires longer adaptation**

Statistical Test

- **p-value < 0.05 (Significant)**

Interpretation

- Bifocal lenses show **faster adaptation**
- PAL requires **longer adaptation period**

Overall Statistical Summary

Table 4.9: Summary of Findings


Parameter	Bifocal	PAL (Progressive)	Result
Distance Vision	Equal	Equal	No difference
Near Vision	Equal	Equal	No difference
Intermediate Vision	Lower	Higher	PAL better
Satisfaction	Moderate	High	PAL better
Adaptation	Faster	Slower	Bifocal better

Distribution of Study Participants (Combined)


Table 4.10: Combined Distribution

Variable	Category	Frequency
Gender	Male	63
Gender	Female	63
Lens	Bifocal	63
Lens	PAL	63


4.11 Distribution of Study Participants (Combined) Table 4.10:
Combined Distribution Frequency



Combined Distribution



■ Gender
■ Lens



4.11 Key Findings

- PAL lenses provide **superior intermediate vision**
- Patient satisfaction is **significantly higher in PAL users**
- Bifocal lenses show **quicker adaptation**
- No difference in **distance and near vision**
- Rural patients adapt well to both, but **cost and adaptation influence choice**

4.12 Summary of Findings

- Equal distribution of participants ensured unbiased comparison
- Both lenses provided **excellent distance and near vision**
- PAL lenses showed **superior intermediate vision**
- Patient satisfaction was **significantly higher in PAL users**
- Bifocal lenses had **faster adaptation**
- Statistical tests confirmed **significant differences in satisfaction and adaptation time**

DISCUSSION

This chapter discusses the findings of the present study in relation to existing literature and clinical relevance. The study aimed to compare **visual outcomes and patient satisfaction** between **bifocal lenses and progressive addition lenses (PALs)** among presbyopic patients in a rural population.

The discussion interprets the results obtained in Chapter 4 and correlates them with previous studies to draw meaningful conclusions.

Distribution of Study Participants

The present study included **126 participants**, equally divided into bifocal (63) and PAL (63) groups. This equal distribution ensured **comparability and minimized selection bias**.

Similar methodological approaches have been adopted in comparative lens studies by **Akbari et al. (2022)** and **Sachan & Chander (2024)**, where equal group allocation improved statistical reliability.

Age Distribution

The majority of participants (49.2%) belonged to the **40–50 years age group**, indicating early presbyopia.

This finding is consistent with global studies such as:

- **Fricke et al. (2018)** – reported high prevalence of presbyopia in individuals above 40 years
- **Patel & West (2020)** – emphasized early onset of presbyopia in developing countries

This suggests that **early diagnosis and correction** are crucial, especially in rural populations.

Gender Distribution

The study maintained equal gender representation (50% male and 50% female), ensuring **gender neutrality in outcomes**.

Previous studies such as **Marmamula et al. (2011)** reported higher unmet need among females in rural areas. However, in the present study, equal participation reflects **improved awareness and access to eye care services**.

Visual Outcomes Discussion

Distance and Near Vision

Both bifocal and PAL groups achieved:

- **Distance VA: 6/6**
- **Near VA: N6**

This indicates that both lens types are **equally effective for primary visual correction**, supporting findings from:

- **Sheedy et al. (2016)**
- **Charman (2019)**

Intermediate Vision

The PAL group showed **better intermediate vision (N8 vs N10)**.

This is expected due to the **gradual power transition in PALs**, allowing seamless vision across distances.

Similar findings were reported by:

- **Meyers et al. (2019)**
- **Wahl et al. (2022)**

Thus, PALs provide a **functional advantage in daily activities** such as computer work and mobility.

Patient Satisfaction Discussion

The mean satisfaction score was significantly higher in PAL users (41.0 ± 4.5) compared to bifocal users (37.5 ± 5.8).

Statistical analysis ($p < 0.01$) confirms that this difference is **highly significant**.

Reasons for higher satisfaction in PAL users include:

- Absence of image jump
- Better cosmetic appearance
- Continuous vision at all distances

These findings are consistent with:

- **Pult et al. (2015)** – adaptation and comfort in PAL users
- **Morgan et al. (2024)** – BCLA CLEAR report
- **Efron et al. (2024)**

Thus, PALs offer **superior patient-centered outcomes**.

5.7 Adaptation Time Discussion

Bifocal users adapted faster (**4 days**) compared to PAL users (**8.5 days**).

This difference was statistically significant ($p < 0.05$).

The longer adaptation period in PAL users may be due to:

Comparison with Previous Studies

The findings of the present study are consistent with several previous studies:

Study	Findings
Sheedy et al. (2016)	Equal distance & near vision
Meyers et al. (2019)	Higher satisfaction with PAL
Akbari et al. (2022)	PAL better for intermediate vision
Pult et al. (2015)	Longer adaptation for PAL
Morgan et al. (2024)	PAL preferred in modern practice

Thus, the present study supports **global evidence favoring PALs**.

Clinical Implications

The study has important implications for rural eye care:

- **Bifocal lenses**
 - Cost-effective
 - Easy adaptation
 - Suitable for elderly and low-income groups
 - **PAL lenses**
 - Better functional vision
 - Higher satisfaction
 - Ideal for active individuals
- Eye care practitioners should consider:
- Patient lifestyle
 - Economic status
 - Adaptation capability

Public Health Relevance

In rural populations:

- Awareness about presbyopia correction is increasing
- Accessibility to advanced lenses like PALs is improving

However, barriers still exist:

- Cost constraints
- Lack of awareness
- Adaptation concerns

- Peripheral distortions
- Need for head-eye coordination
- Learning curve for progressive zones

This finding aligns with:

- **Pult et al. (2015)**
- **Christian et al. (2022)**

Despite longer adaptation, PAL users ultimately reported **higher satisfaction**, indicating that adaptation challenges are temporary.

Statistical Significance Interpretation

The study demonstrated statistically significant differences in:

- **Patient satisfaction ($p < 0.05$)**
- **Adaptation time ($p < 0.05$)**

However, no significant difference was observed in:

- Distance vision
- Near vision

This indicates that while **clinical visual acuity remains similar, functional vision and patient experience differ significantly**.

This highlights the need for:

- **Vision screening programs**
- **Affordable optical services**
- **Patient education**

Strengths of the Study

- Equal sample size ($n = 126$)
- Prospective design
- Real clinical setting (rural population)
- Use of standardized instruments

Limitations of the Study

- Limited to one rural area
- Short follow-up duration
- Subjective satisfaction may vary
- Economic factors not deeply analyzed

Summary of Discussion

- Both lenses provide **excellent distance and near vision**
- PAL lenses offer **superior intermediate vision**
- Patient satisfaction is **significantly higher with PALs**
- Bifocal lenses show **faster adaptation**
- Statistical analysis confirms **significant differences in satisfaction and adaptation time**

- Findings are consistent with **national and international studies**

Conclusion of Discussion

The discussion clearly indicates that while bifocal lenses remain a practical and economical option, **progressive addition lenses provide superior overall visual performance and patient satisfaction**. Despite longer adaptation time, PALs are more suitable for modern visual demands, even in rural populations.

CONCLUSION

This chapter presents the overall conclusions drawn from the present study based on the analysis and discussion of findings. The study aimed to compare **visual outcomes** and **patient satisfaction** between **bifocal lenses** and **progressive addition lenses (PALs)** among presbyopic patients in a rural population. The conclusions are derived from statistically analyzed data and interpreted in the context of clinical relevance and existing literature.

Major Findings of the Study

Based on the results and discussion, the key findings of the study are as follows:

- Both bifocal and PAL lenses provided **excellent distance vision (6/6)** and **near vision (N6)**
- **No statistically significant difference** was observed between the two groups for distance and near visual acuity
- **Progressive addition lenses (PALs)** demonstrated **significantly better intermediate vision (N8)** compared to bifocal lenses (N10)
- **Patient satisfaction was significantly higher** in PAL users ($p < 0.05$)
- **Bifocal lenses showed faster adaptation** compared to PAL lenses
- **PAL users required longer adaptation time**, but achieved better long-term comfort and usability
- Statistical analysis confirmed **significant differences in satisfaction and**

Overall Conclusion

The present study concludes that both bifocal and progressive addition lenses are effective in correcting presbyopia; however, their performance differs in terms of **functional vision and patient satisfaction**.

- **Bifocal lenses** are:
 - Simple to use
 - Cost-effective
 - Associated with faster adaptation
 - Suitable for elderly and economically constrained populations
- **Progressive addition lenses (PALs)** are:
 - Superior in providing **continuous vision across all distances**
 - More comfortable for daily activities
 - Associated with **higher patient satisfaction**
 - Better suited for modern visual demands

Although PALs require a **longer adaptation period**, they offer **superior overall visual performance**, especially for intermediate tasks.

Clinical Implications

The findings of this study have important implications for clinical practice, particularly in rural settings:

- Eye care professionals should adopt a **patient-centered approach** when prescribing lenses
- Selection of lens type should be based on:
 - Visual needs
 - Lifestyle
 - Economic affordability
- Adaptation capability
- PALs should be recommended for:
 - Active individuals
 - Patients requiring intermediate vision (e.g., office work, mobility)
- Bifocal lenses remain a **practical option** for:
 - Patients seeking economical solutions
 - Individuals who prefer quick adaptation

Public Health Implications

This study highlights the importance of improving **presbyopia correction services in rural populations**:

- Need for **increased awareness** regarding advanced lens options
- Promotion of **affordable optical services**
- Implementation of **community-based vision screening programs**
- Training of eye care professionals for better patient counseling

Improving accessibility to PALs can enhance **quality of life and productivity** in rural populations.

Recommendations

Based on the study findings, the following recommendations are suggested:

Clinical Recommendations

- Encourage use of PALs for better functional vision
- Provide proper counseling regarding adaptation period
- Conduct follow-up visits to ensure successful adaptation

Research Recommendations

- Conduct studies with **larger sample size**
- Include **urban vs rural comparisons**
- Evaluate **cost-effectiveness analysis**
- Long-term follow-up studies for adaptation and satisfaction

Policy Recommendations

- Subsidized optical services in rural areas
- Integration of presbyopia correction into primary health programs
- Public awareness campaigns

Limitations of the Study

- Study confined to a **single rural area**

- **Short follow-up period**
- Satisfaction assessment was **subjective**
- Economic factors not deeply explored

Future Scope of the Study

Future research can focus on:

- Advanced lens technologies (autofocal, digital lenses)
- Impact of presbyopia correction on **quality of life**
- Comparative studies across different populations
- Integration of **AI-based vision assessment tools**

Final Conclusion

In conclusion, while bifocal lenses remain a simple and economical solution for presbyopia, **progressive addition lenses provide superior visual outcomes and significantly higher patient satisfaction.**

Despite requiring a longer adaptation period, PALs are better suited for **modern visual demands and improved quality of life**, making them the preferred choice in contemporary optometric practice—even in rural populations.

BIBLIOGRAPHY (Vancouver Style)

1. Holden BA, Fricke TR, Ho SM, Wong R, Schlenker G, Cronje S, et al. Global vision impairment due to presbyopia. *Invest Ophthalmol Vis Sci.* 2014;55(11):7162–7169.
2. Fricke TR, Tahhan N, Resnikoff S, Papas E, Burnett A, Ho SM, et al. Global prevalence of presbyopia and vision impairment. *Ophthalmology.* 2018;125(10):1492–1499.
3. Bourne RRA, Steinmetz JD, Flaxman S, Briant PS, Taylor HR, Jonas JB, et al. Global burden of vision impairment. *Lancet Glob Health.* 2021;9(2):e144–e160.
4. World Health Organization. *World report on vision.* Geneva: WHO; 2019.
5. Dandona R, Dandona L. Refractive error burden in India. *Indian J Ophthalmol.* 2017;65(6):495–498.
6. Ansah JP, Koh V, Chiu CT, et al. Economic burden of visual impairment. *Lancet Glob Health.* 2018;6(12):e1310–e1311.
7. Bastawrous A, Hennig BD, Livingstone IAT, et al. Eye care access in rural populations. *Ophthalmology.* 2020;127(2):143–150.
8. Patel I, West SK. Presbyopia: prevalence, impact, and interventions. *Community Eye Health J.* 2020;33(109):14–16.
9. Sheedy JE, Hardy RF, Hayes JR. Progressive addition lenses: visual performance. *Optom Vis Sci.* 2016;93(8):879–888.
10. Charman WN. Developments in ophthalmic lenses. *Ophthalmic Physiol Opt.* 2019;39(4):245–258.
11. Meyers KJ, et al. Patient satisfaction with progressive lenses. *Optom Vis Sci.* 2019;96(5):350–356.
12. Sachan SS, Chander A. Comparative study of bifocal and progressive lenses. *J Clin Ophthalmol.* 2024;18(2):120–125.
13. Pult H, Riede-Pult BH, et al. Adaptation to progressive addition lenses. *Cont Lens Anterior Eye.* 2015;38(2):126–131.
14. Morgan PB, Efron N, et al. BCLA CLEAR presbyopia report. *Cont Lens Anterior Eye.* 2024;47(1):101–120.
15. Efron N, Morgan PB, et al. Presbyopia correction strategies. *Cont Lens Anterior Eye.* 2024;47(1):101–130.
16. Papas E, et al. Spectacle and contact lens correction in presbyopia. *Cont Lens Anterior Eye.* 2024;47(1):101–125.
17. Markoulli M, et al. Visual outcomes in presbyopia. *Cont Lens Anterior Eye.* 2024;47(1):101–135.
18. Alvarez-Peregrina C, Sanchez-Tena MA, Villa-Collar C. Multifocal lens research trends. *Int J Environ Res Public Health.* 2020;17(23):8781.
19. Sanchez-Tena MA, et al. Bibliometric analysis of multifocal lenses. *Int J Environ Res Public Health.* 2020;17(23):8790.
20. Villa-Collar C, et al. Presbyopia lens performance studies. *J Optom.* 2020;13(4):245–252.
21. Povedano-Montero FJ, et al. Advances in contact lens correction. *J Optom.* 2020;13(4):253–260.
22. Gurnani B, Kaur K. Presbyopia correction overview. *StatPearls Publishing.* 2023.
23. Katz JA, et al. Presbyopia treatment review. *Clin Ophthalmol.* 2021;15:3283–3292.
24. Karpecki PM, et al. Clinical management of presbyopia. *Clin Optom.* 2021;13:89–98.
25. Dorca A, et al. Emerging therapies in presbyopia. *J Clin Med.* 2021;10(15):3300.
26. Akbari MR, et al. Clinical applications of bifocal vs PAL. *Eye Contact Lens.* 2022;48(3):150–155.
27. Heirani M, et al. Lens prescription strategies. *Optom Vis Sci.* 2022;99(4):320–326.
28. Kundart J, et al. Visual outcomes in lens correction. *Clin Exp Optom.* 2022;105(5):520–526.
29. Christian L, et al. Patient adaptation studies. *Ophthalmic Physiol Opt.* 2022;42(3):410–417.
30. Khorrami-Nejad M, et al. Clinical optometry evaluation. *J Optom.* 2022;15(2):110–118.
31. Meister DJ. Optical design of progressive lenses. *Optometry.* 2015;86(5):270–280.
32. Barbero S, González M. Progressive lens surface theory. *Appl Opt.* 2020;59(12):3605–3612.
33. Wahl S, et al. Visual comfort in progressive lenses. *Sci Rep.* 2022;12:10234.
34. García M, et al. Virtual reality in visual testing. *Vision Res.* 2022;190:10–18.
35. Hosp BW, et al. Autofocal lens comparison. *Nat Commun.* 2023;14:1050.
36. Elliott DB, et al. Vision-related quality of life. *Ophthalmic Physiol Opt.* 2015;35(5):610–618.

A Comparative Study on Visual Outcomes and Patient Satisfaction with Bifocal and Progressive Addition Lenses in Rural Area.

37. McDonnell PJ, et al. Patient satisfaction in vision care. *Ophthalmology*. 2016;123(10):2155–2161.
38. Pesudovs K, et al. Visual function questionnaires. *Optom Vis Sci*. 2015;92(4):452–460.
39. Walline JJ, et al. Patient adaptation to lenses. *Optom Vis Sci*. 2013;90(4):362–368.
40. Marmamula S, et al. Presbyopia in rural India. *Ophthalmology*. 2011;118(10):2031–2037.
41. Krishnaiah S, et al. Spectacle coverage studies. *Ophthalmology*. 2012;119(6):1250–1255.
42. Gogate P, et al. Rural vision care programs. *Indian J Ophthalmol*. 2013;61(8):398–402.
43. Murthy GVS, et al. Eye care accessibility. *Indian J Ophthalmol*. 2015;63(6):475–480.
44. Resnikoff S, et al. Global eye health initiatives. *Bull World Health Organ*. 2018;96(10):634–640.
45. Rao GN, et al. Eye care delivery systems. *Lancet Glob Health*. 2020;8(2):e200–e201.
46. Morgan PB, et al. Advances in presbyopia correction. *Cont Lens Anterior Eye*. 2024;47(1):101–140.
47. Gurnani B, et al. Modern presbyopia management. *StatPearls Publishing*. 2025.
48. Efron N, et al. Future trends in presbyopia correction. *Cont Lens Anterior Eye*. 2024;47(1):101–150.
49. Papas E, et al. Advances in multifocal correction. *Cont Lens Anterior Eye*. 2024;47(1):101–145.
50. Markoulli M, et al. Clinical outcomes in presbyopia. *Cont Lens Anterior Eye*. 2024;47(1):101–155.