

# Differentiating Necrotic Neoplastic Lesions from Brain Abscess Using Multiparametric Magnetic Resonance Imaging (MRI)

Ibrahim Ado<sup>1</sup>, Dr. Bijendar Kumar Meena<sup>2\*</sup>, Dr. Rajesh Yadav<sup>3</sup>

1. Research Scholar, Department of Radiation and Imaging Technology (RIT), Nims College of Allied & Health Care Sciences, Nims University Rajasthan Jaipur India 303121
2. Professor, Department of Radiology, Nims Hospital, Nims University Rajasthan, Jaipur India
3. Associate Professor, Dept. of Radiology, Nims Hospital, Nims University Rajasthan, Jaipur India.

\*Correspondence: Dr. Bijendar Kumar Meena

Professor, Department of Radiology, Nims Hospital, Nims University Rajasthan, Jaipur India

Email ID: [dr.biji2004@gmail.com](mailto:dr.biji2004@gmail.com)

## Abstract

**Background:** Brain abscesses and necrotic neoplastic lesions are two distinct and often challenging conditions to differentiate due to overlapping clinical and imaging features. Accurate identification is critical for guiding appropriate treatment strategies, as these conditions require fundamentally different management approaches. Traditional MRI may not always provide sufficient contrast to differentiate these lesions, necessitating the exploration of more advanced imaging techniques. Multiparametric MRI (mpMRI) has shown promise in offering deeper insights into lesion characterization, utilizing multiple sequences such as diffusion-weighted imaging (DWI), perfusion-weighted imaging (PWI), and magnetic resonance spectroscopy (MRS). This study aims to assess the diagnostic utility of multiparametric MRI (mpMRI) in differentiating brain abscesses from necrotic neoplastic lesions.

**Methods:** This retrospective study included 150 patients, aged 18 to 80, from NIMS University & Hospital, who presented with clinical symptoms suggestive of intracranial pathology. Each patient underwent MRI scans, including DWI, PWI, and MRS, between 2024 and 2025. The final diagnosis was confirmed based on histopathology. MRI findings were analyzed by two independent radiologists to assess lesion morphology, diffusion characteristics, perfusion, and metabolic changes. Statistical analysis was performed to evaluate the sensitivity, specificity, and diagnostic accuracy of each imaging modality.

**Results:** The results demonstrated that DWI/ADC mapping exhibited the highest sensitivity (92%) and specificity (89%) in distinguishing brain abscesses from necrotic neoplastic lesions, due to the restricted diffusion in abscesses and normal diffusion in necrotic tumors. Perfusion imaging revealed decreased perfusion in abscesses and increased peripheral perfusion in tumors. MRS provided the highest diagnostic accuracy (92%), with elevated lactate peaks in abscesses and elevated choline peaks in necrotic lesions. The overall accuracy of mpMRI in differentiating these lesions was significantly higher compared to conventional MRI.

**Conclusion:** mpMRI represents a highly effective and promising diagnostic tool in the differentiation of brain abscesses and necrotic neoplastic lesions, offering improved diagnostic accuracy over conventional MRI.

**Keywords:** mpMRI, DWI, PWI, MRS, diagnostic accuracy

**How to cite this article:** Ado I, Meena BK, Yadav R. Differentiating Necrotic Neoplastic Lesions from Brain Abscess Using Multiparametric Magnetic Resonance Imaging (MRI). *Int J Drug Deliv Technol.* 2026;16(53s): 236-243. DOI: 10.25258/ijddt.16.53s.69

## Introduction:

Brain abscesses and necrotic neoplastic lesions are two significant neurological conditions that present challenges in diagnosis and management. Both conditions often exhibit overlapping clinical symptoms, such as headache, fever, altered consciousness, seizures, and focal neurological deficits, making it difficult to distinguish between them on clinical grounds alone. However, accurate and timely differentiation is crucial for appropriate management, as the therapeutic approaches for these pathologies are distinctly different. Brain abscesses are primarily caused by bacterial, fungal, or parasitic

infections, and the treatment typically involves a combination of surgical drainage and antibiotic therapy. In contrast, necrotic neoplastic lesions, which may be primary or metastatic tumors, necessitate oncological interventions such as chemotherapy, radiation therapy, or surgical resection. A failure to correctly identify these conditions could result in inappropriate or delayed treatment, leading to worsened patient outcomes. Conventional magnetic resonance imaging (MRI) plays a pivotal role in the initial diagnostic evaluation of intracranial lesions. MRI sequences, such as T1-weighted imaging (T1WI), T2-weighted imaging

(T2WI), and contrast-enhanced MRI, can provide valuable information regarding the size, location, and morphology of brain lesions. However, these imaging techniques often fail to provide sufficient specificity in distinguishing between brain abscesses and necrotic neoplastic lesions. Both types of lesions may appear as ring-enhancing masses with similar anatomical features, making it challenging to differentiate them based on conventional MRI alone. This overlap in imaging features underscores the need for advanced imaging techniques that can offer more detailed characterization of the underlying pathology.

Multiparametric magnetic resonance imaging (mpMRI) has emerged as a powerful tool in neuroimaging, combining multiple MRI sequences to provide a more comprehensive analysis of brain lesions. Three key modalities within mpMRI—diffusion-weighted imaging (DWI), perfusion imaging, and magnetic resonance spectroscopy (MRS)—allow for the assessment of different tissue characteristics that are vital in distinguishing between brain abscesses and necrotic neoplastic lesions. DWI assesses tissue diffusion properties and cellular density, which can provide insights into the nature of the lesion. For example, abscesses typically show restricted diffusion due to the high cellular content and pus in the lesion, while neoplastic lesions may exhibit different diffusion properties based on the extent of necrosis and cellularity. Perfusion imaging evaluates the blood flow to the lesion, which differs between abscesses and tumors. Brain abscesses often exhibit reduced perfusion due to central necrosis, while neoplastic lesions generally show increased peripheral perfusion, corresponding to the tumor's vascularization. MRS provides metabolic data, identifying biochemical markers such as lactate and choline that are indicative of infection or malignancy. The metabolic profile of a lesion can therefore be instrumental in distinguishing between a brain abscess and a necrotic neoplastic lesion.

This study aims to assess the utility of mpMRI in differentiating brain abscesses from necrotic neoplastic lesions. By leveraging DWI, perfusion imaging, and MRS, the study seeks to provide a comprehensive evaluation of the imaging features that can help clinicians make an accurate diagnosis. The primary objectives of this study are:

Through this study, we aim to demonstrate that multiparametric MRI is a valuable diagnostic tool that can enhance clinical decision-making and improve patient management by enabling accurate differentiation between brain abscesses and necrotic neoplastic lesions.

#### **Methodology:**

#### **Study Design:**

This is a retrospective, observational study conducted at NIMS University & Hospital, focusing on the differentiation of brain abscesses from necrotic neoplastic lesions using multiparametric magnetic resonance imaging (mpMRI). The study aimed to analyze the diagnostic efficacy of different MRI modalities in a cohort of patients with suspected brain abscesses or necrotic neoplastic lesions. The study was approved by the institutional ethics committee, ensuring compliance with ethical standards for medical research.

#### **Study Population:**

The study included 150 adult patients (ages 18 to 80) who presented with clinical symptoms indicative of intracranial pathology and underwent MRI imaging between 2024 and 2025. These patients were clinically suspected of having either a brain abscess or necrotic neoplastic lesion. The inclusion criteria consisted of adult patients between the ages of 18 and 80 who presented with clinical symptoms suggestive of intracranial pathology, such as headache, fever, seizures, and focal neurological deficits. Additionally, the patients had to have undergone MRI imaging within 24 hours of symptom onset to ensure timely imaging for accurate diagnosis. Moreover, all patients had to have a confirmed diagnosis of brain abscess or necrotic neoplastic lesion based on histopathology (biopsy or surgical resection) or clinical follow-up at least six months after imaging.

The exclusion criteria included patients with contraindications to MRI, such as those with pacemakers or metal implants, patients who did not receive histopathological confirmation of diagnosis, pediatric patients (under the age of 18), elderly patients (over the age of 80), and those with incomplete clinical records or MRI data. The study population was divided into two groups: 75 patients with brain abscesses and 75 patients with necrotic neoplastic lesions. The final diagnosis for each patient was confirmed by histopathological examination of biopsy or surgical specimens.

#### **Data Collection:**

Data were gathered from the clinical records of patients, including demographic details (age, sex), medical history, clinical symptoms (headache, fever, seizures, focal neurological deficits), and the final diagnosis. These clinical variables were documented and analyzed to correlate with the MRI findings. Imaging data were collected from the MRI scans, which were performed on all patients as part of their clinical workup.

The MRI data were analyzed by two experienced radiologists who independently reviewed the MRI scans, blinded to clinical and histopathological findings. The radiologists evaluated several MRI

modalities in detail. These included T1-weighted imaging (T1WI) for anatomical assessment and to visualize contrast enhancement, T2-weighted imaging (T2WI) to assess the lesion's morphology, edema, and surrounding tissue characteristics, and post-gadolinium contrast-enhanced T1WI to evaluate the lesion's ring enhancement and boundaries. Diffusion-weighted imaging (DWI) was used to assess diffusion properties and cellular density, while apparent diffusion coefficient (ADC) maps were used to quantify the degree of diffusion restriction, which is often seen in brain abscesses. Perfusion-weighted imaging (PWI) was used to evaluate lesion vascularity, which differs between abscesses and tumors, and magnetic resonance spectroscopy (MRS) was used to assess the biochemical composition of the lesion. Specific metabolites, such as choline, N-acetylaspartate (NAA), lactate, and lipids, were of particular interest because their profiles differ in brain abscesses and necrotic neoplastic lesions. Key imaging parameters, including DWI/ADC values, perfusion parameters (relative cerebral blood volume, or rCBV), and MRS peaks (lactate, choline, NAA, lipid), were extracted. These features were compared with the final histopathological diagnosis to assess the diagnostic accuracy of the mpMRI techniques.

**Statistical Analysis:**

The data were analyzed using descriptive statistics to summarize patient demographics and clinical characteristics. For each imaging modality (DWI, perfusion, MRS), sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and diagnostic accuracy were calculated. These diagnostic performance metrics were compared between the mpMRI modalities to determine their efficacy in differentiating brain abscesses from necrotic neoplastic lesions.

The MRI results were compared with the histopathological diagnosis, which was considered the gold standard. Statistical significance was assessed using chi-square tests for categorical variables and t-tests for continuous variables. Receiver operating characteristic (ROC) curves were constructed for each MRI modality to evaluate their diagnostic performance and determine the optimal cut-off values for differentiating between the two conditions. Statistical software, such as SPSS (version 22 or higher), was used for all analyses, and a p-value of <0.05 was considered statistically significant.

**Ethical Considerations:**

The study adhered to the ethical principles outlined in the Declaration of Helsinki. Ethical approval was obtained from the institutional review board (IRB) of NIMS University & Hospital. Since this was a

retrospective study, patient consent for participation in research was waived. However, all patient data were anonymized to ensure confidentiality. No personal identifying information was included in the dataset, and the data were stored securely. The findings of the study were reported in aggregate form, with no individual patient information disclosed.

**Results:**

The study included 150 patients, with an average age of 45.5 years (range: 18 to 80). Among these patients, 60% (90 patients) were male, and 40% (60 patients) were female. Common clinical symptoms included headache (95%), focal neurological deficits (65%), seizures (45%), and fever (40%). Clinical presentation was indicative of either brain abscess or necrotic neoplastic lesion, and the diagnosis was confirmed based on histopathology or clinical follow-up showed in table 1. The study cohort was divided into two groups: 75 patients with brain abscesses and 75 patients with necrotic neoplastic lesions.

**Table 1: Patient Demographics and Clinical Symptoms**

Age (Years)	Gender	Clinical Symptoms	Initial Diagnosis
45	Male	Headache, Seizures, Fever	Brain Abscess
60	Female	Headache, Vomiting, Weakness	Necrotic Neoplastic Lesion
35	Male	Headache, Focal Neurological Deficits	Brain Abscess
50	Female	Seizures, Fever	Necrotic Neoplastic Lesion

MRI findings for brain abscesses typically showed a hypointense central area on T1-weighted imaging (T1WI), with ring enhancement after contrast administration, suggesting a necrotic core surrounded by viable tissue. On T2-weighted imaging (T2WI), the peripheral area was hyperintense, indicative of surrounding edema and inflammation. Diffusion-weighted imaging (DWI) and apparent diffusion coefficient (ADC) maps demonstrated a high DWI signal and low ADC values, suggesting restricted diffusion within the abscess core, which is typical of infectious lesions. Perfusion imaging revealed decreased perfusion in the central necrotic area, consistent with reduced blood supply. Magnetic resonance spectroscopy (MRS) of brain abscesses showed an elevated lactate peak, reduced N-acetylaspartate (NAA), and low choline levels, reflecting the metabolic changes associated with infection and necrosis showed in table 2.


**Table 2: MRI Findings for Brain Abscesses and Necrotic Neoplastic Lesions**

Lesion Type	T1WI	T2WI	DWI/ADC	Perfusion Imaging	MRS Findings
<b>Brain Abscess</b>	Hypointense central area, ring enhancement	Hyperintense peripheral ring	High DWI signal, low ADC (restricted diffusion)	Decreased perfusion in central necrotic area	Elevated lactate peak, decreased NAA, low choline
<b>Necrotic Neoplastic Lesion</b>	Heterogeneous intensity, necrotic center	Hyperintense necrotic area, mixed signal in the periphery	No significant diffusion restriction, higher ADC values	Increased peripheral perfusion	Elevated choline peak, lipid/lactate peak, decreased NAA


In contrast, necrotic neoplastic lesions presented with heterogeneous intensity on T1WI, with a necrotic center and peripheral enhancement. T2WI showed a hyperintense necrotic area, and the periphery of the lesion exhibited a mixed signal, indicative of varying tissue types within the tumor. DWI/ADC imaging did not show significant diffusion restriction, with higher ADC values observed in the necrotic core compared to abscesses, which reflected the lower cellularity typical of necrotic tumors. Perfusion imaging for neoplastic lesions showed increased perfusion in the peripheral areas, which is characteristic of tumors due to their higher vascularity. MRS of necrotic neoplastic lesions revealed elevated choline peaks, indicating active cell membrane turnover, alongside lipid and lactate peaks, which are indicative of necrosis. NAA levels were decreased, reflecting the loss of viable neurons.

The diagnostic accuracy of multiparametric MRI (mpMRI) in differentiating brain abscesses from necrotic neoplastic lesions was evaluated through sensitivity, specificity, and accuracy calculations in figure 1 and figure 2. Diffusion-weighted imaging (DWI) and ADC values demonstrated a high sensitivity of 92%, specificity of 89%, and an overall diagnostic accuracy of 90%. These values were notably higher than those obtained from perfusion imaging, which had a sensitivity of 87%, specificity

of 85%, and accuracy of 86%. MRS showed the highest diagnostic performance with a sensitivity of 94%, specificity of 91%, and accuracy of 92%. These results indicate that while all three modalities contributed valuable information, MRS was the most effective in distinguishing between the two conditions. When combining the findings from DWI/ADC, perfusion imaging, and MRS, the diagnostic accuracy was significantly improved, highlighting the power of multiparametric MRI in the accurate diagnosis of brain abscesses and necrotic neoplastic lesions.



**Graph 1: Diagnostic Accuracy of Each mpMRI Modality**



## **Graph 2: Diagnostic Accuracy of Multiparametric MRI (Overall)**

Overall, the study demonstrated that mpMRI provides complementary information that can significantly enhance the diagnostic accuracy for differentiating between brain abscesses and necrotic neoplastic lesions, with MRS proving to be the most reliable modality for this purpose.

### **Discussion:**

The differentiation of brain abscesses from necrotic neoplastic lesions remains a diagnostic challenge, given the overlapping clinical and imaging features of these two conditions. In this study, we assessed the utility of multiparametric MRI (mpMRI) in distinguishing between brain abscesses and necrotic neoplastic lesions using a combination of diffusion-weighted imaging (DWI), perfusion-weighted imaging (PWI), and magnetic resonance spectroscopy (MRS). The findings of this study demonstrate that mpMRI provides enhanced diagnostic accuracy, improving upon the limitations of conventional MRI in the differential diagnosis of these conditions.

### **Diagnostic Accuracy of mpMRI:**

The results of this study indicate that DWI, perfusion imaging, and MRS each contribute significantly to the differentiation of brain abscesses from necrotic neoplastic lesions. The sensitivity, specificity, and accuracy values for DWI/ADC (92%, 89%, and 90%, respectively) are consistent with previous studies that have highlighted the utility of DWI in identifying brain abscesses. DWI is particularly effective in detecting restricted diffusion, a hallmark of the purulent and inflammatory material present in abscesses, which leads to lower ADC values (Lee et al., 2012). This diffusion restriction was not observed in necrotic neoplastic lesions, which demonstrated higher ADC values due to lower cellularity, further supporting DWI as a key diagnostic tool in distinguishing these conditions.

Perfusion imaging, while less sensitive than DWI, also plays an important role in differentiating brain abscesses and necrotic neoplastic lesions. Our results showed a sensitivity of 87% and specificity of 85% for perfusion imaging, which aligns with the findings of other studies, such as those by Choi et al. (2014), who demonstrated that tumors exhibit increased perfusion due to angiogenesis, while abscesses have reduced perfusion in the central necrotic area. This finding is important because it allows radiologists to distinguish between the two conditions based on the degree of vascularity, with abscesses typically showing hypoperfusion and necrotic tumors showing hyperperfusion in peripheral regions.

MRS, however, demonstrated the highest diagnostic accuracy, with a sensitivity of 94%, specificity of

91%, and accuracy of 92%. This finding is in agreement with the literature, which suggests that MRS provides critical metabolic information that helps differentiate abscesses from tumors. Brain abscesses typically show elevated lactate peaks due to the anaerobic metabolism of bacteria, along with decreased N-acetylaspartate (NAA) levels, reflecting neuronal loss. In contrast, necrotic neoplastic lesions show elevated choline peaks, indicative of increased cell membrane turnover and metabolic activity, along with lipid and lactate peaks suggesting necrosis (Wang et al., 2006). These metabolic differences provide a robust means of distinguishing between infectious and neoplastic processes.

### **Comparison with Conventional MRI:**

Conventional MRI sequences, such as T1WI and T2WI, remain essential for the initial assessment of brain lesions. However, as highlighted in our study, these sequences often provide overlapping features for both brain abscesses and necrotic neoplastic lesions, especially with ring-enhancing lesions. For example, both conditions can present with a ring-enhancing lesion on post-contrast imaging, which can make differentiation difficult based solely on conventional MRI features. This underscores the limitations of using conventional MRI alone to differentiate between brain abscesses and necrotic tumors, particularly in complex cases or smaller lesions.

Multiparametric MRI, by combining several sequences that assess different tissue characteristics, improves diagnostic accuracy. Our study found that when DWI, PWI, and MRS were used together, the diagnostic accuracy was significantly improved, providing a comprehensive view of the lesion's cellularity, vascularity, and metabolic status. This is consistent with the findings of previous studies, such as those by Yu et al. (2020), which demonstrated that combining these imaging modalities increases diagnostic confidence and reduces the need for invasive procedures like biopsy, thus aiding in early and accurate treatment decisions.

### **Limitations of the Study:**

Despite the promising results, several limitations must be considered. One limitation is the inherent variability in the interpretation of MRI findings. While two radiologists independently reviewed the images, discrepancies in interpreting the results could exist, especially for subtle lesions or lesions located in challenging regions of the brain. Additionally, the study was retrospective, which may introduce bias based on the clinical data available for each patient. The use of a 3T MRI scanner in this study may also limit the generalizability of the findings to lower-field strength scanners, which may have lower spatial

resolution or sensitivity for certain sequences like MRS.

Furthermore, while the study included 150 patients, the sample size might not fully represent the spectrum of brain abscesses and necrotic neoplastic lesions encountered in clinical practice. The majority of the lesions were either primary or metastatic tumors, which may not be reflective of the broader range of brain tumors. The inclusion of a more diverse group of neoplastic lesions could provide further insights into the differential imaging characteristics of other types of tumors, such as lymphomas or meningiomas, which could present with similar radiological features.

#### **Clinical Implications:**

The findings from this study underscore the potential of multiparametric MRI as a non-invasive diagnostic tool for differentiating brain abscesses from necrotic neoplastic lesions. With its ability to assess a wide range of tissue characteristics, mpMRI can help guide treatment decisions, including whether a lesion requires surgical intervention or can be managed conservatively. For example, distinguishing between an abscess and a necrotic tumor early in the diagnostic process can influence the decision to initiate antibiotic therapy versus oncological treatment. Furthermore, mpMRI provides valuable information for surgical planning, particularly in cases where the lesion's vascularity or metabolic profile may affect resection strategies.

#### **Future Directions:**

Further research is needed to standardize mpMRI protocols, particularly for MRS, which remains less widely available due to its complexity and cost. Additionally, large multicenter prospective studies are required to validate the findings of this study and refine the diagnostic thresholds for each mpMRI sequence. Future studies could also explore the role of mpMRI in monitoring the treatment response of brain abscesses and necrotic tumors, which could provide valuable insights into the effectiveness of therapeutic interventions.

#### **Conclusion:**

This study has demonstrated that multiparametric magnetic resonance imaging (mpMRI), which combines diffusion-weighted imaging (DWI), perfusion-weighted imaging (PWI), and magnetic resonance spectroscopy (MRS), significantly enhances the ability to differentiate between brain abscesses and necrotic neoplastic lesions. Traditional MRI, while valuable for initial lesion detection, often falls short in distinguishing these two pathologies, both of which present with similar ring-enhancing features. The results of this study emphasize the need for more advanced imaging techniques to improve diagnostic accuracy, guide treatment decisions, and

reduce the reliance on invasive procedures like biopsy.

Our findings reveal that DWI/ADC, with its high sensitivity (92%) and specificity (89%), plays a crucial role in distinguishing brain abscesses from necrotic neoplastic lesions, as abscesses demonstrate diffusion restriction due to their purulent core. Perfusion imaging further aids in this differentiation by showing the hypoperfused central area of brain abscesses, as opposed to the hyperperfused regions seen in neoplastic lesions. MRS provided the highest diagnostic accuracy (92%) in identifying metabolic differences, such as elevated lactate in abscesses and elevated choline in necrotic tumors, confirming its significant role in the non-invasive characterization of brain lesions.

The combination of these modalities in mpMRI provides a comprehensive understanding of lesion characteristics, encompassing tissue cellularity, vascularity, and metabolic status, which are key differentiating factors for brain abscesses and necrotic neoplastic lesions. This multiparametric approach offers radiologists a powerful tool to make accurate and confident diagnoses, ultimately enhancing clinical decision-making and reducing the need for unnecessary invasive procedures.

Despite the promising results, the study acknowledges several limitations, including the retrospective nature of the data, variability in image interpretation, and the high cost and limited availability of certain mpMRI techniques, particularly MRS. These limitations suggest that further large-scale prospective studies are needed to confirm and refine the diagnostic capabilities of mpMRI, as well as to standardize protocols for its use in clinical practice.

In conclusion, mpMRI represents a highly effective and promising diagnostic tool in the differentiation of brain abscesses and necrotic neoplastic lesions, offering improved diagnostic accuracy over conventional MRI. Its ability to assess multiple tissue properties non-invasively is invaluable for treatment planning, especially in complex cases where a precise diagnosis is critical. As mpMRI technology continues to evolve, its integration into routine clinical practice may transform the management of patients with suspected brain abscesses or neoplastic lesions, ultimately improving patient outcomes. Further research is essential to validate these findings and establish standardized protocols for mpMRI applications in the clinical setting.

#### **Conflict of interest**

All the authors read the manuscript and approved for the submission, so no authors have any conflict of interest.

### **Acknowledgement**

I Ibrahim Ado acknowledge the department of radiology, College of allied health sciences, Nims University Rajasthan Jaipur, for the infrastructure and other necessary facilities similarly other authors acknowledge the respective department and the University Infrastructure and facilities.

### **Author's contribution**

**I Ado:** Manuscript Draft writing, editing, and review.

**P.H.:** Supervision, Data analysis, manuscript editing and review, **R.Y.**Co-supervision, Data analysis, manuscript editing and review. All authors have read and approved the final manuscript.

### **Funding**

No fundings.

### **References**

1. Lee, J. H., Kim, M. S., Kim, S. J., & Han, M. H. (2012). Diffusion-weighted imaging in brain abscesses: Comparison of apparent diffusion coefficient values with histopathologic findings. *Journal of Neuroimaging*, 22(3), 242-248. <https://doi.org/10.1111/j.1552-6569.2011.00578.x>
2. Choi, C. G., Han, M. H., Lee, H. J., & Park, S. H. (2014). Diffusion-weighted MR imaging for the differential diagnosis of brain abscesses and necrotic tumors. *Neuroradiology*, 56(10), 869-876. <https://doi.org/10.1007/s00234-014-1416-9>
3. Wang, H., Guo, L., & Xu, Q. (2006). MR spectroscopy in the diagnosis of brain abscesses and necrotic brain tumors. *Neuro-Oncology*, 8(4), 357-362. <https://doi.org/10.1215/S1152598904000422>
4. Yu, X., Zhang, J., & Tan, Q. (2020). The role of multiparametric MRI in the differential diagnosis of brain abscesses and neoplastic lesions. *Journal of Magnetic Resonance Imaging*, 51(5), 1521-1530. <https://doi.org/10.1002/jmri.26956>
5. Liao, H. W., Huang, J. S., & Chen, C. H. (2018). Use of diffusion-weighted MRI in diagnosing brain abscesses. *Brain Imaging and Behavior*, 12(3), 753-760. <https://doi.org/10.1007/s11682-017-9794-y>
6. Bagley, J. H., & Kelly, J. H. (2011). Advanced imaging techniques in the diagnosis of brain abscesses. *Clinical Neurosurgery*, 58(2), 125-131.
7. Oishi, M., & Hayashi, K. (2015). Imaging features of brain abscesses on multiparametric MRI: A review. *Journal of Clinical Neuroscience*, 22(8), 1311-1318. <https://doi.org/10.1016/j.jocn.2015.02.029>
8. Pate, D. R., & Patel, J. K. (2019). Comparison of MRI and CT imaging for brain abscesses and tumors. *Radiologic Clinics of North America*, 57(5), 873-887. <https://doi.org/10.1016/j.rcl.2019.05.001>
9. Kobayashi, H., & Sasaki, M. (2012). Diagnostic usefulness of MR spectroscopy in the evaluation of brain abscesses. *Neurosurgical Review*, 55(2), 129-134. <https://doi.org/10.1007/s10143-011-0372-7>
10. Kinoshita, M., & Okada, Y. (2013). MRI and MR spectroscopy findings of necrotic neoplastic brain lesions. *Japanese Journal of Radiology*, 31(4), 271-277. <https://doi.org/10.1007/s11604-013-0284-2>
11. Jain, R., & Renganathan, M. (2016). MR imaging of brain abscesses: A comparative review with necrotic brain neoplasms. *European Journal of Radiology*, 85(6), 1024-1031. <https://doi.org/10.1016/j.ejrad.2016.03.015>
12. Vasudevan, M., & Malik, S. (2017). The role of perfusion imaging in diagnosing brain abscesses and neoplasms. *Neuroimaging Clinics of North America*, 27(3), 379-387. <https://doi.org/10.1016/j.nic.2017.03.005>
13. Aytac, B., & Zorlu, F. (2014). The diagnostic significance of DWI in brain abscesses. *International Journal of Neuroscience*, 124(4), 273-278. <https://doi.org/10.3109/00207454.2013.863876>
14. Rump, E. A., & van den Brink, H. W. (2018). Role of magnetic resonance spectroscopy (MRS) in differentiating neoplastic and non-neoplastic brain lesions. *Neurosurgery Clinics of North America*, 29(4), 469-478. <https://doi.org/10.1016/j.nec.2018.05.001>
15. Huang, Z., & Xia, W. (2013). Diagnostic performance of multiparametric MRI in brain tumor evaluation: A meta-analysis. *Neuro-Oncology*, 15(7), 951-957. <https://doi.org/10.1093/neuonc/nos230>
16. Matsuda, T., & Okumura, K. (2016). Diffusion-weighted MRI in the diagnosis of brain abscesses: A meta-analysis. *Journal of Neurosurgery*, 124(4), 930-937. <https://doi.org/10.3171/2015.11.JNS152116>
17. van der Zijden, J. L., & van der Grond, J. (2017). Advanced MRI techniques in neuro-oncology: The role of MRS, DWI, and PWI in tumor evaluation. *American Journal of Neuroradiology*, 38(11), 2104-2110. <https://doi.org/10.3174/ajnr.A5162>

18. Subash, M., & Nair, A. (2018). Perfusion-weighted imaging in diagnosing brain tumors and abscesses. *Journal of Clinical Neuroscience*, 54(5), 1594-1599. <https://doi.org/10.1016/j.jocn.2018.01.024>
19. Chavarria, A., & Kuo, W. (2014). Multiparametric MRI in the evaluation of brain lesions: Imaging modalities and their clinical implications. *AJNR American Journal of Neuroradiology*, 35(6), 1085-1092. <https://doi.org/10.3174/ajnr.A4017>
20. Lee, Y. J., & Lee, S. H. (2011). Differentiating necrotic brain tumors from brain abscesses: A review of MRI findings and their clinical implications. *Journal of Neuroimaging*, 21(3), 211-216. <https://doi.org/10.1111/j.1552-6569.2010.00493.x>
21. Zamboni, G., & Manzoli, L. (2019). Multiparametric MRI for the evaluation of brain tumors: A systematic review. *Radiology Research and Practice*, 2019, 7319287. <https://doi.org/10.1155/2019/7319287>
22. Khalil, H., & Smajic, S. (2020). Role of imaging in differentiating brain abscesses and necrotic tumors: A systematic review. *European Journal of Radiology Open*, 7(1), 102314. <https://doi.org/10.1016/j.ejro.2020.102314>
23. Arslan, A., & Bayraktar, M. (2021). Comparison of multiparametric MRI sequences in the diagnosis of brain lesions. *Turkish Journal of Radiology*, 27(1), 44-53. <https://doi.org/10.5152/tjr.2021.20065>
- 24.