

Machine Learning for Quantum Dots Analysis: Bridging Data-Driven Methods with Quantum Technology

P. DHYVA STUTHI^{1,*}, P. PAVANI KOTESWARI DEVI², K K V SATYANARAYANA³,
CHANDOLU NAGAMANI⁴, SYED FAROOQ⁵, YETI DANA RAO⁵,
P K VENKATA DURGA³, B. KIRAN BABU⁶

^{1,*}Department of Basic Sciences and Humanities,
Usha Rama College of Engineering & Technology, A.P, India.

²Department of Basic Sciences and Humanities,
CMR Technical Campus, Hyderabad, Telangana, India.

³Dept. of Computer Science and Engineering, Aditya University, Surampalem, A. P, India

⁴Department of Computer Science and Engineering,
Koneru Lakshmaiah Education Foundation, Vaddeswaram, A. P, India

⁵Department of Physics, Rajeev Gandhi Memorial College of Engineering and Technology, Nandyal, A. P, India

⁶Department of Botany and Microbiology, Acharya Nagarjuna University, Guntur, A. P, India

*Corresponding Author: *P. DHYVA STUTHI*

Abstract

Quantum dots (QDs) are nanomaterials at the nanoscale that have completely new optical and electronic properties resulting from quantum confinement effects. Present, the traditional approach to quantum dot research involved a cyclical process of experiments, computational simulations, and manual parameter adjustments. Still, with the growing complexity of systems and the demand for higher precision in applications, this approach is rapidly becoming insufficient. Machine learning (ML) is a new technology that is fundamentally changing the way scientists synthesize, characterize and optimize QD devices. This in-depth paper investigates the use of machine learning in quantum dot research experimentally, practically, and computationally. We look at how data-driven methods are bridging and improving traditional physical workflows, examine the variety of algorithms used in QD research, share in-depth studies and most importantly impact the field. In addition, the opportunities and challenges that this interdisciplinary field faces was discussed.

Keywords: Quantum dots, Machine learning, Data-driven optimization, Synthesis.

How to cite this article: Dhyva Stuthi P, Pavani Koteswari Devi P, Satyanarayana KKV, Nagamani C, Farooq S, Dana Rao Y, Venkata Durga PK, Kiran Babu B. Machine Learning for Quantum Dots Analysis: Bridging Data-Driven Methods with Quantum Technology. Int J Drug Deliv Technol. 2026;16(54s): 708-716. DOI: 10.25258/ijddt.16.54s.62

1. Introduction

Over the last few decades, the uncovering and advancement of quantum dots have resulted in a highly productive synergy between the basic quantum physics and applied materials engineering. Semiconductor quantum dots (QDs) or nanocrystals usually of 2 to 10 nm in size have tunable optical properties and size that are due to quantum confinement[1]. If the carriers of charges are confined in the three spatial dimensions to a scale close to the exciton Bohr radius, the energy bands of the bulk semiconductors change to discrete energy levels like atoms. The properties of the material due to this confinement are highly tunable by just controlling the particle size shape composition and surface chemistry, which makes quantum dots extremely versatile for applications including bioimaging photovoltaics LEDs, quantum computing and sensing[2].

These remarkable characteristics, excluding the actual implementation of high-performance QD-based devices is still limited due to the almost unparalleled complexity of the parameter space. QD synthesis is a multivariate process that entails dozens of variables, the amounts of the precursors, temperatures of reactions, rates of injection, identities of ligands, composition of solvents, and the conditions of post-processing which together, nonlinearly and in a coupled manner, affect the final article[3]. Conventional optimization methods, which rely on changing one variable at a time in the experiments or conducting expensive ab initio simulations, not only have great difficulty in effectively handling the complexity but also, as a consequence, have given rise to a research pattern marked by very slow advancement, massive waste of materials, and overreliance on the expert intuition which has been built up through years of laboratory work[4,5].

Machine learning is really changing the way things are done here on a deep level. By getting patterns from big sets of experimental data, ML methods can find unsuspected forms in the high-dimensional parameters, create predictive models that work beyond the training data and steer experimental planning to reach best results with only a small fraction of the experimental work[6].

2. Quantum Dot Physics: Opportunities and Challenges

2.1 Quantum Confinement and Size-Dependent Properties

The main feature that sets quantum dots apart is the quantum confinement effect or the limitation of charge carrier movement in all three spatial dimensions to the sizes smaller than the bulk exciton Bohr radius. In bulk semiconductors, electrons and holes are so widely spread throughout the crystal lattice the bands of their energies are almost continuous. When a nanocrystal size is close to the one of an exciton Bohr radius, the increase in energy due to confinement and the energy arising from interaction between a pair of charge carriers have similar values and the energy spectrum changes from continuous to discrete[7].

This type of discretization has major implications practically. Most importantly, the optical bandgap of a quantum dot goes up as its size goes down and this property, coupled with the ability to control particle sizes, enables one to choose very accurately the spectral properties. The CdSe QDs emission wavelength 'for instance-the probably most widely studied QD system can be smoothly varied between about 450 nm (blue) for 2 nm particles and 650 nm (red) for 7 nm particles. The emission linewidth of a single QD can be very narrow (< 15 meV even at cryogenic temperatures) and ensemble linewidths have been brought down to less than 25 meV by improvements in synthesis monodispersity[8].

Besides optical properties, quantum confinement also changes carrier transport, Auger recombination rates, phonon coupling and spin dynamics in such ways that they create opportunities as well as challenges for device applications. The reduction or elimination of the nonradiative Auger recombination - the major loss mechanism in bulk semiconductors at high carrier densities - can be achieved by tailoring QD size and shape, Because of this providing the possibility of high-efficiency single-photon emission and low threshold stimulated emission[9].

2.2 The Parameter Space Challenge

Quantum dots' tunability is the very feature that also poses the greatest difficulty in controlling the vast array of parameters that define their behavior.

Colloidal synthesis - the chief method for making premium-quality QDs in large quantities - involves a nucleation and growth process directed through the LaMer mechanism. A nucleation burst needs to be time-wise separated from the growth phase to result in narrow size distributions and this separation is highly dependent on the interplay between monomer consumption rates, diffusion kinetics and surface energy aspects[10,11]. In reality, it is typically that a single synthesis method comprises over twenty individual factors like several precursor concentrations, injection temperatures, growth temperatures, reaction times, types and concentrations of ligands, compositions of solvent, and conditions of post-synthetic treatments.

The relationships between these factors are very nonlinear and change with the context; altering one parameter usually makes it necessary to change a few others to keep the quality of the product. This combination explosion renders brute-force experimental optimization almost impossible[12].

The characterization struggle is similar in difficulty. A thorough QD characterization involves the combination of various methods of X-ray diffraction for crystal structure, transmission electron microscopy for size and shape, UV-visible absorption spectroscopy for bandgap, photoluminescence spectroscopy for emission properties, time-resolved fluorescence for dynamics and inductively coupled plasma mass spectrometry for composition - each producing large, diverse datasets that need to be analyzed and interpreted together[13,14].

2.3 Application-Driven Demands

The expanding range of QD applications demands higher and higher performance levels, which in turn increase the difficulties in synthesis and characterization. For the display applications of quantum dot LEDs (QLEDs), external quantum efficiencies well above 20% are required, emission linewidths should be lower than 20 nm, and operation lifetimes over 10,000 hours with reproducible color coordinates that meet ITU-R BT.2020 color space standards are also necessary[15].

For quantum dot solar cells, the morphology of the QD film, surface passivation, and charge transport layers need to be designed carefully so that light absorption, carrier collection and stability are all maximized at the same time. Biomedical QDs, Then again, should not only be brightly and stably fluorescent but also capable of being biocompatible, delivered in a targeted manner, and have their pharmacokinetics controlled[16–18].

3. Machine Learning Methodologies in QD Research

3.1 Supervised Learning

Supervised learning is the major form of machine learning in quantum dot (QD) research. The approach involves training a machine learning model with a labeled data set -- pairs of input (synthesis parameters, structural descriptors, or spectra) and output for the model then making output predictions based on new inputs.

The training is aimed at minimizing a loss function quantifying the divergence of model predictions from the actual labels. Random forests, an ensemble method which consists in pooling the predictions of many decision trees that have been trained independently, have become a prime algorithm for predicting QD properties[19,20]. Their blend of high prediction accuracy, resistance to overfitting, and ability to estimate feature importance internally renders them Inparticular suitable for QD datasets which being moderate in size, high in dimensionality, and noisy, are quite typical in these respects.

Gradient boosted trees, including popular implementations like XGBoost and LightGBM, provide similar benefits, with the added advantage of often delivering better performance on **Table 1**[21].

Table 1 The comparative aspect of algorithm and its best use in QD research

Algorithm	Best use in QD research
Random forest	Predicting emission wavelength and from synthesis parameters; importance ranking
Gradient boosted trees XGBoost, LightGBM	Property prediction from synthesis/composition data; QD bandgap correlations
Convolutional neural networks CNNs	Spectral analysis (PL, UV-Vis); image segmentation and particle phase identification via Raman
Recurrent networks RNNs / LSTMs / GRUs	Time-resolved PL decay analysis; dynamic synthesis monitoring; temporal property tracking
Generative adversarial networks GANs	Ligand-QD surface interaction modeling; predicting effect of surface chemistry on optical properties

Deep learning, Mostly through convolutional neural networks (CNNs), has opened up new possibilities for handling QD spectral and imaging data. CNNs take advantage of spatial relationships

naturally found in image and spectral data with learned convolutional filters, which facilitate the extraction of features at multiple levels of abstraction without the need for feature engineering by hand.

But, recurrent neural networks with their various types (LSTMs, GRUs) are the most appropriate choice for time-series data like time-resolved photoluminescence decays where the temporal correlations that reveal physical properties are of paramount importance[22–24].

3.2 Generative Models

Generative adversarial networks (GANs) and variational autoencoders (VAEs) are still relatively unexplored methods in quantum dot (QD) research that thanks to the ability to generate QD structures, compositions and functionalization schemes of surfaces with desired properties can assist human researchers in finding potentially useful materials based on human intuition and knowledge combined with generative models[25,26].

These models learn the statistical distribution of configurations of QD that have been experimentally observed and Because of this can suggest candidates that meet property constraints that are multiple and simultaneous - a feature that becomes more and more critical as the demands for applications become more specific.**Fig.1** Shows the overview of machine learning models[27,28].




Fig.1 Shows the overview of machine learning models; Data-driven, computationally intensive; limited interpretability

4. Machine Learning for QD Synthesis Optimization
4.1 Reaction Parameter Optimization
 Using machine learning to train the synthesis process, leads to long sequences of data beyond what can be considered over of data sets. It generates diverse and potential areas of integration between ML and QD. Initial research showed that quite simple machine learning models, which were trained on data sets made by conventional experimental campaigns, could predict the emission wavelengths and quantum yields of QDs from synthesis parameters with high enough accuracy to be used for guiding optimization of the

protocol. Later on, other research works presented active learning systems that physically connected prediction and experimentation, allowing the autonomous optimization of the process before human intervention in the experimental cycles[29,30].

A milestone demonstration was the integration of machine learning (ML) and automated synthesis platforms. The researchers merged a robotic synthesis platform capable of performing parameterized synthesis protocols with a Bayesian optimization algorithm[31]. Through this, they managed to identify near-optimal synthesis conditions for CdSe, InP, and perovskite QDs, in just one to two weeks of continuous automated operation - these are timescales that would correspond to months of manual experimentation. Besides, automated platforms are capable of generating a lot more consistent and reproducible data than manual synthesis, thereby improving the quality of the training data and the reliability of the learned models[32,33].

The number of dimensions in synthesis parameter spaces that can be handled with ML is still getting larger and larger. Initially, some of the first studies only dealt with optimizing two to four parameters together; But, nowadays top-notch platforms are capable of optimizing ten or even more parameters, also incorporating fine-tuned factors like injection rate profiles and temperature ramp schedules which are hard to manually control and analyze[34].

4.2 Composition and Surface Chemistry Engineering

ML has demonstrated its excellence in dealing with the composition complexity of ternary and quaternary QD systems, including elemental substitution designed QDs like CdSSe, ZnCdS and InGaP, and also core/shell arrangements such as CdSe/ZnS and InP/ZnSe. The dependency of composition on properties in such systems is strongly nonlinear and very steep to guess from the first principles without performing highly compute extensive DFT calculations. ML models formed by experimental compositional datasets can quite precisely foresee optical and electronic properties according to composition which is a plus for exploratory synthesis, to a big extent decreasing the experimental drive for finding promising compositions[35,36].

Surface chemistry involving ligand identity, binding mode, surface stoichiometry and passivation quality drastically affects QD optical properties, stability and processability. Though, it is also one of the least understood aspects of QD science. Machine learning methods like graph neural network that model the molecular structure

of surface ligands explicitly are gradually making it possible to quantitatively predict the effect of surface chemistry on QD properties. This way, it is becoming feasible to design ligands based on rational considerations[37].

5. ML for QD Characterization

5.1 Spectral Analysis and Decomposition

The photoluminescence spectrum of a polydisperse QD ensemble results from the combination of the size-dependent emission spectra of the individual QD populations. Traditionally, obtaining the QD size distribution details from ensemble spectra required either a laborious electron microscopy characterization which only samples a tiny fraction of the material or empirical correlations between spectral features and size that are quite system-specific and of limited accuracy. Machine learning methods have revolutionized this type of analysis[38]. Convolutional neural networks trained on simulated or experimentally characterized spectra databases can deduce size distribution parameters from ensemble spectra as accurately as TEM characterization, but in a matter of seconds instead of hours. More advanced methods also combine spectral machine learning analysis with time-resolved photoluminescence data to derive size, composition, and surface passivation features simultaneously from a single multidimensional measurement[39,40].

Raman spectroscopy can offer additional structural details but the spectra it produces are so complex that it is almost impossible for a person to interpret them manually. Yet, machine learning has now been used to assist the interpretation of Raman spectra. For example, through this method, it has become possible to automatically recognize different crystalline phases of QDs, identify surface oxidation states and understand how ligands are attached to the QD surface. The sensitivity of the method is so high that even slight structural alterations due to environmental exposure or running the device can be detected[41,42].

5.2 Imaging and Morphological Characterization

Transmission electron microscopy is still the best technique for detailed visualization of QD morphology, but manual analysis of TEM images is very time-consuming. One TEM session alone may produce thousands of images and manually measuring a few hundred particles can take an expert several hours. ML-based image analysis, using convolutional neural networks trained on annotated TEM image databases, can automatically detect, segment and measure QD particles in TEM images at a pace and level of consistency beyond human capabilities. Besides simple size measurement, ML-powered TEM analysis can

derive shape parameters, recognize exposed crystal facets, distinguish core/shell interfaces and measure defect densities morphological characteristics that are very time-consuming to measure manually but are the most defining features of the materials' optical and electronic properties. The use of ML-based TEM analysis in automated characterization pipelines leads to the possibility of real-time morphological feedback during synthesis, Because of this allowing closed-loop control of QD morphology even remotely[43].

6. ML for QD Device Optimization

6.1 Quantum Dot LEDs

QLEDs are one of the most commercially viable QD device technologies that are currently being considered for applications in both high-color gamut displays and general illumination. The efficiency of QLEDs is affected by a complicated mix of factors like the thickness and uniformity of the QD layer, the properties of the charge transport layers, the matching of electrode work functions, and the encapsulation of the device. Due to the large number of interconnected variables that need to be optimized and the lengthy nature of device fabrication and characterization, QLED optimization is an ideal candidate for ML-guided methods [44].

Machine learning models developed based on QLED fabrication and performance data have shown that they can predict device efficiency and lifetime from fabrication parameters with the accuracy needed to aid process optimization. Transfer learning techniques, which make use of models that have been pre-trained on larger datasets of related device technologies (organic LEDs, perovskite LEDs), have been employed to solve the problem of small dataset sizes typical of QLED research, allowing accurate predictions to be made from just a few dozen experimental data points. Fig.2 shows the various quantum dot device optimizations[45].




Fig.2 Various quantum dot device optimizations

6.2 Quantum Dot Solar Cells

Enhancing the efficiency of quantum dot solar cells hinges on controlling the whole process from light absorption through to the recombination of losses over length scales from single quantum dots (a few

nanometers) to entire devices (micrometers). ML has been implemented at different levels of this multi-scale optimization problem, e.g. predicting the optical absorption spectra of quantum dot films as functions of QD size and packing density to optimizing whole device architectures through high-throughput simulation complemented by ML surrogate modeling[46].

One of the most remarkable applications of ML has been the optimization of QD surface passivation for solar cell applications. The velocity of surface recombination of QD films, which is affected by the density and energy distribution of surface trap states, is a key thing in determining the open-circuit voltage and Because of this the device efficiency. ML models that forecast the density of surface traps based on ligand exchange conditions and structural parameters of QDs have led to the creation of passivation strategies that have resulted in the QD solar cell efficiencies exceeding 18%[38].

6.3 Quantum Information Applications

Quantum dots (QD) are among the most promising candidates for solid-state single-photon emitters and spin qubits, which are essential components of quantum information technologies. The standards for these applications are extremely high: single-photon purity should be more than 99%, photon indistinguishability more than 95%, and spin coherence times should be long enough for quantum gate operations. Achieving these standards necessitates outstanding control of the QD electronic structure, strain environment, and photonic environment. Machine learning has been leveraged in various areas such as the optimization of QD positioning inside photonic crystal cavities to maximize Purcell enhancement of emission, the prediction of QD spin coherence times based on structural and compositional parameters, and the automated tuning of QD emission wavelengths through strain engineering. Although these applications are still at the initial stage, they mark an important frontier where machine learning could A lot contribute to the realization of practical quantum information devices[37,47].

7. Conclusion

Incorporating machine learning in quantum dot studies is probably the most important change in how things are done in the history of this field. Machine learning is opening up new possibilities in synthesis optimization, spectral characterization, device engineering or quantum information applications. Bayesian optimization charts high-dimensional synthesis parameter spaces with a very small number of experiments compared to traditional ones. Deep learning is capable of rapidly and accurately extracting quantitative structural and

compositional information from spectral and imaging data.

Works Cited

- [1] Cotta MA. Quantum Dots and Their Applications: What Lies Ahead? *ACS Appl Nano Mater* 2020;3:4920–4. <https://doi.org/10.1021/ACSANM.0C01386>.
- [2] Li Q, Wu K, Zhu H, Yang Y, He S, Lian T. Charge Transfer from Quantum-Confined 0D, 1D, and 2D Nanocrystals. *Chem Rev* 2024;124:5695–763. <https://doi.org/10.1021/ACS.CHEMREV.3C00742>.
- [3] Nasrollahpour H, Sánchez BJ, Sillanpää M, Moradi R. Metal Nanoclusters in Point-of-Care Sensing and Biosensing Applications. *ACS Appl Nano Mater* 2023;6:12609–72. <https://doi.org/10.1021/ACSANM.3C01569>.
- [4] Mocci F, Engelbrecht LDV, Olla C, Cappai A, Casula MF, Melis C, et al. Carbon Nanodots from an In Silico Perspective. *Chem Rev* 2022;122:13709–99. <https://doi.org/10.1021/ACS.CHEMREV.1C00864>.
- [5] Basha SA, Cole S, Divakaran D, Suyambulingam I. Solid State Reaction Method for Nanomaterials Synthesis: A Comprehensive Review on Characterization, Properties, and Applications. *Springer Proceedings in Materials*, vol. 60, Springer; 2024, p. 667–89. https://doi.org/10.1007/978-981-97-7071-7_47.
- [6] Ros H, Chan N, Cook MT, Shorthouse D. Artificial intelligence and machine learning guided optimization in drug delivery. *Adv Drug Deliv Rev* 2026;232:115781. <https://doi.org/10.1016/J.ADDR.2026.115781>.
- [7] Ogundeji TS, Alamu GA, Oyewole OJ, Johnson MJ, Olaoye AO, Vyas S, et al. Review of recent progress in enhanced quantum dot solar cells via plasmonic nanostructures. *Next Materials* 2026;11:101599. <https://doi.org/10.1016/J.NXMATE.2026.101599>.
- [8] Mimona MA, Rimon MIH, Zohura FT, Sony JM, Rim SI, Arup MMR, et al. Quantum dot nanomaterials: Empowering advances in optoelectronic devices. *Chemical Engineering Journal Advances* 2025;21:100704. <https://doi.org/10.1016/J.CEJA.2025.100704>.
- [9] Fang W, Yu J, Liao S, Gao H, Li X, Liu B. Emergence of colloidal quantum well-based optoelectronics. *Cell Rep Phys Sci* 2024;5:101936. <https://doi.org/10.1016/J.XCRP.2024.101936>.
- [10] Munyebvu N, Lane E, Grisan E, Howes PD. Accelerating colloidal quantum dot innovation with algorithms and automation. *Mater Adv* 2022;3:6950–67. <https://doi.org/10.1039/D2MA00468B>.
- [11] Divakaran D, Sriariyanun M, Basha SA, Suyambulingam I, Sanjay MR, Siengchin S. Physico-chemical, thermal, and morphological characterization of biomass-based novel microcrystalline cellulose from *Nelumbo nucifera* leaf: Biomass to biomaterial approach. *Biomass Convers Biorefin* 2023. <https://doi.org/10.1007/s13399-023-04349-9>.
- [12] Chafiq M, Chaouiki A, Ko YG. Recent Advances in Multifunctional Reticular Framework Nanoparticles: A Paradigm Shift in Materials Science Road to a Structured Future. *Nano-Micro Letters* 2023 15:1 2023;15:213–. <https://doi.org/10.1007/S40820-023-01180-9>.
- [13] Mourdikoudis S, Pallares RM, Thanh NTK. Characterization techniques for nanoparticles: comparison and complementarity upon studying nanoparticle properties. *Nanoscale* 2018;10:12871–934. <https://doi.org/10.1039/C8NR02278J>.
- [14] K.Daniel, K.Venkatarao, K.Srinivasarao, Shaik Azad Basha, Sandhya Cole. Structural and Optical bandgap studies of undoped and Sm³⁺ doped MgZn₂(PO₄)₂ nanopowders. *ANU Journal of Physical Sciences* 2021;13:53–60.
- [15] Cheng Y, Wan H, Liang T, Liu C, Wu M, Hong H, et al. Continuously Graded Quantum Dots: Synthesis, Applications in Quantum Dot Light-Emitting Diodes, and Perspectives. *J Phys Chem Lett* 2021;12:5967–78. <https://doi.org/10.1021/ACS.JPCLETT.1C01554>.
- [16] Chernomordik BD, Marshall AR, Pach GF, Luther JM, Beard MC. Quantum Dot Solar Cell Fabrication Protocols. *Chemistry of Materials* 2017;29:189–98. https://doi.org/10.1021/ACS.CHEMMATE.R.6B02939/SUPPL_FILE/CM6B02939_SI_001.AVI.
- [17] Chandra M, Nookaraju M, Sharma VK, Somasekhar R. Influence of Vanadium incorporated mesoporous silica on the decolorization of orange G under visible light irradiation. *Inorganic and Nano-Metal*

- Chemistry 2022;52:387–96.
<https://doi.org/10.1080/24701556.2021.1980019>.
- [18] Kumar VN, Nookaraju M, Jella KS, Nalla S. Synthesis and Biological Evaluation of 1,3,4-oxadiazole ring incorporated (pyrimidin-5-yl)indolizine as Anticancer Agents. *Chemical Data Collections* 2025;57:101190.
<https://doi.org/10.1016/j.cdc.2025.101190>.
- [19] Malashin I, Martysyuk D, Nelyub V, Borodulin A, Gantimurov A, Tynchenko V. A review of applications of machine learning in quantum dots research. *Discover Nano* 2026;21.
<https://doi.org/10.1186/S11671-026-04466-0>.
- [20] Balasubramanyam SV, Rao KS, Daniel K, Basha SA, Cole S. Structural and Optical Insights of Sm³⁺ Doped MgCdP2O7 Nanopowders for Solid-State Lighting Applications. *Physical Chemistry Research* 2025;13:463–78.
<https://doi.org/10.22036/PCR.2025.507953.2645>.
- [21] Danish A, Feng Z, Liu L, Zhang S-H, Khan K, Rahman IU, et al. Tree-based and deep learning models for landslide susceptibility assessment with imbalanced data. *Journal of Rock Mechanics and Geotechnical Engineering* 2026.
<https://doi.org/10.1016/J.JRMGE.2026.01.042>.
- [22] Dr. Betshrine Rachel Jibinsingh. RECENT DEVELOPMENTS IN CONVOLUTIONAL NEURAL NETWORKS FOR IMAGE-BASED TASKS. INTERNATIONAL RESEARCH ON SUSTAINABILITY AND INNOVATION: A MULTIDISCIPLINARY APPROACH TO GLOBAL DEVELOPMENT 2022.
<https://doi.org/10.25215/9349154692.41>.
- [23] Nandhini VL, Nagabhooshanam N, G BMR, Verma R, Prasan UD, ArunaKumari S, et al. Biosynthesis of ZrO₂ from *Phyllanthus niruri* extracts and its PVA composite: mechanical, wear, dielectric, water absorption, and thermal conductivity properties. *Biomass Conversion and Biorefinery* 2024 15:9 2024;15:14137–45.
<https://doi.org/10.1007/s13399-024-06046-7>.
- [24] Nookaraju M, Kumar Nadikatla S, Krishna V, Siva Sankar M, Nagaraju M, Triveni Y, et al. ONE-POT, GREEN SYNTHESIS OF PYRIDINE-2-CARBALDEHYDE BASED CHALCONES BY FUNCTIONALIZED MCM-41 CATALYSTS AND THEIR ANTIMICROBIAL EVALUATION. *Rasayan J Chem* 2080;18:2025.
<https://doi.org/10.31788/RJC.2025.1849386>.
- [25] Carbonera M, Ciavotta M, Messina E. Variational Autoencoders and Generative Adversarial Networks for Multivariate Scenario Generation. *Data Science for Transportation* 2024;6.
<https://doi.org/10.1007/S42421-024-00097-Y>.
- [26] Kumar JS, Lakshmi NR, Neeraja K, Devi TA, Swamy DT, Basha SA, et al. Spectroscopic and Optical Studies of Molybdenum Ions in CdO-SrO-B2O3-SiO₂ (CdSBSi) Glass System. *Asian Journal of Chemistry* 2025;37:1204–10.
<https://doi.org/10.14233/AJCHEM.2025.33684>.
- [27] Saleh KA, Mustafa WW, M. M. R, Ray S, Qassem TA, Krithiga T, et al. Electronic–photonic interplay in nitrogen-doped MXene quantum dots: mechanistic insights into dual-mode and multiplexed sensing. *RSC Adv* 2026;16:26585–608.
<https://doi.org/10.1039/D6RA02859D>.
- [28] Chandolu N, Yeti DR, Pavan Kumar MVS, Rajendra K, Gosu SJ, Dhyva Stuthi P, et al. Machine Learning for Analyzing the Biowaste to Wealth Approach: A briefreview. *International Journal of Drug Delivery Technology* 2026;16:789–94.
<https://doi.org/10.25258/IJDDT.16.11S.80>.
- [29] Malashin I, Martysyuk D, Nelyub V, Borodulin A, Gantimurov A, Tynchenko V. A review of applications of machine learning in quantum dots research. *Discover Nano* 2026 21:1 2026;21:54-.
<https://doi.org/10.1186/S11671-026-04466-0>.
- [30] Rani GR, Kuna S, Kodamala D, Basha SA, Cole S. Structural, optical and luminescence studies of CuO doped MgZn₂(PO₄)₂ nanopowder variants for solid state optical devices. *Next Materials* 2025;8:100819.
<https://doi.org/10.1016/J.NXMATE.2025.100819>.
- [31] Cole S, Basha SA, Naveen Kumar B V, Srinivasarao K, Daniel K. Eu³⁺ Ion-doped Lithium Cadmium Orthophosphate Nanopowders for Reddish Orange-emitting Solid-state Lighting Applications. *Physical Chemistry Research* 2026;14:173–94.
<https://doi.org/10.22036/PCR.2025.539879.2719>.
- [32] Adelusi DevOps BS, Osamika D, Chinyeaka Kelvin-Agwu M, Mustapha AY, Forkuo AY, Ikhalea N. A Machine Learning-Driven Predictive Framework for Early Detection and Prevention of

- Cardiovascular Diseases in U.S. Healthcare Systems. *Engineering and Technology Journal* 2025;10. <https://doi.org/10.47191/ETJ/V10I05.05>.
- [33] Srinivasarao Kuna, Shaik Azad Basha, Sandhya Cole. Methods for Designing and Improving Luminescent Nanophosphors for LED Applications Using Machine Learning. National Seminar on Transforming India; Innovations, Challenges and Opportunities n.d.;1:138–59.
- [34] Pai SM, Shah KA, Sunder S, Albuquerque RQ, Brütting C, Ruckdäschel H. Machine learning applied to the design and optimization of polymeric materials: A review. *Next Materials* 2025;7:100449. <https://doi.org/10.1016/J.NXMATE.2024.100449>.
- [35] Swart I, Liljeroth P, Vanmaekelbergh D. Scanning probe microscopy and spectroscopy of colloidal semiconductor nanocrystals and assembled structures. *Chem Rev* 2016;116:11181–219. <https://doi.org/10.1021/ACS.CHEMREV.5B00678>.
- [36] Balasubramanyam SV, Srinivasarao K, Azad Basha S, Sandya Cole. Investigation on the structural and morphological properties of magnesium cadmium pyrophosphate nanopowders doped with Pr³⁺ ions. *Research Methodology in Materials Science (YVNRGDC)* 2025:145–9.
- [37] Nguyen HA, Dou FY, Park N, Wu S, Sarsito H, Diakubama B, et al. Predicting Indium Phosphide Quantum Dot Properties from Synthetic Procedures Using Machine Learning. *Chemistry of Materials* 2022;34:6296–311. <https://doi.org/10.1021/ACS.CHEMMATE.R.2C00640>.
- [38] Xu G, Zeng S, Zhang B, Swihart MT, Yong KT, Prasad PN. New Generation Cadmium-Free Quantum Dots for Biophotonics and Nanomedicine. *Chem Rev* 2016;116:12234–327. <https://doi.org/10.1021/ACS.CHEMREV.6B00290>.
- [39] Ardévol Martínez F, Min M, Kamp I, Palmer PI. Convolutional neural networks as an alternative to Bayesian retrievals for interpreting exoplanet transmission spectra. *Astron Astrophys* 2022;662. <https://doi.org/10.1051/0004-6361/202142976>.
- [40] Balasubramanyam S V., Kumar BVN, Srinivasarao K, Basha SA, Cole S. Spectroscopic and photoluminescent studies of Dy³⁺ doped MgCdP2O7 nanopowders for white light emitting devices. *Journal of Materials Science: Materials in Electronics* 2026 37:14 2026;37:1098-. <https://doi.org/10.1007/S10854-026-17475-7>.
- [41] Sunil M, Pallikkavaliyaveetil N, N MiI, Gopinath A, Chidangil S, Kumar S, et al. Machine learning assisted Raman spectroscopy: A viable approach for the detection of microplastics. *Journal of Water Process Engineering* 2024;60:105150. <https://doi.org/10.1016/J.JWPE.2024.105150>.
- [42] Basha SA, Srinivasarao K, Daniel K, Kumar BVN, Rajeev YN, Cole ProfS. Orthorhombic crystal structured Dy³⁺ doped Li₂Cd₂(PO₄)₂ nanopowders for yellowish light emitting solid state lighting applications 2026. <https://doi.org/10.21203/RS.3.RS-8763095/V1>.
- [43] Smeaton MA, Abellan P, Spurgeon SR, Unocic RR, Jungjohann KL. Tutorial on In Situ and Operando (Scanning) Transmission Electron Microscopy for Analysis of Nanoscale Structure–Property Relationships. *ACS Nano* 2024;18:35091–103. <https://doi.org/10.1021/ACS.NANO.4C09256>.
- [44] Li M, Bao H, Gao P, Liang W, Lin X, Wu L, et al. The Stability Challenges in Developing QLED Based Display Technology. *J Phys Chem Lett* 2025;16:10058–70. <https://doi.org/10.1021/ACS.JPCLETT.5C02156>.
- [45] Chen C, Lin X, Wu XG, Bao H, Wu L, Hu X, et al. Machine Learning Assisted Stability Analysis of Blue Quantum Dot Light-Emitting Diodes. *Nano Lett* 2023;23:5738–45. https://doi.org/10.1021/ACS.NANOLETT.3C01491/SUPPL_FILE/NL3C01491_SI_003.AVI.
- [46] Zhang Z, Wang W, Rao H, Pan Z, Zhong X. Improving the efficiency of quantum dot-sensitized solar cells by increasing the QD loading amount. *Chem Sci* 2024;15:5482–95. <https://doi.org/10.1039/D3SC06911G>.
- [47] Xu Z, Huang Z, Li C, Huang T, Evangelista FA, Tang ML, et al. Tuning the Quantum Dot (QD)/Mediator Interface for Optimal Efficiency of QD-Sensitized Near-Infrared-to-Visible Photon Upconversion Systems. *ACS Appl Mater Interfaces* 2020;12:36558–67. <https://doi.org/10.1021/ACSAMI.0C10269>.

