

Evaluation of changes in intraocular lens power following corneal refractive surgery

Dr Sujata R Charel¹, Dr Vishva Sheth^{2*}, Dr Jeet Patel³

¹Associate Professor, (MS Ophthalmology) Department of Ophthalmology, SBKS Medical College and Research Centre, Sumandeep Vidyapeeth deemed to be university, Piparia Village, Taluka: Waghodia, District: Vadodara, Gujarat - 391760, India. Ph. No. 9726253089.
Email: sujatacharel117@gmail.com. ORCID: 0000-0002-1856-9946

²*2nd Year Resident (MS Ophthalmology) Department of Ophthalmology, SBKS Medical College and Research Centre, Sumandeep Vidyapeeth deemed to be university, Piparia Village, Taluka: Waghodia, District: Vadodara, Gujarat - 391760, India. Ph. No. 6354130044. Email: vishvasheth11@gmail.com

³Alumini, (MS Ophthalmology) Department of Ophthalmology, SBKS Medical College and Research Centre, Sumandeep Vidyapeeth deemed to be university, Piparia Village, Taluka: Waghodia, District: Vadodara, Gujarat - 391760, India.

***Corresponding author: Dr Vishva Sheth, 2nd Year Resident (MS Ophthalmology) Department of Ophthalmology, SBKS Medical College and Research Centre, Sumandeep Vidyapeeth deemed to be university, Piparia Village, Taluka: Waghodia, District: Vadodara, Gujarat - 391760, India**

Ph. No. 6354130044 | Email: vishvasheth11@gmail.com

Conflicts of interest: There are no conflicts of interest.

Financial support and sponsorship: Nil.

I declare that no artificial intelligence tools were used in the creation of this article, and all content has been independently written by the author.

Ethical approval: The research/study was approved by the Institutional Review Board at the Institutional Ethics Committee, approval number SVIEC/ON/MEDI/BNPG18/019224, dated 27th December 2019.

Informed patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patients have given their consent for their clinical information to be reported in the journal. The patients understand that their names and initials will not be published and due efforts will be made to conceal their identity, but anonymity cannot be guaranteed.

Author contributions:

Dr. Sujata R Charel conceptualized the study design, provided overall supervision, and critically reviewed the manuscript for important intellectual content and also conducted patient recruitment, performed all clinical examinations, analysed the data, and drafted the initial manuscript.

Dr. Vishva Sheth coordinated laboratory investigations, performed statistical analysis validation, and revised the manuscript for final submission.

Dr. Jeet Patel assisted with data collection, maintained the study database, and contributed to literature review.

The authors would like to thank all the patients for their participation in the study.

Received: 28th Feb, 2026; **Revised:** 6th March, 2026; **Accepted:** 7th April, 2026; **Available Online:** 20th April, 2026

ABSTRACT

Introduction

LASIK is one of the most commonly performed refractive surgeries worldwide because of patient's comfort and surgeons skills. This young generation undergoing LASIK surgery will also have to undergo cataract surgery in their old age. Patients undergoing cataract surgery require intraocular lens (IOL) calculation for the IOL implantation. As LASIK changes the corneal curvatures the keratometry and effective lens position will be affected leading to change in IOL power. This study aims to identify the change in IOL power after LASIK surgery.

Method

We performed a prospective and observational study on 75 patients (150 eyes) undergoing LASIK surgery, who met inclusion criteria. All the patient were categorized on basis of degrees of myopia and all went under same preoperative and postoperative treatment and data was collected using performa. All the collected data were used to see the change in IOL power before and after LASIK.

Conclusion

IOL power after LASIK surgery changes significantly due to changes occurring at the level of cornea. Higher the degrees of myopia more the difference in IOL power was be seen. The mean increase change in IOL power among 150 eyes using SRK I was 2.95 D \pm 1.20, while in SRK II, SRK T, Hoffer Q, Holladay and Binkhorst was 3.03 D \pm 1.41, 3.45 D \pm 1.45, 4.73 D \pm 1.91, 3.91 D \pm 1.55 and 4.57 D \pm

RESEARCH PAPER

1.80 respectively. Over all maximum change was seen in Hoffer Q and minimum change in SRK I formula. All the formulae mentioned may produce residual refractive errors, for that the patient has to undergo cataract surgery, but we have only taken the postoperative LASIK patients so it cannot be confirmed which formula yields the least residual refractive error after cataract surgery in post LASIK eyes.

Keywords: LASIK, intraocular lens, IOL power, corneal refractive surgery, cataract surgery, myopia.

How to cite this article: Charel SR, Sheth V, Patel J. Evaluation of changes in intraocular lens power following corneal refractive surgery. *Int J Drug Deliv Technol.* 2026;16(61s):1717-1724. DOI: 10.25258/ijddt.16.61s.194

Source of support: Nil.

Conflict of interest: None

INTRODUCTION

Uncorrected refractive errors (URE) is defined as a presenting visual acuity of less than 6/12 in the better eye with an improvement of at least 0.2 logMAR (equivalent to two lines) after refraction.⁽¹⁾ Refractive error (RE) is one of the most common ocular conditions affecting all age groups and a priority under the VISION 2020 initiative. Worldwide, uncorrected refractive errors are one the major causes of visual impairment.⁽²⁾ Around 12.8 million population in age group of 5-15 years were suffering from uncorrected or inadequately corrected refractive errors, which leads to global prevalence of 0.96%.⁽³⁾

Refractive errors like myopia, hypermetropia and astigmatism are the commonest causes of defective vision. Most REs are easily corrected by the help of spectacles at the primary care level. In spite of the availability of a cost effective intervention, uncorrected refractive error (URE) is a leading public health challenge. Defective visual and blindness which are caused by URE in adults leads to impact severely on their social & economic well-being, which causes limitation in their educational and employment opportunities of economically active persons.⁽⁴⁾

Among all the REs, myopia is the commonest RE. In myopic eyes, the parallel rays of light coming from infinity tend to focus anterior to the retina, leading to diminution in vision. In myopia the cornea and the lens refract the rays too much for the given axial length or the axial length might be too much for normal refracting optical system. For correction of myopia the commonly used methods include spectacles and contact lens. These methods provide temporary solution for correction. They both have certain functional limitations such as the problems encountered while wearing glasses in rainy season, while playing sports, limited field of vision and chromatic aberration while using higher power. Contact lens also have inconvenience of carrying the solutions, storage containers and might also cause sight threatening corneal infection.⁽⁵⁾ There are

certain surgical procedures for correcting myopia. These procedures involve an operation over cornea (corneal refractive surgeries) or lens (lenticular refractive surgeries), they work by reducing the focusing power of cornea or lens.

However in recent times there is an emerging trend of correcting refractive errors with surgical procedures like laser assisted in-situ keratomileusis (LASIK) and photorefractive keratectomy (PRK) which are corneal refractive surgeries.⁽⁶⁾ Overall, demand for laser refractive surgeries is expected to rise at annual rate of 7.7% from 2009 to 2014 and the number of procedures increasing from 3.4 million to 4.9 million. The majority of growth is occurring in the developing countries of Asia. In India refractive surgeries have also taken pace with 1,28,000 surgeries or more per year.⁽⁷⁾ These procedures cause changes in many aspects of the cornea like corneal curvature, surface, rigidity and thickness in addition to mild changes in axial length of the eye.⁽⁸⁾ Refractive surgery is generally done in younger age group. However as the age advances all of these patients will develop cataract and need cataract surgery. Cataract extraction along with intra ocular lens (IOL) implantation is the only way to manage cataracts.

To calculate IOL power for implanting IOL after extraction of cataract, some parameters like keratometry, anterior chamber depth and axial length of the eye ball need to be measured and then IOL power is calculated by using various formulae.⁽⁹⁾ Intraocular lens (IOL) implantation after refractive surgery is challenging because standard IOL power formulae leave some significant errors in calculation of IOL power and prevents achievement of postoperative emmetropia.⁽¹⁰⁾

The aim of the present study is to study the changes occurring in the calculation of IOL power with commonly used formula in patients after corneal refractive surgery.

AIM

RESEARCH PAPER

- To study the change in IOL power calculation after Laser Assisted in-situ Keratomileusis (LASIK) surgery.

OBJECTIVES

- To study change in corneal keratometry, preoperative and postoperative LASIK surgery.
- To compare various IOL calculation formula results in LASIK surgery cases.

MATERIALS AND METHODS

Study design

- Study was a prospective and observational.

Study period

- The date of approval from ethics committee to 2 years or completion of sample size, whichever is the earliest.

Sample size

- Total number of 75 cases (150 eyes), who are undergoing Laser Assisted In-Situ Keratomileusis (LASIK) surgery. Study setting
- The entire study was conducted in Ophthalmology department, Tertiary hospital Gujarat.

Inclusion criteria

- Age of >18 years to < 40 years
- Refractive power of >-0.5 D to <-8.00 D
- Patients willing to participate
- Patients undergoing only LASIK surgery
- Having stable refraction for atleast one year
- Atleast 1 year of contact lens free period.

Exclusion criteria

- Monocular patients
- Age of <18 years and >40 years
- Patients undergoing refractive surgery other than LASIK
- History of any ocular trauma
- Extended use of contact lens
- History of herpes keratitis
- Severe dry eye or any kind of tear film abnormalities
- Ocular diseases like blephritis, meibomian gland dysfunction, corneal opacities, corneal dystrophies, allergic conjunctivitis etc
- Patients with projected residual corneal stromal bed thickness after ablation is less than 270 microns
- Glaucoma patients
- Keratoconus
- Autoimmune disorders
- Post LASIK corneal ectasia or any kind of post LASIK complications like epithelial ingrowth, keratitis etc
- Patient with posterior segment abnormalities like lattice with or without hole
- Patients not willing for participation
- Patients with unreasonable expectations
- Patient who lost follow ups

RESULTS

This was a clinical, interventional and prospective study held in Department of ophthalmology, at tertiary care hospital Gujarat, in which we examined 150 eyes of 75 patients. Among them male were 43 (57.33%) and female were 32 (42.67%).

The age groups of the patients which are included in are study was divided in 4 groups. Among which 14 (18.67%) patients were ≤ 20 years of age, 34 (45.33%) patients were between 21-25 years, 21 (28%) were in 26-30 years of age group and only 6 (8%) were in >31 years of age group.

The mean age in male group was 24.67 ± 4.21 and in female group was 24.13 ± 3.40 . Thus the mean age was 24.44 ± 3.88 and it was not statistically significant difference between them (p value 0.378).

As shown in Table 1, preoperative mean K1 value for mild myopia was $43.67D \pm 1.39$ which changed to $41.52 D \pm 1.31$ and remained constant at 1 week and 1 month postoperative respectively. In moderate myopic cases, mean K1 was $43.85 D \pm 1.28$ preoperatively, which was reduced to $39.93 D \pm 1.63$ and was constant at 1 week and 1 month. In high myopic cases, preoperative mean K1 was $43.33 D \pm 1.41$ and reduced to $38.78 D \pm 1.09$ which was constant on follow-ups. Thus there was significant difference in preoperative and postoperative K1 values after LASIK surgery in all groups (p value 0.001).

The average reduction of K1 in all our cases after LASIK was $3.19 D \pm 1.32$, while average reduction of K1 value was $2.18 D \pm 0.58$, $3.92 D \pm 1.17$ and $4.61 D \pm 1.11$ in mild myopic eyes, moderate myopic and high myopic eyes respectively. We can note that there was more reduction in K1 values (flattening) after LASIK surgery with higher degree of myopia.

Table 1: Comparison between preoperative and postoperative K1 (Flat Keratometry) in different groups of myopia

	Mild		Moderate		High	
K1 Diopters	Mean	Std. deviation	Mean	Std. deviation	Mean	Std. deviation
Preoperative	43.67	1.39	43.85	1.28	43.33	1.41
Postoperative 1 week	41.52	1.31	39.93	1.63	38.78	1.09

RESEARCH PAPER

Postoperative month	1	4	1.3	39.9	1.6	3	8.7	1.0
p value		0.001		0.001		0.001		0.001


As shown in Table 2, preoperative mean K2 value for mild myopia was 44.48 D ±1.248 which changed to 42.34 D ±1.32 and remained constant at 1 week and 1 month postoperative respectively. In moderate myopic cases, mean K2 was 44.73 D ±1.20 preoperatively, which was reduced to 40.61 D ±1.60 and was constant at 1 week and 1 month. In high myopic cases, preoperative mean K2 was 45.11 D ±1.69 and reduced to 39.67 D ±1.41 which was constant during follow-ups. Thus there was significant difference in preoperative and postoperative K2 values after LASIK surgery in all groups (p value 0.001).

The average reduction of K2 value in all our case after LASIK was 3.33 D ±1.35 and average reduction in mean K2 value in mild myopic eyes, moderate myopic and high myopic eyes was 2.24 D ±0.51, 4.05 D ±1.10 and 5.51 D ±0.76 respectively. We can note that there was more reduction in K2 values (flattening) after LASIK surgery with higher degree of myopia.

Table 2: Preoperative and postoperative K2 (steep keratometry) in different groups of myopia

K2 Diopter (D)	Mild		Moderate		High	
	Mean	Std. deviation	Mean	Std. deviation	Mean	Std. deviation
Preoperative K2	44.48	1.25	44.73	1.20	45.11	1.69
Postoperative 1 week K2	42.34	1.32	40.61	1.60	39.67	1.41
Postoperative 1 month K2	42.34	1.32	40.61	1.60	39.67	1.41

p value	0.001	0.001	0.001
---------	-------	-------	-------



Change in Preoperative and Postoperative K2

Table 3: Preoperative mean IOL power (D)

	Mild	Moderate	High
SRK I	19.88 ± 1.77	17.01 ± 1.79	15.22 ± 0.97
SRK II	20.03 ± 1.83	16.84 ± 2.06	15.33 ± 1.00
SRK T	19.94 ± 2.02	16.49 ± 2.20	14.67 ± 1.50
HOFFER Q	19.96 ± 2.57	16.19 ± 2.58	14.11 ± 1.69
HOLLADAY	19.91 ± 2.26	16.47 ± 2.29	14.67 ± 1.50
BINKHORST	19.96 ± 2.69	15.86 ± 2.78	13.78 ± 1.48
p value	0.098	0.05	0.05

Table 3 shows the mean intraocular lens power (IOL) of preoperative patients in various degrees of myopia using various formulae. There was no significant difference between various formulae in mild myopic cases (p value 0.098). While there was some significant difference found in moderate and high myopia (p value 0.05 & 0.05 respectively).

Table 4: Postoperative mean IOL power (D)

Formulae	Mild	Moderate	High
SRK I	21.94 ± 1.80	20.65 ± 1.70	20.0 ± 1.23

RESEARCH PAPER

SRK II	22.07 ± 1.79	20.65 ± 2.04	19.44 ± 1.33
SRK T	22.25 ± 1.88	20.84 ± 1.87	19.78 ± 1.48
HOFFER Q	23.19 ± 2.46	21.91 ± 2.23	21.56 ± 1.59
HOLLADAY	22.63 ± 2.07	21.27 ± 2.02	20.67 ± 1.41
BINKHORST	23.22 ± 2.55	21.41 ± 2.35	20.67 ± 1.73
p value	0.001	0.001	0.001

As shown in Table 4, after LASIK surgery there was variation in IOL power within different formulae of same group. IOL power in mild myopic cases after LASIK was 21.94 D ± 1.80, 22.07 D ± 1.79, 22.25 D ± 1.88, 23.19 D ± 2.46, 22.63 D ± 2.07 and 23.22 ± 2.55 in SRK I, SRK II, SRK T, Hoffer Q, Holladay and Binkhorst formula. The same was applicable in moderate and high groups. In postoperative cases there was significant difference among all the formulae in all the groups. (p value 0.001) (Table 4)

Table 5: Comparison of mean IOL power (D) before and after LASIK in Mild myopic eyes

	Preoperative	Postoperative	Mean increase in power of IOL	p value
SRK I	19.88 ± 1.77	21.94 ± 1.80	2.05 ± 0.52	0.001
SRK II	20.03 ± 1.83	22.07 ± 1.79	2.04 ± 0.46	0.001
SRK T	19.94 ± 2.02	22.25 ± 1.88	2.31 ± 0.54	0.001
HOFFER Q	19.96 ± 2.57	23.19 ± 2.46	3.19 ± 0.67	0.001
HOLLADAY	19.91 ± 2.26	22.63 ± 2.07	2.72 ± 0.60	0.001

BINKHO RST	19.96 ± 2.69	23.22 ± 2.55	3.20 ± 0.86	0.001
------------	--------------	--------------	-------------	-------

In mild myopic eyes preoperative IOL power using SRK I formula was 19.88 D ± 1.77 and postoperative it was 21.94 D ± 1.80. So the mean difference of IOL power using SRK I formula 2.05 D ± 0.52. The change was significant (p value 0.001). Similarly, the mean difference of diopters in SRK II, SRK T, Hoffer Q, Holladay and Binkhorst was 2.04 D ± 0.46, 2.31 D ± 0.54, 3.19 D ± 0.67, 2.72 D ± 0.60 and 3.20 D ± 0.86 respectively, which proved to be significant. Maximum difference was seen in Binkhorst formula for mild myopic cases, while minimum difference was seen in SRK II.

In moderate myopic cases the mean preoperative IOL power was 17.01 D ± 1.79, 16.84 ± 2.06, 16.49 ± 2.20, 16.19 ± 2.58, 16.47 ± 2.29 and 15.86 ± 2.78 using SRK I, SRK II, SRK T, Hoffer Q, Holladay and Binkhorst, while postoperative IOL power using the same formulae was 20.65 D ± 1.70, 20.65 D ± 2.04, 20.84 D ± 1.87, 21.91 D ± 2.23, 21.27 D ± 2.02 and 21.41 D ± 2.35 respectively. This resulted in mean difference of 3.55 D ± 1.07, 3.74 D ± 1.44, 4.24 D ± 1.27, 5.80 D ± 1.64, 4.74 D ± 1.38 and 5.52 D ± 1.60 respectively in moderate myopic eyes.

Table 6: Comparison of IOL power (D) before and after LASIK in Moderate myopic eyes

	Preoperative	Postoperative	Mean difference	p value
SRK I	17.01 ± 1.79	20.65 ± 1.70	3.55 ± 1.07	0.001
SRK II	16.84 ± 2.06	20.65 ± 2.04	3.74 ± 1.44	0.001
SRK T	16.49 ± 2.20	20.84 ± 1.87	4.24 ± 1.27	0.001
HOFFER Q	16.19 ± 2.58	21.91 ± 2.23	5.80 ± 1.64	0.001
HOLLADAY	16.47 ± 2.29	21.27 ± 2.02	4.74 ± 1.38	0.001
BINKHORST	15.86 ± 2.78	21.41 ± 2.35	5.52 ± 1.60	0.001

Table 7: Comparison of mean IOL power (D) before and after LASIK in High myopic eyes

	Preoperative	Postoperative	Mean difference	p value
SRK I	15.22 ± 0.97	20 ± 1.23	4.67 ± 0.79	0.001
SRK II	15.33 ± 1.00	19.44 ± 1.33	4.56 ± 0.73	0.001
SRK T	14.67 ± 1.50	19.78 ± 1.48	5.39 ± 1.05	0.001
HOFFER Q	14.11 ± 1.69	21.56 ± 1.59	7.39 ± 1.14	0.001
HOLLADAY	14.67 ± 1.50	20.67 ± 1.41	6.00 ± 1.15	0.001
BINKHORST	13.78 ± 1.48	20.67 ± 1.73	6.89 ± 1.02	0.001

Similarly in Table 7, in high myopic cases, the mean difference of IOL power after LASIK was 4.67 D ± 0.79, 4.56 D ± 0.73, 5.39 D ± 1.05, 7.39 D ± 1.14, 6.00 D ± 1.15 and 6.89 ± 1.02 respectively using the same formulae. There was significant difference in IOL power after LASIK surgery. The difference between the preoperative and postoperative IOL power was seen more in eyes with high diopters of myopia. Maximum difference of IOL power was seen in Hoffer Q while minimum was seen in SRK II formula in high myopic eyes.

DISCUSSION

This study was performed on 150 eyes of 75 patients over period of one and half year. It included patients with myopia in range of >-0.50 D to <-8.00 D. No consensus on the upper limit of myopia has been established yet, while some studies recommend myopia LASIK <-12.00 D.⁽¹¹⁾

In our study, we took 150 eyes of 75 patients, among which there were 43 (57.33%) males and 32 (42.67%) were females, thus there was mild preponderance of male over female patients, but it was not significant.

The age group of the patients which were enrolled in our study was divided into 4 groups, <=20, 20-25, 26-30 and >31 years of age. Among which 14 (18.67%), 34 (45.33%), 21 (28%) and 6 (8%) number of patients fell under the above mention groups respectively. The mean age in male group was 24.67±4.21 and in female group was 24.13±3.40 . Kato N et al⁽¹²⁾ reported myopic LASIK surgery on 779 eyes with mean age of 34.6±8.3 years. Yuen LH et

al⁽¹³⁾ performed LASIK surgery in 37,932 eyes with mean age of 33 ± 7.9 years. The average age of the patients in other studies were significantly higher compared to our study. Difference of opinion does exist regarding the upper limit of age, while most of the authors agreed that LASIK is not to be performed below age of 18 years.

All the similar studies have consensus regarding rest of the patient selection criteria like, refractive status should be stable for at-least 1 year before undergoing LASIK, lens has to be clear, and patient should understand that this procedure will not prevent presbyopia and cataract formation from developing in future.

In our study, all the eyes undergoing LASIK were differentiated in mild, moderate and high myopia on basis of preoperative refractive errors, is <-3.00 Diopters (D), -3.00 D to -6.00 D and >-6.00 D respectively. Among which 67 (44.67%) eyes had mild myopia, 74 (49.33%) eyes was under moderate myopia and only 9 (6%) eyes had high myopia. Thus the mean diopters myopia was -3.69 D ± 1.57. Kato et al underwent LASIK surgery with the mean diopters of myopia as -6.40 D ± 2.58, which was much higher than our study.

In our study the mean K1 and K2 value for mild myopic cases was 43.67 D ± 1.39 and 44.48±1.25 respectively, which reduced to 41.52 D ± 1.31 and 42.34±1.32 at 1st week and 1st month follow-up and remained constant after 1st week. Average reduction noted in K1 and K2 value after LASIK for mild myopia was 2.18 D ± 0.58 and 2.24 D ± 0.51 for moderate myopia was 3.93 D ± 1.17 and 4.05 D ± 1.10. While in Maldonado Bas A⁽¹⁴⁾ study preoperative keratometry was 44.09 D ± 1.65 which changed to 39.11 D ± 1.61 with average reduction of 4.98 D ± 0.4 for myopia up-to -5.00 D which was greater reduction in K values than in our study. This less reduction would have been due to use of newer generation of laser excimer machine which was used in our study and it can also be due to different type of flap repositioning technique.

In present study the preoperative mean K1 and K2 values were 43.33 D ± 1.41 and 45.11 D ± 1.69 which reduced to 38.78 D ± 1.09 and 39.67 D ± 1.41 respectively, which was constant on 1st week and 1st month follow-ups. The mean reduction K1 and K2 values was 4.61 D ± 1.11 and 5.51 D ± 0.76, while in Maldonado Bas A⁽¹⁴⁾ study average K value reduction was 6.78 D ± 0.8 and 8.18 D ± 0.5 was observed in myopia of -5.00 to -10.00 D and -10.00 to -15.00 D. Reduction in this study was more than observed in our study.

The topographic stability were achieved after 6 month follow-up in Maldonado Bas A⁽⁵⁰⁾ this

observed delay in achieving stability could be due to increased amount of attempted correction. It is known that after higher attempted myopic corrections, it takes longer for topography to stabilize than after smaller correction.^(15,16) This is in contrast to our study as there was no significant change keratometry values after 1 week follow-up to 1 month follow-up.

In our study of 150 eyes we found no significant difference in IOL power calculated by various formulae in preoperative mild myopic cases ($p = 0.098$), but there was some significant difference between the formulae used in moderate and high myopic cases.

In our study, mild myopic eyes had IOL power of $19.96 \text{ D} \pm 2.69$ preoperative and postoperative it was $23.22 \text{ D} \pm 2.55$ using Binkhorst formula. The resulted mean difference was $3.20 \text{ D} \pm 0.86$ which was maximum among all the formulae in mild myopic group. The change was significant (p value-0.001). Similarly, the mean difference of diopters in SRK I, SRK II, SRK T, Hoffer Q and Holladay was $2.05 \text{ D} \pm 0.52$, $2.04 \text{ D} \pm 0.46$, $2.31 \text{ D} \pm 0.54$, $3.19 \text{ D} \pm 0.67$ and $2.72 \text{ D} \pm 0.60$ respectively, which proved to be significant. Minimum difference was seen in SRK II ($2.04 \text{ D} \pm 0.46$). In moderate myopic cases the resulted mean difference between the IOL power before and after surgery was $3.55 \text{ D} \pm 1.07$, $3.74 \text{ D} \pm 1.44$, $4.24 \text{ D} \pm 1.27$, $5.80 \text{ D} \pm 1.64$, $4.74 \text{ D} \pm 1.38$ and $5.52 \text{ D} \pm 1.60$ respectively using SRK I, SRK II, SRK T, Hoffer Q, Holladay and Binkhorst. The maximum difference was seen in Hoffer Q formula ($5.80 \text{ D} \pm 1.64$) while minimum was SRK I ($3.55 \text{ D} \pm 1.07$).

Similarly, in high myopic cases, the mean difference of IOL power after LASIK was $7.39 \text{ D} \pm 1.14$ using Hoffer Q. The mean difference of IOL power was seen maximum in Hoffer Q and minimum in SRK II ($4.56 \text{ D} \pm 0.73$) among the other formulae.

The mean change in IOL power among 150 eyes using SRK I was $2.95 \text{ D} \pm 1.20$, while in SRK II, SRK T, Hoffer Q, Holladay and Binkhorst was $3.03 \text{ D} \pm 1.41$, $3.45 \text{ D} \pm 1.45$, $4.73 \text{ D} \pm 1.91$, $3.91 \text{ D} \pm 1.55$ and $4.57 \text{ D} \pm 1.80$ respectively. Maximum change was seen in Hoffer Q and minimum change in SRK I formula.

There was significant difference in IOL power after LASIK surgery. The difference between the preoperative and postoperative IOL power was seen more in eyes with high diopters of myopia. We were not able to compare various different studies with our study as we were not able to find studies comparing various formulae before and after LASIK surgery. Our study was

first to study change in IOL power before and after LASIK surgery so more research is yet to be done on this topic.

Limitation of study:

There are considerable limitations of our study, among which one of the major limitation was none of our patient studied underwent actual cataract surgery. As the IOL power to be implanted can be better be judged after cataract surgery as post surgery residual errors gives surprising results.

LASIK surgery changes the anterior curvature of the cornea only; posterior curvature remains the same even after the surgery. We use corneal topograph to measure the keratometric values after LASIK surgery. Corneal topography measure the anterior curvature by taking normal anterior posterior curvature ratio (7.8mm/6.7mm). This ratio is altered after the LASIK surgery thus the keratometric values measured on the manual keratometry and topograph were overestimated than the actual true power of cornea. To find out the actual true power of cornea we require different formulae. This was the second most important area which was lacking in our study.

For better understanding the IOL power after LASIK surgery more research work has to be carried out with higher number of patients as a subject.

Conclusion:

Corneal refractive surgery significant change in keratometry values which was higher in high myopia cases, as the ablation required to correct the refractive error was more in high myopia, so as a result making standard intraocular lens power calculation formulas less accurate. The Keratometry values decreases after LASIK because ablation of cornea causes flattening of the cornea. So the IOL power after LASIK will be higher than the preoperative IOL power. So as a result patients who have undergone refractive surgery are at increased risk of postoperative refractive error following cataract surgery. The use of specialized post-refractive surgery IOL power calculation methods, modern formulas and advanced biometry can improve predictive accuracy and visual outcomes.

REFERENCE

1. Lou L, Yao C, Jin Y, Perez V, Ye J. Global Patterns in Health Burden of Uncorrected Refractive Error. *Investig Ophthalmology Vis Sci.* 2016 Nov 17;57(14):6271.
2. Pascolini D, Mariotti SP. Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol.* 1;96(5):614-8.(2012 May).

RESEARCH PAPER

3. Krishnaiah S, Srinivas M, Khanna RC, Rao GN. Prevalence and risk factors for refractive errors in the South Indian adult population: The Andhra Pradesh Eye disease study. *Clinical ophthalmology* (Auckland, NZ). 2009;3:17.
4. Sheeladevi S, Seelam B, Nukella P, Borah R, Ali R, Keay L. Prevalence of refractive errors, uncorrected refractive error, and presbyopia in adults in India: A systematic review. *Indian J Ophthalmol*. 2019;67(5):583.
5. Shortt AJ, Allan BD, Evans JR. Laser-assisted in-situ keratomileusis (LASIK) versus photorefractive keratectomy (PRK) for myopia. *Cochrane Database of systematic reviews*. 2013(1).
6. Seitz B, Langenbacher A. Intraocular lens power calculation in eyes after corneal refractive surgery. *Journal of refractive surgery*. 2000 May 1;16(3):349-61.
7. comprehensive report on the global refractive surgery market;http://dev.marketscope.com/market_reports/2010/9/2010-comprehensive-report-on-t-7.html.
8. Juhasz T, Loesel FH, Kurtz RM, Horvath C, Bille JF, Mourou G. Corneal refractive surgery with femtosecond lasers. *IEEE Journal of Selected Topics in Quantum Electronics*. 1999 Jul;5(4):902-10.
9. Aristodemou P, Cartwright NE, Sparrow JM, Johnston RL. Formula choice: Hoffer Q, Holladay 1, or SRK/T and refractive outcomes in 8108 eyes after cataract surgery with biometry by partial coherence interferometry. *Journal of Cataract & Refractive Surgery*. 2011 Jan 1;37(1):63-71.
10. Lee AC, Qazi MA, Pepose JS. Biometry and intraocular lens power calculation. *Current opinion in ophthalmology*. 2008 Jan 1;19(1):13-7.
11. Sharma N, Singhvi A, Sinha R, Vajpayee RB. Reasons for not performing LASIK in refractive surgery candidates.
12. Kato N, Toda I, Hori-Komai Y, Sakai C, Tsubota K. Five-year outcome of LASIK for myopia. *Ophthalmology*. 2008 May 1;115(5):839-44.
13. Yuen LH, Chan WK, Koh J, Mehta JS, Tan DT, SingLasik Research Group. A 10-year prospective audit of LASIK outcomes for myopia in 37 932 eyes at a single institution in Asia. *Ophthalmology*. 2010 Jun 1;117(6):1236-44.
14. Maldonado-Bas A, Onnis R. Results of laser in situ keratomileusis in different degrees of myopia. *Ophthalmology*. 1998 Apr 1;105(4):606-11.
15. Magallanes R, Shah S, Zadok D, Chayet AS, Assil KK, Montes M, Robledo N. Stability after laser in situ keratomileusis in moderately and extremely myopic eyes. *Journal of Cataract & Refractive Surgery*. 2001 Jul 1;27(7):1007-12.
16. Kymionis GD, Tsiklis NS, Astyrakakis N, Pallikaris AI, Panagopoulou SI, Pallikaris IG. Eleven-year follow-up of laser in situ keratomileusis. *Journal of Cataract & Refractive Surgery*. 2007 Feb 1;33(2):191-6.
17. Rosa N, Capasso L, Lanza M, Romano A. Axial eye length evaluation before and after myopic photorefractive keratectomy. *Journal of Refractive Surgery*. 2005;21(3):281-7.
18. Winkler Von Mohrenfels C, Gabler B, Lohmann CP. Optical Biometry before and after Excimer Laser Epithelial Keratomileusis (LASEK) for Myopia. *Eur J Ophthalmol*. 2003 Apr;13(3):257-9.