

Phytochemical Profiling and *In-Vitro* Antioxidant Activity *Curcuma longa* Extract Fractions

Priyanka U. Telang¹, Surendra G. Gattani^{2*}

1. Ms. Priyanka U. Telang Ph.D. Research scholar, School of Pharmacy, SRTMU, Nanded.
priyankatelang99@gmail.com

2. Dr. Surendra Gattani- Senior Professor, School of Pharmacy, SRTMU, Nanded.
sggattani@gmail.com

*Corresponding author: Surendra G. Gattani, Senior Professor, School of Pharmacy, S. R. T. M. University, Nanded- 43606, Maharashtra, India Tel.: +919970819627, Email: sggattani@gmail.com

ABSTRACT

Background:

Metabolic syndrome (MetS) is a complex illness characterized by obesity, insulin resistance, dyslipidemia, and hypertension, which elevates the risk of cardiovascular disease and type 2 diabetes mellitus. Natural products like *Curcuma longa* (Curcumin) are recognized for their multifaceted therapeutic potential, encompassing antioxidant, anti-inflammatory, antidiabetic, and cardioprotective activities. Although traditionally utilized, comprehensive phytochemical standardization and spectrum characterization are crucial for ensuring purity, repeatability, and medicinal efficacy.

Methods:

Rhizome of *Curcuma longa* were gathered, shade-dried, pulverized, and underwent sequential solvent extraction. Extracts were standardized by assessing foreign organic matter, moisture content, ash, and extractive values. Initial phytochemical analysis revealed bioactive compounds. Thin-layer chromatography (TLC) and column chromatography were utilized to isolate active fractions, which were further characterized by high-performance thin-layer chromatography (HPTLC). Structural elucidation was conducted utilizing UV-Vis, FTIR, ¹H NMR, and mass spectrometry. *In vitro* antioxidant activity of *curcuma longa* rhizome ethanolic extract using DPPH and nitric oxide scavenging assays

Major Findings:

Turmeric extracts were yellowish semisolids with a yield of 9.85% ethanol. Phytochemical investigation verified the existence of flavonoids, terpenoids, alkaloids, steroids, and tannins. TLC and HPTLC demonstrated several unique spots and R_f patterns. Spectroscopic examination revealed Curcumin analogue from *Curcuma longa*. These findings demonstrate the potent antioxidant potential of *Curcuma longa* rhizome extract and support its potential as a natural source of bioactive antioxidant compounds.

Conclusion:

The research standardized and described extracts of *Curcuma longa*, emphasizing their abundant phytochemical compositions and antioxidant potential, thereby endorsing their application in the management of metabolic illnesses.

Keywords: *Curcuma longa*, antioxidant, phytochemical characterization, curcumin.

How to cite this article: Priyanka U. Telang¹ Surendra G. Gattani^{2*}. Phytochemical Profiling and In-Vitro Antioxidant Activity *Curcuma longa* Extract Fractions. Int J Drug Deliv Technol. 2026;16(85s):1458-1468. DOI: 10.25258/ijddt.16.85s.179.

INTRODUCTION

Metabolic syndrome (MetS) is a multifaceted disorder defined by a combination of metabolic irregularities, including obesity, insulin resistance, dyslipidemia, and hypertension, which combined elevate the risk of cardiovascular disease and type 2 diabetes mellitus. The increasing global incidence of MetS has emerged as a significant public health issue, requiring the formulation of effective preventative and therapeutic measures.

Conventional pharmaceutical therapies exist, although they frequently address isolated components of MetS and may have long-term deleterious consequences. This has resulted in heightened interest in natural products and plant-based therapies owing to their multi-targeted mechanisms, enhanced safety profiles, and supplementary therapeutic advantages.

Curcumin (CUR), a bioactive polyphenol derived from the rhizome of *Curcuma longa*, has garnered considerable interest owing to its broad medicinal potential. Curcumin has been documented to provide protective effects against various pathological situations, including neurological illnesses, infections, cancer, and metabolic disorders [11–13]. *Curcuma longa*, generally referred to as turmeric, is extensively utilized as a culinary spice, natural dye, and dietary supplement. Curcumin possesses antioxidant, anti-inflammatory, antidiabetic, and anticancer attributes, rendering it a significant contender for the prevention and control of chronic illnesses [14,15]. Its enduring application in traditional medicinal systems, including Ayurveda, Thai medicine, and ancient Chinese medicine, further substantiates its therapeutic significance [16,17]. Despite the significant medicinal potential of *Curcuma longa*, comprehensive phytochemical standardization and spectrum characterization of their extract fractions are crucial for ensuring repeatability, quality control, and therapeutic efficacy. The current study seeks to conduct phytochemical and spectral characterisation of standardized extracts of *Curcuma longa* fractions, establishing a solid foundation for their prospective use in managing metabolic diseases.

2. Materials and Methods

2.1 Materials

All chemicals, including chloroform, hydrochloric acid, ethanol, petroleum ether, sulfuric acid, 5% ferric chloride, toluene, ethyl acetate, acetic acid, and formic acid, were procured from Merck and Cosmo Chem Pvt. Ltd. All solvents are of analytical grade.

2.2 Methods

2.2.1. Collections and Drying

Curcuma longa (rhizome) was harvested from the Latur district of Maharashtra, India, and subsequently shade-dried under controlled settings to prevent exposure to direct sunlight. The desiccated plant matter was subsequently pulverized into a coarse powder. Fine particles were eliminated by running the powdered substance through a 120-mesh sieve.

2.3. Standardization of plants materials

2.3.1. Determination of foreign organic matter

A thin layer of 5 g of coarsely powdered, air-dried drug material was distributed on a clean surface. The specimen was inspected visually and using a 6× magnifying lens. Foreign organic materials were meticulously isolated to the fullest extent practicable. The quantity of foreign organic matter was ascertained by weighing the isolated material and expressing it as a proportion of the initial sample weight [18].

2.3.2. Determination of moisture content

An accurately weighed, clean, dry weighing bottle equipped with a glass stopper was utilized.

Approximately 2 g of the sample was introduced into the bottle, uniformly distributed to a maximum depth of 10 mm, after which the bottle was sealed and weighed. The filled bottle was thereafter positioned in an oven and dehydrated until a stable weight was achieved. Upon drying, the bottle was extracted from the oven and permitted to cool to ambient temperature within a desiccator. The loss on drying was determined and represented as a percentage weight loss (% w/w) [19].

2.3.3. Ash value

The residual ash from the burning of the medicinal plant material was assessed utilizing three distinct methodologies: total ash, acid-insoluble ash, and water-soluble ash, in accordance with established protocols [20].

2.3.4. Determination of Total ash

Approximately 2 g of the air-dried crude medication was precisely measured and positioned in a pre-weighed silica dish. The material was burned at a temperature not above 450 °C until devoid of carbon. Upon permitting the dish to attain room temperature, the mass of the ash was documented. The total ash content was determined based on the air-dried substance and expressed as % w/w [21].

2.3.5. Determination of Water- soluble ash

The ash produced by the aforementioned process was subjected to boiling with 25 mL of water for 5 minutes, thereafter filtered, and the insoluble residue was gathered on ashless filter paper. The residue was rinsed with hot water and incinerated for 15 minutes at a temperature not surpassing 450 °C. Upon cooling, the mass of the insoluble residue was documented. The water-soluble ash was determined by subtracting the weight of the insoluble residue from the total ash weight and was represented as % w/w relative to the air-dried medication.

2.3.6. Determination of Acid -insoluble ash

The ash was subjected to boiling with 25 mL of 2 M hydrochloric acid for 5 minutes, thereafter filtered, and the insoluble residue was retained on ashless filter paper. The residue was rinsed with hot water, incinerated at a temperature not beyond 450 °C, cooled in a desiccator, and subsequently weighed. The acid-insoluble ash concentration was determined and represented as a percentage by weight (% w/w) relative to the air-dried medication.

2.3.7. Extractive values

Multiple extractive values, encompassing water-soluble and alcohol-soluble extractive values, were assessed utilizing established pharmacopoeial methodologies.

A) Determination of water-soluble extractive value

One hundred milliliters of chloroform water was incorporated into 1.5 g of air-dried, coarsely powdered medication and macerated in a sealed

container for 24 hours. The mixture was agitated intermittently for the initial 6 hours and subsequently permitted to rest for the ensuing 18 hours. Following filtration, 25 mL of the filtrate was placed in a shallow, flat evaporating dish, evaporated to dryness, and subsequently dried at 105 °C until a consistent weight was achieved. The water-soluble extractive value was determined and represented as % w/w relative to the air-dried substance [24].

B) Determination of Alcohol-soluble extractive value

In a sealed flask, 1.5 g of air-dried, finely powdered medication was macerated with 100 mL of ethanol of appropriate concentration for 24 hours. The mixture was agitated intermittently for the initial 6 hours and subsequently permitted to rest for the ensuing 18 hours. Following meticulous filtration to avert ethanol loss, 25 mL of the filtrate was placed in a shallow, flat evaporating dish, evaporated to dryness, and subsequently dried at 105 °C until a consistent weight was achieved. The ethanol-soluble extractive value was determined and represented as % w/w relative to the air-dried substance [25].

2.4. Extraction

Sequential extractions were conducted utilizing solvents of ascending polarity: petroleum ether (60–80 °C), ethanol, and water. The procedure was conducted in several batches utilizing a Soxhlet extractor to guarantee thorough extraction for each solvent fraction. At intervals, aliquots from the siphon tube were applied to thin-layer chromatography (TLC) plates and subjected to iodine vapor exposure. The lack of colorful dots signified the completion of extraction with the corresponding solvent. The solvent from each extract was then eliminated using distillation, utilizing lower pressure as required, and the concentrated residue was permitted to air dry. The desiccated extract was subsequently stored in hermetically sealed containers to maintain its integrity. The identical approach was adhered to for each subsequent solvent. The residual plant material (marc) was meticulously oven-dried to eliminate any remaining solvent before to advancing to the subsequent extraction phase. The desiccated plant material underwent reflux with distilled water for around three hours to provide the aqueous extract.

2.4.1. Phytochemical Test

The phytochemical tests are given in table 1

Table 1: Phytochemical Test

Sr	Phytochemical Test	Procedure	Observation
1			
2			

1	Test for Flavonoids	Dispense 1–2 mL of the plant extract into a sterile test tube. Carefully add several drops of pure sulfuric acid (H ₂ SO ₄). Monitor any alteration in hue, which signifies the existence of particular phytochemical elements.	Yellow/orange color
2	Test for Terpenoids	Dispense 1–2 mL of the plant extract into a sterile test tube. Incorporate 2 mL of chloroform into the extract. Gradually introduce 1–2 mL of concentrated sulfuric acid (H ₂ SO ₄) down the interior wall of the test	reddish-brown layer

		tube to create a distinct layer. Examine the contact between the layers for any chromatic alteration, which signifies the presence of particular phytochemical ingredients.	
3	Test for alkaloids	Dispense 1–2 mL of the plant extract into a sterile test tube. Introduce 1 mL of Dragendorff's reagent to the extract. Note the emergence of a precipitate, signifying the presence of alkaloids.	orange/reddish-brown precipitate

4	Test for steroids	Dispense 1–2 mL of the plant extract into a sterile test tube. Introduce 2 mL of chloroform to the extract. Gradually introduce 2 mL of concentrated sulfuric acid (H_2SO_4) along the interior wall of the test tube to create a distinct layer. Examine the interface for the emergence of a colorful ring, signifying the presence of particular phytochemical elements.	reddish-brown ring
5	Test for tannins	Dispense 1–2 mL of the plant extract into a sterile	Blue black/greenish color

		test tube. Introduce several drops of a 5% ferric chloride (FeCl_3) solution. Monitor any alteration in hue, which signifies the existence of phenolic compounds or tannins.	
--	--	--	--

2.4.2. Characterization of Bioactive Fractions by TLC

Subsequent to the pharmacological assessment of the ethanol fractions, the active fractions underwent thin-layer chromatography (TLC) for the identification of phytocomponents. The TLC examination of *Curcuma longa* extract, utilizing chloroform: methanol (9.8:2), revealed several spots, signifying the presence of various components.

Table 2: TLC studies of plant extracts

Sr. No	Extracts	Solvent system (v:v:v)	No. of spots
01	<i>Curcuma longa</i>	chloroform and methanol (9.8:2)	03

2.4.3. Column Chromatography of Active Extracts

Initial phytochemical analysis of *Curcuma longa* extracts revealed the existence of several phytochemicals in both ethanol and chloroform fractions. Column chromatography was conducted on these active extracts to isolate specific phytochemicals for subsequent study.

2.4.4. High Performance Thin Layer Chromatography (HPTLC) Fingerprint Analysis

1. Preparation of Extract

Each plant extract was dissolved in its respective solvent at a concentration of 5 mg/mL.

2. Application of Extract

The obtained extracts (5 mg/mL) were applied to HPTLC plates (10 × 10 cm) with a Linomat syringe. The samples were applied in bands measuring 5–6 mm in width, with a separation of 6 mm between them.

3. Development of the Chromatogram

The HPTLC plates were generated in a CAMAG twin-trough chamber with an optimum solvent mixture, as determined for TLC analysis. Upon reaching a development of 80 mm, the plates were extracted from the chamber and permitted to air-dry.

4. Scanning of the Chromatogram

The prepared plates were analyzed with a CAMAG HPTLC Scanner in absorbance mode at wavelengths of 254, 366, and 560 nm. The scanned data were examined utilizing the WinCATS planar chromatography manager software. Fingerprint chromatograms were produced to document the R_f values, band colors, and other distinctive characteristics of the chemicals found in the extracts. Bands at 254, 366, and 560 nm were seen without any supplementary derivatizing agents [26]

2.5. Structural Elucidation of Isolated Fractions

The fractions associated with TLC spots were subsequently analyzed using spectroscopic methods, including UV–Vis, mass spectrometry (MS), Fourier-transform infrared spectroscopy (FTIR), and proton nuclear magnetic resonance (^1H NMR), to ascertain their structural characteristics.

2.6. In vitro antioxidant activity:

2.6.1. DPPH method:

Principle: The 1,1-diphenyl-2-picrylhydrazyl (DPPH) is a stable free radical and is commonly used to evaluate the radical scavenging activity of antioxidant agents. The principle of this method is based on the fact that, decrement of DPPH in methanol solution in the presence of a hydrogen donating antioxidant due to the formation of the non-radical form DPPH.

Procedure:

- The free radical scavenging activity of test sample was determined by DPPH scavenging method.
- 0.1mM DPPH solution was prepared in methanol by adding 39.4 mg of DPPH in 1000 ml of methanol, and to 0.5 mL of this solution, 1.5 mL of test sample

dissolved in DMSO was added at various concentrations (1000, 500, 250, 125, 62.5, 15.62, 1.95 µg/ml).

- The mixtures were shaken vigorously and allowed to stand at room temperature for 30 minutes.
- Then the absorbance was measured at 517 nm using non-coated 96 well plate on microplate reader (EPOCH, Agilent BioTek, US). Ascorbic acid was used as standard compound.
- Reduction in absorbance by test compounds indicates radical scavenging activity.
- The scavenging activity by the DPPH radical was determined by

$$\text{DPPH scavenging effect (\% inhibition)} = \{(A_0 - A_1)/A_0\} \times 100\}$$

Where, A₀ is the absorbance of the control reaction, and A₁ is the absorbance test compound or Ascorbic acid

2.6.2: Nitric Oxide method:

Principle: Nitric oxide (NO) is a free radical that produced by interaction of NO with oxygen or reactive oxygen species. It is free radical due to its unpaired electron and exhibits similar properties like superoxide free radicals. This assay is based on the reduction of nitrate to nitrite by a reducing agent at 37 °C. Converted nitrite and endogenous nitrite are collectively converted by Griess reagent to a blue colored azo compound. This compound can be measured spectroscopically between 580-630 nm and absorbance is directly proportional to the total nitric oxide concentration.

Procedure:

- Nitric oxide (NO) radical scavenging activity of test extracts (PG & HI), were done, various concentrations of test compounds test extracts (PG & HI), such as 1000, 500, 250, 125, 62.5, 15.62, 1.95 µg/ml were dissolved in dimethyl sulphoxide (DMSO).
- To which, 0.5 mL of 10 mM sodium nitroprusside in phosphate buffered saline was added, then, 1 ml of various concentrations 1000, 500, 250, 125, 62.5, 15.62, 1.95 of test compounds were mixed, and to this equal volume of freshly prepared Griess reagent was added, solution was then incubated at 25°C for 3 hours.
- From above solution, 100 µl of the reaction mixture was transferred to a noncoated 96-well plate, and the absorbance was read at 546 nm using a microplate reader (EPOCH, Agilent BioTek, US). Ascorbic acid was used as standard control.

- The percentage of nitrite radical scavenging activity of test compounds was calculated by

$$\text{Nitric oxide scavenging activity} = \frac{\text{Absorbance of control} - \text{Absorbance of test compounds}}{\text{Absorbance of control}} \times 100$$

Absorbance of control

3. RESULTS AND DISCUSSION

3.1. Collections and Drying

The specimens of *Curcuma longa* (rhizome) were successfully gathered from the Latur region of Maharashtra, India. Shade drying in controlled settings produced dried plant material with negligible browning and no evidence of microbial contamination. The desiccated plants were further pulverized into coarse powder, and finer particles were eliminated by sifting through a 120-mesh screen. The resultant powdered substance exhibited a consistent texture, making it appropriate for subsequent extraction and phytochemical investigations.

3.2. Authentication

The identities of the collected plants were verified by comparing their morphological characteristics. Authentication was confirmed by the Botanical Survey of India, Pune, Maharashtra state, India (BSI/WRC/Iden.Cer./2021).

3.3. Standardization of plants materials

3.3.1. Determination of foreign organic matter

The examination of foreign organic matter indicated *Curcuma longa* rhizome exhibited a reduced contamination level of 0.1% w/w. Plant materials comply with the acceptable purity standards, since the allowable limit for foreign organic materials does not exceed 2% w/w.

3.3.2. Determination of moisture content

The moisture content of the extracts was assessed, yielding findings within acceptable parameters. The loss on drying (LOD) analysis indicated that *Curcuma longa* (rhizome) commenced at 0.00% w/w, escalated to 0.20% at 1 hour, and attained 0.22% at 2 hours, sustaining this concentration through 3 and 4 hours. The results demonstrate negligible and consistent moisture loss over time for extracts, affirming that all plant extracts remain below the permitted limits of detection.

3.3.3. Determination of Ash value

The ash content of the plant materials was analysed to determine their inorganic residue profiles. The *Curcuma longa* (Rhizome) exhibited a marginally elevated total ash level of 3.0% w/w, comprising acid-insoluble ash at 1.6% w/w and water-soluble ash at 0.2% w/w. The results reveal significant disparities in the inorganic makeup of the plant materials.

3.3.4. Determination of Extractive values

The extractive qualities of the plant materials were assessed to determine the solubility of their contents in various solvents. The *Curcuma longa*

(rhizome) exhibited marginally elevated values, with an ethanol-soluble extractive value of 12.5% w/w and a water-soluble extractive value of 8.1% w/w, signifying superior solubility of its components in both solvents.

3.4. Extraction

Extraction was conducted in several batches utilizing different solvents. The characteristics and yield percentages of each extract are summarized below:

3.4.1. Yield of Various Extracts from rhizome powder (*Curcuma longa*)

All solvent fractions of *Curcuma longa* yielded yellowish, semisolid extracts. Ethanol had the maximum extraction yield at 9.85%, followed by chloroform at 7.45%, ethyl acetate at 6.52%, petroleum ether at 6.22%, and the aqueous extract, which exhibited the lowest yield at 6.07%.

3.5. Preliminary Phytochemical Screening

The initial phytochemical study of the extracts revealed the existence of many bioactive compounds, such as triterpenoids, steroids, glycosides, saponins, alkaloids, flavonoids, tannins, proteins, free amino acids, carbohydrates, and vitamins. The data indicate that both plant extracts are abundant in various phytometabolites with potential pharmacological properties.

3.5.1. Phytochemical test *Curcuma longa* Extract

The initial phytochemical analysis of the *Curcuma longa* (rhizome) extract verified the existence of terpenoids, although flavonoids, alkaloids, steroids, and tannins were not detected. The affirmative terpenoid reaction indicates the existence of curcumin, a polyphenolic molecule generated from terpenoids that accounts for the primary pharmacological effects of the plant. The results suggest that the biological effects of the extract are mostly due to terpenoid components.

Table 4: Phytochemical test of *Curcuma longa* Extract

Sr. No.	Phytochemical test	Test Extracts
1.	Test for Flavonoids	
	H ₂ SO ₄ Test	-
2.	Test for Terpenoids	
	Choloroform test	+
3	Test for alkaloids	
	Dragondroff's reagent	-
4	Test for steroids	
	Salkowski test	-
5	Test for tannins	
	FeCl ₃ test	-

The phytochemical study of *Curcuma longa* identified the presence of flavonoids, terpenoids, alkaloids, steroids, and tannins.

3.6. Thin layer chromatography of extract

The thin layer chromatography of the plant extracts revealed well-defined and distinct spots corresponding to alkaloids, flavonoids, tannins, and steroids under optimum mobile phases. The measured R_f values and distinct colour reactions post-spraying validated the existence and varied distribution of these phytoconstituents across different solvent extracts. The disparity in R_f values among solvents indicates variations in polarity and chemical makeup of the extracts. The TLC results affirm the repeatability and reliability of phytochemical profiling of the extracts.

Table 5: Thin Layer Chromatography of all extracts.

Sr. No.	Chemical constituent	Mobile Phase	Visualization Sprayin g reagent	Color of spot	Rf-value
1.	Alkaloids	Ethyl acetate: Formic acid: Toluene (40:10:50)	10% H ₂ SO ₄ in ethanol	Violet - blue	Petroleum ether :0.71 Ethanol: 0.87 Chloroform : 0.86
2.	Flavonoid	Ethyl acetate: Toluene : Water: Glacial acetic acid (100:11:26:11)	Anisaldehyde – Sulfuric acid.	Yellowish green	Petroleum ether: 0.73 Ethanol: 0.81 Chloroform : 0.89
3.	Tannin	Water: acetic acid: ethyl	5 % FeCl ₃ in 0.1N HCl	Black	Ethanol : 0.62

		acetate: formic acid (26:11:1 00:11)			
4.	Steroi ds	Methan ol: Acetic acid: Ethyl acetate (20:10:7 0)	Vanilli n – Sulfuric acid.	Pink	Petrole um ether: 0.83 Ethano l: 0.75 Chloro form: 0.46 Ethyl acetate : 0.82

Figure 1: TLC chromatography for Alkaloid, flavonoid, tannins and steroids

3.7. Column Chromatography of Active Extract of both Plant Extract

Preliminary phytochemical tests conducted on the selected plant materials using ethanol and chloroform extracts have shown the presence of alkaloids, steroids, flavonoids, and terpenoids. The TLC of the extract also indicates the presence of several phytometabolites. The ethanol and chloroform extract demonstrates significant potential phytoconstituents. Isolation of phytocomponents from ethanol was necessary.

3.8. Percent yield and Appearance of all fractions.

The fractionation of plant extracts was carried out using different mobile phases. For Moringa extract, ethyl acetate and methanol in a 5:5 ratios were used, resulting in three fractions: MEF1 (0.78 g, 2.9% w/w), MEF2 (0.69 g, 2.8% w/w), and MEF3 (0.72 g, 3.2% w/w). For Curcumin extract (95% powder), methanol was used as the mobile phase, yielding three fractions: CEF1 (0.94 g, 5.4% w/w), CEF2 (0.89 g, 4.2% w/w), and CEF3 (0.82 g, 4.9% w/w).

A) Ethanol Extract Column Chromatography

Table 6: Appearance and percent yield of all fractions.

Plant Name	Mobile phase	Fraction Designation	Weight of fraction (gm.)	% Yield w/w
<i>Curcuma longa</i> extract	Methanol	CEF1	0.94	5.4
		CEF2	0.89	4.2
		CEF3	0.82	4.9

Figure 2: TLC chromatograph of *Curcuma longa* extract


Rf values of plant extracts

The Rf values *curcuma longa* extract are 0.22, 0.32, and 0.43. The Thin Layer Chromatography (TLC) examination of diverse plant extracts was conducted at different detection wavelengths, demonstrating unique Rf values for each extract. *Curcuma longa* (CEF3), identified at 366 nm, exhibited 8 spots with Rf values of 0.07, 0.26, 0.32, 0.43, 0.50, 0.62, 0.81, and 0.84. Each plant extract displayed distinct Rf patterns, underscoring the variability in their chemical contents.

3.9. High Performance Thin Layer Chromatography (HPTLC)

3.9.1 HPTLC fingerprints study

The HPTLC fingerprint analysis at 366 nm exhibited clearly defined and consistent



chromatographic patterns for both samples. *Curcuma longa* (CEF3) exhibited nine distinct bands with Rf values ranging from 0.07 to 0.84, indicating a chemically diverse profile aligned with curcuminoids and associated compounds. The unique HPTLC fingerprints validate the complexity, identity, and consistency of both extracts, affirming their appropriateness for quality control and additional pharmacological research.




Figure 3: [a] HPTLC chromatogram of MEF1 extract measured at 366nm [b] HPTLC chromatogram of CEF3 extract measured at 366nm

3.10. Structural Elucidation of Compounds isolated by Column Chromatography




Figure:5 [a] UV graph of Compound – I [b] IR spectra of compound-I [c] Proton NMR spectrum of Compound –I [d] MASS spectrum of Compound –I

The chromatographic behavior and the distinctive IR, ^1H NMR, and mass spectroscopic characteristics of the isolated molecule exhibited significant concordance with the documented spectral data of curcumin in the literature. The identified functional groups, proton signals, and molecular ion peak corresponded with those of conventional curcumin. Consequently, the isolated chemical can be unequivocally identified as a Curcumin analogue

3.10. In vitro antioxidant activity by DPPH and NO Method

Table 7: Observation table for Ascorbic acid

Sr. No	Conc.($\mu\text{g/ml}$)	DPPH	λ_{max}	Ascorbic acid
1	5	0.5	514	69.50 $\pm$ 0.00
2	10	0.5	514	71.88 $\pm$ 0.01
3	15	0.5	514	75.83 $\pm$ 0.01
4	20	0.5	514	81.71 $\pm$ 0.01
5	25	0.5	514	86.64 $\pm$ 0.01





Table 08: In-vitro Antioxidant Study of ethanolic extracts of Curcuma longa by DPPH and NO Method:

Sr. No	Conc. $\mu\text{g/ml}$	Abs. at 596	Scavenging activity (%) NO	Abs. at 596	Scavenging activity (%) DPPH
1	1.95	0.087	10.31	0.091	1.086
2	15.62	0.083	14.43	0.085	07.608
3	62.5	0.076	21.65	0.074	19.565
4	125	0.072	25.77	0.065	29.347
5	250	0.056	42.27	0.053	42.391
6	500	0.051	47.42	0.041	55.434
7	1000	0.005	48.45	0.018	80.366



4. CONCLUSION

This study effectively standardized and described extracts of *Curcuma longa*, validating their abundant phytochemical makeup, which includes flavonoids, tannins, phenolics, and terpenoids. TLC and HPTLC investigations exhibited the repeatability and diversity of phytoconstituents via unique Rf values and fingerprint patterns. Column chromatography facilitated the extraction of bioactive fractions, CEF3 from *Curcuma longa*, which were further structurally characterized using UV-Vis, FTIR, ¹H NMR, and mass spectrometry. Spectroscopic results, corroborated by literature, verified that these substances are counterparts of curcumin. The study offers an extensive phytochemical and spectral analysis of these plant extracts, establishing a solid foundation for their potential therapeutic use in controlling metabolic disorders and facilitating additional pharmacological and clinical research. The ethanolic extract of *Curcuma longa* rhizomes demonstrated significant *in vitro* antioxidant activity in DPPH and nitric oxide radical-scavenging assays. These findings indicate its promising free-radical-scavenging potential and suggest that *Curcuma longa* may serve as a natural source of antioxidant compounds with potential relevance to the management of oxidative stress-associated metabolic disorders. Further *in vivo* and clinical investigations are warranted to establish its therapeutic potential.

Conflict of Interest

The authors declare no Conflict of Interest.

References

1. K. Pal, S. Chowdhury, S. K. Dutta, et al., "Analysis of Rhizome Colour Content, Bioactive Compound Profiling and Ex-Situ Conservation of Turmeric Genotypes (*Curcuma longa* L.) From Sub-Himalayan Terai Region of India," *Industrial Crops and Products* 150 (2020): 112401, <https://doi.org/10.1016/j.indcrop.2020.112401>.
2. A. Rauf, M. Imran, I. E. Orhan, and S. Bawazeer, "Health Perspectives of a Bioactive Compound Curcumin: A Review," *Trends in Food Science and Technology* 74 (2018): 33–45.
3. J. Sharifi-Rad, Y. E. Rayess, A. A. Rizk, et al., "Turmeric and Its Major Compound Curcumin on Health: Bioactive Effects and Safety Profiles for Food, Pharmaceutical, Biotechnological and Medicinal Applications," *Frontiers in Pharmacology* 11 (2020): 1021, <https://doi.org/10.3389/fphar.2020.01021>.
4. S. Rathore, M. Mukim, P. Sharma, S. Devi, J. C. Nagar, and M. Khalid, "Curcumin: A Review for Health Benefits," *International Journal of Research and Review* 7, no. 1 (2020): 273–290.
5. Alhar MS, El-Sofany WI, AlRashidi AA, Hamden K. Protective Effects of Isolated Curcumin from *Curcuma longa* on Key Enzymes Involved in the Insulin Signaling Pathway and Digestive and Metabolic Enzymes Associated with Obesity, Type 2 Diabetes, and Hypertension. *Journal of Diabetes Research*. 2025;2025(1):8050374.
6. Sharifi-Rad, J.; Rodrigues, C.F.; Sharopov, F.; Docea, A.O.; Can Karaca, A.; Sharifi-Rad, M.; Calina, D. Diet, lifestyle and cardiovascular diseases: Linking pathophysiology to cardioprotective effects of natural bioactive compounds. *Int. J. Environ. Res. Public Health* 2020, 17, 2326.
7. Matter FO. *Methods of Analysis*. Retrieved from. 1984.
8. Kays AO. Determination of Moisture Content in Composites by Dielectric Measurements. 1979 Jul 1.
9. Harris GK, Marshall MR. Ash analysis. In *Food analysis* 2017 Jun 7 (pp. 287-297). Cham: Springer International Publishing.
10. Rohmah M, Saragih B, Amaliah N, Kristopal K, Putra YH, Rahmadi A. Determination of moisture, ash, protein, polyphenolic, flavonoids, and amino acid contents and antioxidant capacity of dried Mekai (*Pycnarrhena tumefacta* Miers) leaf as potential herbal flavor enhancers. In *International Conference on Tropical Agrifood, Feed and Fuel (ICTAFF 2021)* 2022 Jan 7 (pp. 149-158). Atlantis Press.
11. Querol X, Umaña JC, Alastuey A, Bertrana C, Lopez-Soler A, Plana F. Extraction of water-soluble impurities from fly ash. *Energy Sources*. 2000 Sep 1;22(8):733-49.
12. Sales J, Janssens GP. Acid-insoluble ash as a marker in digestibility studies: a review. *Journal of Animal and Feed Sciences*. 2003 Jul 15;12(3):383-401.
13. Sluiter A, Ruiz R, Scarlata C, Sluiter J, Templeton DJ. Determination of extractives in biomass. *Laboratory analytical procedure (LAP)*. 2005 Jul 17;1617(4):1-6.
14. Chen SF, Mowery RA, Scarlata CJ, Chambliss CK. Compositional analysis of water-soluble materials in corn stover. *Journal of agricultural*

- and food chemistry. 2007 Jul 25;55(15):5912-8.
15. Joshi VK, Joshi A, Dhiman KS. The Ayurvedic Pharmacopoeia of India, development and perspectives. Journal of ethnopharmacology. 2017 Feb 2; 197:32-8.
 16. Ramaley L, Vaughan MA, Jamieson WD. Characteristics of a thin-layer chromatogram scanner-mass spectrometer system. Analytical Chemistry. 1985 Jan 1;57(1):353-8.