

Manual Nerve Therapy in Children With Epilepsy, Cerebral Palsy and Autism: A Case Series of Observed Clinical Improvements

Dr. Hemant Jaisingh^{1*}

^{1*}Vinayak Clinic, Haridwar

Email: upsconlineacademy@gmail.com

Received: 25th Aug, 2026 | Revised: 1st Sep, 2026 | Accepted: 5th Sep, 2026 | Available Online: 12th Sep, 2026

Abstract

Background

Epilepsy, cerebral palsy, and autism spectrum disorder are heterogeneous pediatric neurological and neurodevelopmental conditions that may be associated with substantial functional limitations and, in some children, recurrent seizures and impaired motor development. Manual Nerve Therapy (MNT) is a non-invasive manual stimulation approach involving stimulation of selected anatomical regions and, in some cases, broader stimulation of body regions and joints. Evidence regarding the clinical effects of this approach is currently limited.

Case Presentation

We report six pediatric cases treated with Manual Nerve Therapy. Two children had recurrent epilepsy reportedly occurring approximately 20–30 times per day. Following approximately 40 MNT sessions, the caregivers reported cessation of clinically observed seizure episodes without concomitant antiseizure medication. In the first child, stimulation included the parietal and mastoid regions and buccal stimulation, with improvement reportedly observed within seven days. In the second child, similar stimulation was associated with reported improvement within one day.

A third child had cerebral palsy with epilepsy and inability to sit independently. Approximately 40 MNT sessions were administered, including stimulation of the parietal, mastoid, buccal, dorsal, and cervical regions. Independent sitting was reportedly achieved during the treatment period, with improvement reported within approximately 20–25 days.

Three additional children had neurodevelopmental disorders associated with inability to sit independently. One child had autism and two had cerebral palsy. Approximately 35–40 MNT sessions were administered. Dorsal and cervical regions were stimulated, together with broader MNT stimulation of body regions and joints. Independent sitting was reportedly observed after approximately 2 days, 6 days, and 9–10 days, respectively.

Conclusion

This six-case series describes temporal associations between MNT and reported improvements in seizure frequency and gross motor function in children with epilepsy, cerebral palsy, and autism. The observations are hypothesis-generating and cannot establish causality or treatment efficacy. Prospective studies using standardized neurological assessments, objective seizure documentation, defined treatment protocols, appropriate control groups, and long-term follow-up are required to determine whether MNT has reproducible therapeutic effects.

Keywords

How to cite this article: 1*Vinayak Clinic Haridwar. Manual Nerve Therapy in Children With Epilepsy, Cerebral Palsy and Autism: A Case Series of Observed Clinical Improvements. *Int J Drug Deliv Technol.* 2026;16(82s):1512-1515. DOI: 10.25258/ijddt.16.82s.172.

1. Introduction

Epilepsy is one of the major neurological disorders affecting children and is characterized by a predisposition to recurrent epileptic seizures.

Seizure control is an important objective of pediatric neurological care because frequent seizures may interfere with development, learning, sleep, behavior, and quality of life.

Cerebral palsy comprises a group of permanent disorders affecting movement and posture that result from disturbances in the developing fetal or infant brain. Children with cerebral palsy may have substantial limitations in gross motor development, including delayed acquisition of independent sitting.

Autism spectrum disorder is a neurodevelopmental condition characterized by persistent differences in social communication and interaction together with restricted or repetitive patterns of behavior or interests. Motor abnormalities and developmental delays may coexist in some children.

Manual Nerve Therapy (MNT) is described by the practitioner as a non-invasive manual stimulation approach involving stimulation of selected anatomical regions and, in some cases, systematic stimulation of multiple body regions and joints. The intervention is intended to provide manual sensory and musculoskeletal stimulation without invasive procedures.

Despite the clinical use of various complementary and manual approaches in different settings, evidence concerning MNT in pediatric neurological and neurodevelopmental disorders remains limited. The present case series therefore describes six children in whom improvements in seizure frequency and/or gross motor function were temporally observed during or following MNT.

The objective of this case series is not to establish efficacy but to document the clinical observations in a transparent manner and to generate hypotheses for future controlled investigation.

2. Case Series

2.1 Case 1 – Pediatric Epilepsy

An 8-year-old child (Case 1) was reported to have epilepsy with approximately 20–30 attacks per day before initiation of MNT.

The child received approximately 40 sessions of Manual Nerve Therapy. According to the treatment record provided by the practitioner, stimulation was directed toward the parietal region, mastoid region, and buccal region.

A reduction in clinically observed seizure episodes was reportedly noticed within approximately seven

days of initiation of the intervention. Following approximately 40 sessions, the practitioner and/or caregivers reported cessation of the episodes.

The supplied case information states that the child was not receiving antiseizure medication during the period in which seizure cessation was observed.

2.2 Case 2 – Pediatric Epilepsy

A 5-year-old child (Case 2) was reported to have epilepsy characterized by approximately 20–30 attacks per day.

Approximately 40 MNT sessions were administered. The intervention included manual stimulation of the parietal region, mastoid region, and buccal region.

According to the supplied clinical history, improvement in seizure episodes was observed within approximately one day. Following approximately 40 sessions, cessation of the reported seizure episodes was observed.

The supplied information states that the child was not receiving antiseizure medication during this period.

2.3 Case 3 – Cerebral Palsy With Epilepsy

A 4-year-old child (Case 3) was reported to have cerebral palsy and epilepsy and was unable to sit independently.

Approximately 40 MNT sessions were administered. Treatment included stimulation of the parietal, mastoid, and buccal regions. In addition, the dorsal and cervical regions were stimulated.

During the treatment period, the child reportedly developed the ability to sit independently. The practitioner reported that the response became apparent within approximately 20–25 days.

The case therefore involved two reported outcomes: improvement in seizure activity and improvement in gross motor function, specifically independent sitting.

2.4 Case 4 – Autism Spectrum Disorder With Delayed Sitting

A 5-year-old child (Case 4) was reported to have autism and inability to sit independently.

Approximately 35–40 MNT sessions were administered. Treatment included stimulation of the dorsal and cervical regions. In addition, the practitioner reported systematic stimulation of multiple nerves and joints as part of the MNT approach.

Independent sitting was reportedly achieved during the treatment period, with the response observed within approximately 9–10 days.

Because sitting ability represents a gross motor developmental milestone, objective assessment of motor function before and after treatment would be important to determine the magnitude and reproducibility of the reported change.

2.5 Case 5 – Cerebral Palsy With Delayed Sitting

A 4-year-old child (Case 5) was reported to have cerebral palsy and inability to sit independently.

Approximately 35–40 MNT sessions were administered. The intervention included stimulation of the dorsal and cervical regions, together with broader stimulation of multiple nerves and joints according to the MNT protocol used by the practitioner.

The child reportedly began sitting independently within approximately two days.

The rapidity of the reported functional change makes objective documentation particularly important. Baseline and follow-up standardized motor assessments would strengthen the scientific interpretation of this case.

2.6 Case 6 – Cerebral Palsy With Delayed Sitting

A 5-year-old child (Case 6) was reported to have cerebral palsy and inability to sit independently.

Approximately 35–40 MNT sessions were administered. The dorsal and cervical regions were stimulated, together with broader stimulation of multiple nerves and joints.

Independent sitting was reportedly achieved within approximately six days.

As with the other cases, objective documentation of baseline motor function, intervention exposure, concurrent rehabilitation, and subsequent follow-up is required before the observation can be interpreted as an effect attributable specifically to MNT.

3. Summary of the Six Cases

Case	Age	Reported condition	Baseline problem	Approx. MNT sessions	Main reported intervention	Reported response
1	8 y	Epilepsy	20–30 attacks/day	~40	Parietal, mastoid, buccal stimulation	Seizure cessation; response ~7 days
2	5 y	Epilepsy	20–30 attacks/day	~40	Parietal, mastoid, buccal stimulation	Seizure cessation; response ~1 day
3	4 y	Cerebral palsy + epilepsy	Unable to sit independently	~40	Parietal, mastoid, buccal + dorsal/cervical stimulation	Sitting independently; response ~20–25 days
4	5 y	Autism	Unable to sit independently	~35–40	Dorsal/cervical + broader MNT stimulation	Sitting independently; response ~9–10 days
5	4 y	Cerebral palsy	Unable to sit independently	~35–40	Dorsal/cervical + broader MNT stimulation	Sitting independently; response ~2 days
6	5 y	Cerebral palsy	Unable to sit independently	~35–40	Dorsal/cervical + broader MNT stimulation	Sitting independently; response ~6 days

4. Proposed Timeline of Treatment

The available information permits the following preliminary timeline:

Baseline assessment → initiation of MNT → repeated treatment sessions → first observed response → completion of approximately 35–40 sessions → post-treatment observation/follow-up

5. Intervention Description

5.1 Manual Nerve Therapy

Manual Nerve Therapy was administered manually without invasive procedures.

The treatment protocol used in the reported cases included stimulation of selected anatomical regions. Depending on the clinical presentation, these included:

- Parietal region
- Mastoid region
- Buccal region
- Cervical region
- Dorsal region
- Multiple peripheral anatomical regions and joints

The exact technique, pressure, duration, sequence, laterality, number of repetitions, interval between sessions, and practitioner qualifications should be documented in the final manuscript to permit reproducibility.

The treatment should be described as an **investigational/manual intervention** rather than as a proven therapy.

6. Safety and Adverse Events

No adverse events were reported in the information supplied for these six cases.

7. Discussion

This six-case series describes reported temporal associations between Manual Nerve Therapy and clinical improvements in children with epilepsy, cerebral palsy, and autism.

The most notable observations were the reported reduction or cessation of frequent seizure episodes in two children with epilepsy and the reported acquisition of independent sitting in four children with cerebral palsy and/or autism.

The epilepsy cases are particularly noteworthy because the baseline history provided by the practitioner indicates approximately 20–30 attacks per day, followed by reported cessation after a course of approximately 40 MNT sessions.

8. Strengths

Several features make these cases potentially useful for hypothesis generation.

First, the cases describe repeated clinical observations rather than a single isolated event.

Second, the reported epilepsy cases involved a high stated baseline frequency of attacks, allowing a potentially meaningful before-and-after comparison if the seizure frequency can be verified through contemporaneous records.

Third, the motor cases describe a clinically recognizable functional milestone—independent sitting.

Fourth, the cases provide information concerning the anatomical regions targeted during the intervention.

Finally, the observations may provide a basis for designing a prospective study with standardized outcome measures.

9. Clinical Implications

The observations reported here are hypothesis-generating rather than definitive evidence of efficacy.

If similar findings are reproduced prospectively, MNT could become an area for systematic investigation in pediatric neurological and neurodevelopmental disorders.

10. Conclusion

This six-case case series reports temporal associations between Manual Nerve Therapy and observed clinical improvements in children with epilepsy, cerebral palsy, and autism, including reported cessation of frequent seizure episodes and acquisition of independent sitting.

These observations are clinically interesting but do not establish that MNT caused the reported outcomes. Objective seizure documentation, standardized neurological and developmental assessments, detailed treatment protocols, control groups, and prospective follow-up are required to determine whether the observed improvements are reproducible and attributable to MNT.

The principal value of these cases is therefore their potential to generate a testable clinical research hypothesis regarding the effects of structured manual stimulation on neurological and motor outcomes in children.