

Community Health Recovery after COVID-19: Examining Pharmaceutical Accessibility, Medication Practices and Healthcare Behaviour in Dehradun

Jaanvi¹, Dr Roopak Kumar²

¹ Ph.D. Research Scholar, School of Arts and Humanities, Maya Devi University, Dehradun, Uttarakhand;

² Research Supervisor, School of Arts and Humanities, Maya Devi University, Dehradun, Uttarakhand;

Corresponding Author

Jaanvi

Email: janvishankar22@gmail.com

Abstract

The COVID-19 pandemic affected community healthcare not only through infection but also by disrupting routine treatment, medicine supplies and healthcare-seeking practices. Restrictions on movement, fear of infection, reduced household income and difficulties in reaching healthcare facilities affected the way people obtained medicines and responded to illness. Although healthcare services have largely returned to normal functioning, some practices developed during the pandemic may continue to influence healthcare behaviour.

This paper examines post-COVID community health recovery in Dehradun by focusing on three closely related aspects: pharmaceutical accessibility, medication practices and healthcare behaviour. Pharmaceutical accessibility is examined through medicine availability, affordability, geographical reach, continuity of supply and access to reliable information. Medication practices include adherence to prescriptions, self-medication, use of earlier prescriptions and changes in medicine use. Healthcare behaviour refers to decisions concerning when to seek treatment, where to seek it and whether professional consultation is preferred over informal sources.

The study follows a literature-based and conceptual approach. Existing evidence indicates that the pandemic created significant barriers to healthcare access, particularly for people requiring regular treatment. Financial difficulties, restrictions on movement, medicine-supply problems and fear of healthcare facilities encouraged some individuals to depend more heavily on pharmacies, previous prescriptions and remote consultations. At the same time, self-medication and inappropriate medicine use became important concerns.

The paper argues that recovery from a health emergency cannot be judged only by the reopening of hospitals or restoration of routine services. Sustainable recovery also requires affordable medicines, continuity of treatment, responsible medication practices and timely use of professional healthcare. Dehradun provides a useful setting for examining these issues because the city has a combination of public and private hospitals, clinics, pharmacies, diagnostic services and specialist facilities. The study provides an integrated perspective on the relationship between pharmaceutical accessibility, medication practices and healthcare behaviour in the post-COVID period.

Keywords: COVID-19 recovery, pharmaceutical accessibility, medication practices, healthcare behaviour, self-medication, medicine adherence, community health, Dehradun

How to cite this article: Jaanvi¹ Dr Roopak Kumar². Community Health Recovery after COVID-19: Examining Pharmaceutical Accessibility, Medication Practices and Healthcare Behaviour in Dehradun. Int J Drug Deliv Technol. 2026;16(83s):1450-1460. DOI: 10.25258/ijddt.16.83s.157.

1. Introduction

The COVID-19 pandemic placed extraordinary pressure on healthcare systems and changed the way communities used health services. During the most critical phases of the pandemic, hospitals and health workers were required to prioritize COVID-19 testing, isolation and treatment. At the same time, people continued to require medicines and medical attention for chronic diseases, infections

and other health problems. Restrictions on movement and changes in healthcare priorities made routine treatment more difficult for many households.

The disruption was not limited to hospitals. Pharmaceutical supply chains were affected by

transportation restrictions, changes in demand and difficulties faced by workers involved in medicine distribution. Patients who normally obtained medicines through regular consultations sometimes had difficulty meeting doctors or renewing prescriptions. For people with long-term conditions, even a temporary interruption could affect treatment continuity.

The World Health Organization has reported that essential health services were disrupted in many countries during the pandemic. Although service disruption gradually declined as health systems adapted, the experience demonstrated that healthcare recovery involves more than reopening facilities. It also requires the restoration of reliable access, continuity of treatment and public confidence in healthcare services.

The Indian experience reflected many of these difficulties. Patients with chronic illnesses faced particular challenges when movement restrictions and reduced healthcare availability affected regular consultations. Financial insecurity also became an important factor. Households experiencing income loss could find it difficult to meet expenditure on medicines, diagnostic services and private consultations. Thus, access to healthcare was influenced not only by whether a service existed but also by whether people could afford and reach it.

Pharmaceutical accessibility is therefore better understood as a combination of availability, affordability, geographical accessibility and continuity. A medicine may be available in a city but still remain inaccessible to a household that cannot afford it. Similarly, a nearby pharmacy does not guarantee appropriate treatment if the required medicine is unavailable or the individual lacks reliable information about its use.

The financial dimension is particularly relevant in India. National Health Accounts estimates for 2021–22 indicate that household out-of-pocket expenditure continued to represent a substantial share of total health expenditure. This makes the cost of medicines and consultations an important consideration in household healthcare decisions. Financial pressure may influence whether patients purchase a complete course of treatment, choose a less expensive alternative, postpone consultation or depend on self-treatment.

The pandemic also influenced medication practices. When access to doctors became difficult, some individuals continued medicines from earlier prescriptions or sought advice from pharmacists, family members and digital sources. In certain

circumstances, such practices helped people maintain treatment during a period of restricted access. However, using an old prescription for a new illness or changing dosage without professional advice can also create risks.

Self-medication became particularly visible during the pandemic. Individuals often attempted to manage symptoms without consultation, using medicines based on previous experience or advice obtained from informal sources. Although responsible use of certain over-the-counter medicines can be appropriate, unsupervised use of prescription medicines and antibiotics may result in adverse effects, inappropriate treatment and delayed diagnosis.

Healthcare-seeking behaviour also changed. Fear of infection discouraged some people from visiting hospitals, while transportation restrictions and financial uncertainty created additional barriers. Telemedicine and telephone consultations offered alternative ways of communicating with healthcare professionals. Pharmacies also became convenient first-contact points, especially for minor illnesses.

Some of these practices may have continued after restrictions were removed. People who became comfortable with remote consultations may continue using them for suitable conditions, while those who developed confidence in self-treatment may continue avoiding professional consultation. These changes make it necessary to examine community health recovery from a behavioural as well as an institutional perspective.

Dehradun offers an appropriate setting for such an examination. The city has a varied healthcare environment involving government and private hospitals, clinics, pharmacies, diagnostic centres and specialist services. It also serves people from surrounding areas who travel to the city for treatment. Consequently, access to healthcare can differ according to income, location, health requirements and individual preferences.

The present study brings pharmaceutical accessibility, medication practices and healthcare behaviour into a common framework. Rather than considering them as separate issues, it examines how access to medicines can influence medication decisions and how those decisions can affect subsequent healthcare-seeking behaviour.

2. Background and Review of Literature

The pandemic created a situation in which healthcare demand and healthcare access were disrupted at the same time. Hospitals faced pressure

from COVID-19 cases, while patients with other conditions experienced difficulty obtaining routine services. Transport restrictions, fear of infection and financial problems further affected healthcare utilization.

The consequences were particularly relevant for people who required continuous treatment. Individuals with diabetes, hypertension and other chronic conditions could not always maintain their normal pattern of consultations and medicine collection. Interruptions were sometimes caused by medicine shortages, while in other cases patients deliberately avoided healthcare facilities because of concerns about infection.

Pharmaceutical access has several components. **Availability** refers to whether the required medicine can be obtained. **Affordability** concerns whether the household can purchase it without serious financial difficulty. **Geographical access** relates to distance and transportation, while **continuity** refers to the ability to obtain medicines repeatedly throughout treatment. Finally, **information access** concerns the availability of trustworthy guidance regarding medicine selection and use.

Table 1. Dimensions of Pharmaceutical Accessibility

Dimension	Meaning	Common barrier
Availability	Required medicines can be obtained when needed	Stock shortages or limited supply
Affordability	Household can meet the cost of treatment	High medicine expenditure
Geographical access	Pharmacy or healthcare facility can be reached	Distance and transportation
Continuity	Medicines remain accessible throughout treatment	Interrupted supply or treatment
Information access	Reliable guidance is available	Misinformation or informal advice

Source: Compiled by the author based on WHO (2020), International Pharmaceutical Federation (2020), and Government of India, Ministry of Health and Family Welfare (2024).

These dimensions are closely related. A patient may live near several pharmacies but still experience poor access if the prescribed medicine is expensive. Likewise, an inexpensive medicine may not provide a suitable solution if the patient does not have adequate information about its use.

The Indian healthcare system makes affordability especially important. A substantial proportion of healthcare expenditure is borne directly by households. Consequently, the cost of medicines can influence treatment decisions. Some households may purchase smaller quantities, select cheaper alternatives or delay treatment when the cost becomes difficult to manage.

Medication practices form the behavioural link between access and health outcomes. Appropriate practice generally involves following professional instructions, taking medicines at the recommended dose and completing treatment where required. Problems can arise when individuals change doses, discontinue treatment after symptoms improve, use medicines prescribed for someone else or continue an old prescription without reassessment.

The pandemic created conditions that could encourage such practices. Limited access to doctors made previous prescriptions more valuable to patients who were unable to obtain new consultations. Pharmacies offered a convenient source of medicines and basic advice. Digital information also became more accessible. While these alternatives helped people manage some immediate problems, they could not fully substitute for professional diagnosis and clinical assessment.

Self-medication deserves particular attention because it can have both practical and harmful dimensions. For minor, familiar conditions, people may appropriately use medicines that are available without prescription. The concern arises when individuals use prescription medicines without current medical advice or attempt to treat conditions that require diagnosis.

Antibiotic use is an important example. COVID-19 is a viral infection, and antibiotics do not treat the virus itself. Nevertheless, inappropriate antibiotic use can occur when individuals assume that symptoms such as fever or cough require antibiotic treatment. Such practices contribute to antimicrobial resistance and may expose patients to unnecessary adverse effects.

Healthcare-seeking behaviour is influenced by several factors. People assess their symptoms, perceived seriousness, cost, convenience and previous experiences before deciding whether to seek professional treatment. During COVID-19, fear of infection became an additional barrier. The perceived risk of visiting a hospital could outweigh the perceived benefit when symptoms appeared mild.

Telemedicine reduced some of these barriers. Remote consultations allowed patients to communicate with doctors without travelling to healthcare facilities. This was particularly useful for follow-up consultations and some routine health concerns. However, telemedicine has limitations when physical examination, diagnostic testing or emergency intervention is necessary.

Pharmacies also occupy an important position in community healthcare. Their widespread availability and convenient operating hours make them accessible to people who may find formal consultation difficult. This can be beneficial when pharmacists provide appropriate guidance and referral. However, pharmacy-based treatment should not replace medical consultation when symptoms require professional diagnosis.

The literature therefore indicates that healthcare recovery has several dimensions. Restoration of hospital services is important, but it does not automatically restore appropriate medicine use or healthcare-seeking behaviour. The long-term impact of the pandemic must also be accessed through household experiences.

3. Community Health Recovery

Community health recovery can be understood as the restoration of the conditions that allow people to obtain appropriate healthcare and maintain treatment without unnecessary barriers. In this context, recovery includes the availability of medicines, affordability of treatment, responsible medicine use and timely healthcare-seeking.

These components influence one another. Difficulty obtaining medicines can encourage people to search for alternatives or depend on self-medication. Financial barriers may lead to delayed consultation. Previous negative experiences with healthcare facilities may reduce willingness to seek professional treatment in the future. Conversely, affordable medicines and accessible healthcare can encourage appropriate treatment behaviour.

The relationship among the major variables can be summarized as follows:

Pharmaceutical accessibility → Medication practices → Healthcare behaviour → Community health recovery

The relationship is not entirely one-directional. Healthcare behaviour can also affect medication practices. For example, a person who regularly consults a doctor is more likely to receive updated

prescriptions and professional guidance. A person who avoids consultation may depend more heavily on previous prescriptions or informal advice.

Table 2. Relationship among the Major Study Variables

Study variable	Main elements	Contribution to recovery
Pharmaceutical accessibility	Availability, affordability, continuity, information	Supports uninterrupted treatment
Medication practices	Adherence, self-medication, previous prescriptions, substitution	Influences safety and effectiveness of treatment
Healthcare behaviour	Consultation, provider choice, delay, information-seeking	Influences timely use of healthcare
Community health recovery	Continuity, appropriate treatment and healthcare utilization	Reflects overall recovery

Source: Developed by the author based on Andersen (1995), WHO (2020), WHO (2023), and the conceptual framework of the present study.

This integrated perspective is particularly useful for understanding the post-COVID period. A healthcare system may have resumed normal operations while some households continue to experience financial or behavioural barriers. Recovery should therefore be assessed from the perspective of both healthcare institutions and community members.

4. Rationale, Objectives and Research Questions

The need for the present study arises from the fact that pharmaceutical accessibility, medication practices and healthcare-seeking are frequently examined independently. In everyday life, however, these factors are closely connected. Difficulty in accessing a doctor may encourage pharmacy-based treatment, while medicine costs may influence whether treatment is completed.

The pandemic provided a unique environment in which these relationships became more visible. Examining the post-COVID period can help determine whether the practices developed during the emergency were temporary responses or have become part of routine healthcare behaviour.

The study is particularly relevant to Dehradun because the city has a diverse healthcare system and serves populations from both the city and nearby areas. Understanding local patterns can provide useful information for community health planning.

Objectives

The study aims to examine the role of pharmaceutical accessibility, medication practices and healthcare behaviour in community health recovery after COVID-19 in Dehradun. Specifically, it seeks to:

1. Examine the availability, affordability and continuity of medicines;
2. Understand medication practices, including adherence and self-medication;
3. Examine changes in healthcare-seeking behaviour after COVID-19;
4. Identify economic, informational and accessibility factors influencing medicine use and healthcare decisions;
5. Examine the relationship among pharmaceutical accessibility, medication practices and healthcare behaviour; and
6. Suggest measures for strengthening community health recovery.

Research Questions

The study addresses six main questions:

1. How accessible are medicines to residents of Dehradun?
2. What medication practices have continued or developed after COVID-19?
3. To what extent do people rely on self-medication or previous prescriptions?
4. What factors influence post-COVID healthcare-seeking behaviour?
5. How are pharmaceutical accessibility and medication practices related to healthcare behaviour?
6. What measures can strengthen community health recovery in Dehradun?

5. Theoretical Framework

The relationship between access to medicines and healthcare behaviour can be understood through established healthcare-utilization perspectives. The present study mainly draws on the **Andersen Healthcare Utilization Model**, which explains healthcare use through predisposing, enabling and need-related factors. Age, education and health beliefs may influence a person's tendency to seek

care, while income, availability of services, transportation and affordability determine whether healthcare can actually be accessed. Perceived illness and its seriousness further influence the decision to seek treatment.

This framework is relevant because pharmaceutical accessibility can be considered an enabling factor. Even when a person recognizes the need for treatment, limited income, medicine prices or difficulty reaching a pharmacy may prevent appropriate treatment. Similarly, easy access to pharmacies may encourage individuals to manage minor illnesses without visiting a doctor.

The study also draws on the **Health Belief Model**, particularly its emphasis on perceived susceptibility, perceived severity, perceived benefits and perceived barriers. COVID-19 changed people's perception of healthcare facilities. During the pandemic, hospitals could be perceived not only as places of treatment but also as places where infection might be encountered. Such perceptions could discourage consultation even when healthcare was available.

These perspectives together provide a useful explanation of post-COVID healthcare behaviour. Structural conditions determine the options available to individuals, while personal beliefs, previous experiences and perceived risks influence the choices they make.

6. Research Framework and Hypotheses

The study considers pharmaceutical accessibility as an important factor influencing medication practices and healthcare behaviour. Accessibility includes availability, affordability, continuity and access to reliable information. Medication practices include adherence to prescriptions, self-medication, use of previous prescriptions and medicine substitution. Healthcare behaviour includes decisions concerning consultation, provider choice, delayed treatment and reliance on informal sources.

The proposed relationship can be represented as:

Pharmaceutical Accessibility → Medication Practices → Healthcare Behaviour → Community Health Recovery

A direct relationship between pharmaceutical accessibility and healthcare behaviour is also possible. For example, readily available medicines may encourage people to treat minor symptoms themselves, whereas difficulty obtaining medicines

may lead some individuals to seek alternative providers or delay treatment.

The framework also recognizes the influence of socioeconomic characteristics. Income can affect affordability, education may influence understanding of medicines, and age or chronic health requirements may increase the need for regular treatment.

The following hypotheses are proposed for empirical testing:

H1: Pharmaceutical accessibility is significantly associated with medication practices among residents of Dehradun.

H2: Pharmaceutical accessibility is significantly associated with healthcare-seeking behaviour.

H3: Medication practices are significantly associated with healthcare-seeking behaviour.

H4: Medicine affordability significantly influences continuity of medication use.

H5: Self-medication is associated with delayed professional healthcare consultation.

H6: Socioeconomic characteristics significantly influence pharmaceutical accessibility and healthcare behaviour.

7. Research Methodology

The study adopts a **descriptive and analytical research design**. Its purpose is to understand the post-COVID experiences of community members concerning access to medicines, medicine-use practices and healthcare-seeking behaviour. A literature-based conceptual approach is used in the present paper, while the proposed framework can also be tested through primary survey data.

The study focuses on **Dehradun, Uttarakhand**, because the city has a mixed healthcare environment comprising government and private hospitals, clinics, pharmacies, diagnostic facilities and specialist services. It also attracts patients from neighbouring areas. These characteristics make it suitable for examining differences in healthcare access and preferences.

The target population for a future empirical survey would consist of adult residents who have used medicines or healthcare services during the post-COVID period. A structured questionnaire could be administered across different localities to capture

variation in income, occupation, education and healthcare access. The questionnaire would contain questions on medicine availability, affordability, continuity, self-medication, prescription adherence, healthcare consultation and use of telemedicine.

Secondary information for the present study is drawn from government reports, international health organizations and published academic literature. Sources concerning healthcare disruption, pharmaceutical access, medication adherence and healthcare-seeking behaviour provide the basis for the conceptual discussion.

For quantitative analysis, frequencies and percentages can first be used to describe respondents' characteristics and healthcare practices. Cross-tabulation can then be used to examine differences between demographic groups. Chi-square tests can assess associations between categorical variables, while correlation and regression techniques may be applied where the data permit. A 5% significance level may be adopted for hypothesis testing.

Because a cross-sectional survey captures responses at a particular point in time, statistical association should not be interpreted automatically as causation. Longitudinal research would be more suitable for establishing how behaviour changes over an extended period.

Ethical principles should be followed if primary data are collected. Participation should be voluntary, respondents should be informed about the purpose of the research, and personal information should be kept confidential. Questions concerning health experiences should be handled with appropriate privacy.

8. Analysis and Discussion

The literature reviewed in this study indicates that post-COVID community health recovery involves more than restoration of physical healthcare facilities. Access to medicines, the way those medicines are used and decisions about seeking professional care are closely connected.

Pharmaceutical Accessibility

The pandemic highlighted weaknesses in medicine access that may not be visible under normal circumstances. During periods of restricted movement, patients could face difficulties reaching pharmacies, obtaining prescriptions or purchasing medicines. These difficulties were particularly

important for people requiring continuous treatment.

In the post-COVID period, physical access to medicines in an urban centre such as Dehradun is likely to be better than during lockdown. However, physical availability does not necessarily mean effective accessibility. Medicine prices remain important, particularly for households with limited income or people requiring long-term treatment.

Affordability can affect treatment in several ways. A household may purchase a smaller quantity than prescribed, choose a cheaper alternative, postpone a purchase or discontinue treatment once symptoms improve. Such decisions may be driven by economic necessity rather than a lack of understanding.

Information is another part of pharmaceutical accessibility. The availability of medicines without adequate information does not ensure safe treatment. Patients may receive advice from pharmacists, relatives, friends, websites or social media. While some information may be useful, unreliable advice can contribute to inappropriate medicine selection.

The post-COVID period therefore requires attention to both supply and demand. Ensuring that medicines are stocked is important, but so is ensuring that people can afford them and understand when professional advice is necessary.

Medication Practices

The pandemic changed the circumstances under which people used medicines. Regular doctor visits became difficult for many patients, which encouraged some to continue earlier prescriptions or seek advice from alternative sources. Such behaviour may have helped maintain treatment during a period of limited access.

However, the continuation of an old prescription is not always appropriate. A medicine prescribed for an earlier condition may not be suitable for a new illness, and the required dose may change over time. Consequently, what began as a practical response to restricted access can become unsafe if it continues without professional review.

Self-medication is another important issue. People often use medicines based on previous experience, advice from family members or pharmacists, or information found online. The practice can save time and reduce consultation costs, but it may also result in incorrect treatment or delayed diagnosis.

Antibiotic use deserves particular attention. Antibiotics are intended for bacterial infections and do not treat COVID-19 itself. Their unnecessary use can expose individuals to avoidable adverse effects and contributes to antimicrobial resistance. Public awareness should therefore focus on appropriate use rather than simply increasing medicine availability.

Medication adherence is equally important. Access barriers may cause non-adherence even when the patient wants to follow medical advice. This distinction matters because the solution to economically driven non-adherence is different from the solution to deliberate refusal of treatment. Reducing medicine costs and improving continuity can therefore contribute directly to better treatment adherence.

Healthcare Behaviour

Healthcare-seeking behaviour changed considerably during the pandemic. Fear of infection, restricted transportation, financial uncertainty and concerns about crowded facilities encouraged some individuals to postpone consultations. Minor illnesses were often managed at home or through pharmacies.

Some of these patterns may continue after the emergency. People who found pharmacy-based treatment convenient may continue using pharmacies as their first point of contact. Similarly, individuals who experienced teleconsultation may prefer it for follow-up visits or routine advice.

Such changes are not necessarily negative. Healthcare systems can benefit when appropriate cases are managed efficiently through primary care, pharmacy services or telemedicine. The concern arises when these alternatives replace professional assessment for conditions that require physical examination, diagnostic testing or urgent treatment.

Trust also plays a role. Individuals who have confidence in a particular doctor, hospital or pharmacy are more likely to return to that provider. Previous experiences with waiting time, cost and quality of care can influence future choices.

The recovery of healthcare behaviour is therefore likely to be uneven. Some people may have returned completely to pre-pandemic patterns, while others may have adopted a combination of conventional consultation, pharmacy advice and digital healthcare.

9. Illustrative Analysis

Since the present manuscript does not contain an original field-survey dataset, the numerical results in this section are presented only as an **illustrative empirical model**. They demonstrate how the findings can be presented after actual data collection and should not be interpreted as the actual responses of Dehradun residents.

The analysis focuses on six indicators directly related to the study variables: medicine availability, medicine affordability, self-medication, delayed healthcare consultation, pharmacy use and teleconsultation. These indicators represent the three central dimensions of the study—pharmaceutical accessibility, medication practices and healthcare behaviour.

For the purpose of illustration, a hypothetical sample of 200 respondents is considered. The percentages are used consistently in the discussion and in Figure 1.


Indicator	Illustrative percentage	Interpretation
Medicines generally available when required	78%	Medicine availability appears relatively strong
Medicines considered affordable	61%	Affordability remains a concern for some households
Practised self-medication	46%	Self-medication remains an important practice
Delayed healthcare consultation	38%	A considerable proportion may postpone professional care
Used pharmacy as first source for minor illness	57%	Pharmacies remain important first-contact points
Used teleconsultation after COVID-19	31%	Digital healthcare continues to have some relevance

The illustrative pattern indicates that **medicine availability is higher than medicine affordability**. This distinction is important because physical availability alone does not guarantee effective pharmaceutical access. A medicine may be readily available but still remain difficult to obtain for households facing financial constraints.

The results also indicate the continuing importance of behavioural factors. The illustrative self-medication rate of 46% suggests that independent treatment could remain an important feature of post-COVID healthcare behaviour. Similarly, 38%

delayed professional consultation, indicating that accessibility, convenience, cost or perceptions about illness may continue to influence healthcare decisions.

Pharmacies appear to have an important community role, with 57% of the illustrative respondents using them as the first source for minor illness. Teleconsultation, at 31%, represents a smaller but still meaningful component of post-COVID healthcare use.



Source: Author's illustrative presentation based on the conceptual framework of the study; percentages are hypothetical and should be replaced with actual field-survey results.

10. Implications for Community Healthcare

The illustrative analysis suggests that community health recovery requires attention to both pharmaceutical access and healthcare behaviour. The relatively high level of medicine availability, compared with affordability, indicates that physical access alone may not ensure effective treatment. Economic barriers can still influence whether individuals purchase medicines regularly or continue prescribed treatment.

Community education should therefore focus on rational medicine use, particularly the risks associated with inappropriate antibiotics, unsupervised prescription-medicine use and prolonged self-medication. Information should also help people understand when professional medical consultation is necessary.

Pharmacies have an important role because they are easily accessible and may be the first source of advice for minor illnesses. However, pharmacy-based support should be accompanied by appropriate referral when symptoms are serious, persistent or unclear. Stronger coordination between pharmacies and healthcare providers can improve continuity of care.

The continued use of teleconsultation also indicates that digital healthcare can remain useful after the

pandemic. It can support follow-up consultations and routine advice, while patients requiring physical examination or diagnostic testing should be referred to appropriate healthcare facilities.

Affordable primary healthcare is equally important. When consultation is accessible in terms of cost, distance and waiting time, individuals may be less likely to depend entirely on self-medication or delay professional treatment.

11. Major Findings

The conceptual and illustrative analysis produces the following major findings:

1. **Medicine availability appears relatively strong.** The illustrative result of 78% suggests that physical access to medicines may have recovered considerably after the pandemic.
2. **Affordability remains an important concern.** Although 61% of the illustrative respondents considered medicines affordable, the lower percentage compared with availability indicates that financial barriers may continue to affect effective pharmaceutical access.
3. **Self-medication remains significant.** The illustrative figure of 46% indicates that independent medicine use may continue to influence post-COVID medication practices.
4. **Pharmacies remain important healthcare contact points.** The illustrative finding of 57% indicates that many people may approach pharmacies first for minor illnesses because of convenience and accessibility.
5. **Delayed healthcare consultation remains a concern.** The illustrative figure of 38% suggests that a considerable proportion of individuals may postpone professional consultation.
6. **Teleconsultation continues to have a role.** The illustrative result of 31% indicates that digital healthcare practices developed during the pandemic may continue during the recovery period.
7. **Pharmaceutical accessibility and healthcare behaviour are interconnected.** Availability and affordability influence how individuals obtain and use medicines, while medication practices can affect decisions about professional consultation.
8. **Post-COVID recovery is both institutional and behavioural.** Restoration of healthcare facilities does

not necessarily mean that all pandemic-related healthcare practices have returned to earlier patterns.

12. Conclusion

The COVID-19 pandemic disrupted routine healthcare and changed the way many individuals obtained medicines, managed illness and sought professional care. The effects extended beyond hospitals to pharmacies, households and community-level healthcare practices.

This study examined community health recovery in Dehradun through pharmaceutical accessibility, medication practices and healthcare behaviour. The analysis shows that these dimensions are closely connected. Medicine availability is important, but affordability and continuity determine whether people can actually maintain treatment. Similarly, self-medication and pharmacy-based care may provide convenience, but inappropriate use or delayed consultation can create health risks.

The illustrative analysis used in this paper indicates six important patterns: medicines are generally available to a large proportion of respondents (78%), affordability is somewhat lower (61%), self-medication remains common (46%), delayed consultation is reported by a considerable proportion (38%), pharmacies are frequently used as the first source for minor illness (57%), and teleconsultation continues to be used (31%). These percentages are hypothetical and are intended only to demonstrate the proposed empirical presentation.

The findings suggest that post-COVID recovery should not be measured only by the restoration of hospitals and routine medical services. Effective recovery also requires affordable medicines, appropriate medication practices, timely healthcare-seeking and reliable health information.

For Dehradun, stronger coordination among pharmacies, primary healthcare providers, hospitals and digital healthcare services could improve community-level healthcare. Public awareness programmes should particularly address unsafe self-medication, inappropriate antibiotic use and unnecessary delays in seeking professional care.

Overall, **pharmaceutical accessibility, medication practices and healthcare behaviour should be treated as interconnected components of community health recovery.** Addressing these areas together can contribute to a more accessible, responsive and resilient healthcare system.

13. Recommendations

The study recommends the following measures for strengthening post-COVID community healthcare in Dehradun:

1. Improve the continuous availability of essential medicines at community level.
2. Reduce financial barriers by increasing awareness of affordable and quality generic medicines.
3. Conduct community awareness programmes on responsible self-medication and appropriate antibiotic use.
4. Encourage pharmacies to refer individuals to qualified healthcare professionals when symptoms require clinical assessment.
5. Improve the accessibility and affordability of primary healthcare services to reduce unnecessary treatment delays.
6. Continue telemedicine for appropriate follow-up and routine consultations, while maintaining referral pathways for physical examination and diagnostic services.
7. Give particular attention to households experiencing financial difficulty in maintaining regular medication.
8. Monitor medicine availability and community healthcare barriers during future public health emergencies.
9. Include self-medication, delayed consultation, pharmacy use and teleconsultation in local assessments of healthcare behaviour.
10. Conduct further empirical research comparing post-COVID healthcare behaviour in urban and rural areas of Uttarakhand.

14. Limitations and Future Research

The present study is primarily conceptual and literature-based and therefore does not claim that the illustrative percentages represent the actual population of Dehradun. The numerical values used in Section 9 are hypothetical and must be replaced with genuine survey findings before the paper is submitted as an empirical study.

The cross-sectional approach proposed for future primary research would also limit the ability to establish long-term changes in healthcare behaviour. Respondents may have difficulty recalling their experiences during the pandemic, and self-reported responses may be affected by memory or social desirability.

Future research should collect primary data from different parts of Dehradun and examine

differences according to age, education, income and healthcare requirements. Longitudinal studies could compare healthcare practices during the pandemic, immediately after the pandemic and several years into the recovery period.

Comparative research involving rural areas and smaller towns of Uttarakhand would also help identify differences in medicine availability, affordability, geographical access and healthcare-seeking behaviour. Qualitative interviews with pharmacists, doctors and community members could further explain why people choose self-medication, delay consultation or prefer particular healthcare providers.

References

1. World Health Organization. (2024). *Global report on primary health care: Close to communities, closer to people*. Geneva: World Health Organization.
2. World Health Organization. (2024). *World health statistics 2024: Monitoring health for the SDGs, sustainable development goals*. Geneva: World Health Organization.
3. World Health Organization. (2023). *Fourth round of the global pulse survey on continuity of essential health services during the COVID-19 pandemic*. Geneva: World Health Organization.
4. World Health Organization. (2023). *Health systems resilience toolkit: A new perspective on health system resilience*. Geneva: World Health Organization.
5. Government of India, Ministry of Health and Family Welfare. (2024). *National Health Accounts Estimates for India 2021–22*. New Delhi: Government of India.
6. World Health Organization. (2022). *The impact of the COVID-19 pandemic on noncommunicable disease resources and services*. Geneva: World Health Organization.
7. World Health Organization Regional Office for South-East Asia. (2022). *Medicines and health technologies*. New Delhi: WHO South-East Asia Regional Office.
8. Zhang, Y., et al. (2022). Medication adherence and healthcare utilization during the COVID-19 pandemic: A systematic review. *Patient Preference and Adherence*.
9. Kretchy, I. A., Asiedu-Danso, M., & Kretchy, J. P. (2021). Medication management and adherence during the

- COVID-19 pandemic. *Clinical and Translational Science*, 14(3), 601–610.
10. World Health Organization. (2021). *COVID-19 and noncommunicable diseases*. Geneva: World Health Organization.
11. Ministry of Health and Family Welfare. (2020). *Telemedicine practice guidelines: Enabling registered medical practitioners to provide healthcare using telemedicine*. New Delhi: Government of India.
12. International Pharmaceutical Federation. (2020). *Pharmacy at a glance: COVID-19 and pharmacy practice*. The Hague: International Pharmaceutical Federation.
13. World Health Organization. (2020). *Access to essential medicines and health technologies*. Geneva: World Health Organization.
14. Chatterjee, P., Nagi, N., Awasthi, S., et al. (2020). The 2019 novel coronavirus disease (COVID-19) pandemic: A review of the current evidence. *Indian Journal of Medical Research*, 151(2–3), 147–159.
15. Yadav, S., Rawal, G., & Baxi, M. (2020). Self-medication practices in the COVID-19 pandemic: A review. *Journal of Family Medicine and Primary Care*, 9(7).
16. Organisation for Economic Co-operation and Development. (2020). *Health at a glance: Asia/Pacific 2020*. Paris: OECD Publishing.
17. World Health Organization. (2019). *WHO guideline on country pharmaceutical pricing policies*. Geneva: World Health Organization.
18. World Health Organization. (2019). *WHO global action plan on physical activity 2018–2030: More active people for a healthier world*. Geneva: World Health Organization.
19. World Health Organization. (2017). *WHO report on surveillance of antibiotic consumption: 2016–2018 early implementation*. Geneva: World Health Organization.
20. World Health Organization. (2016). *Global action plan on antimicrobial resistance*. Geneva: World Health Organization.
21. World Health Organization. (2015). *Global action framework for strengthening the role of health systems in improving access to essential medicines*. Geneva: World Health Organization.
22. World Health Organization. (2014). *Access to affordable medicines: A priority for achieving universal health coverage*. Geneva: World Health Organization.
23. World Health Organization. (2011). *The world medicines situation 2011: Medicines prices, availability and affordability*. Geneva: World Health Organization.
24. World Health Organization. (2004). *The world medicines situation*. Geneva: World Health Organization.
25. Andersen, R. M. (1995). Revisiting the behavioral model and access to medical care: Does it matter? *Journal of Health and Social Behavior*, 36(1), 1–10.
26. Rosenstock, I. M., Strecher, V. J., & Becker, M. H. (1988). Social learning theory and the Health Belief Model. *Health Education Quarterly*, 15(2), 175–183.
27. Hardon, A., Hodgkin, C., & Fresle, D. (2004). *How to investigate the use of medicines by consumers*. Geneva: World Health Organization and University of Amsterdam.
28. Horne, R., Weinman, J., Barber, N., Elliott, R., & Morgan, M. (2005). *Concordance, adherence and compliance in medicine taking*. London: National Co-ordinating Centre for NHS Service Delivery and Organisation.
29. Wirtz, V. J., Hogerzeil, H. V., Gray, A. L., et al. (2017). Essential medicines for universal health coverage. *The Lancet*, 389(10067), 403–476.
30. Kruk, M. E., Gage, A. D., Arsenault, C., et al. (2018). High-quality health systems in the Sustainable Development Goals era: Time for a revolution. *The Lancet Global Health*, 6(11), e1196–e1252.