Iron Oxide Nanoparticle-Assisted Green Synthesis of Nitriles from Aldehydes in Water: Scope, Selectivity and Catalyst Reusability" Research

Shraddha Upadhyay [1]*, Anjali [2]
[2]: Keral Verma Subharti College of Science, Swami Vivekanand Subharti University, Meerut, Uttar Pradesh, India – 250005
[1]: Keral Verma Subharti College of Science, Swami Vivekanand Subharti University, Meerut, Uttar Pradesh, India – 250005
* Corresponding author: dr.shrupa@gmail.com

Abstract: Interestingly, One pot conversion of heteroaromatic 2-chloro-3-formylquinoline into their cyano derivatives using first time Fe_2O_3 nanoparticle in aqueous ammonia has been discussed.

Keywords: One pot, oxidant, nitrile, transformation, 2-chloro-3- formyl-quinolines, Nanoparticle
How to cite this article: Shraddha Upadhyay 1* Anjali 2. Iron Oxide Nanoparticle-Assisted Green Synthesis of Nitriles from Aldehydes in Water: Scope, Selectivity and Catalyst Reusability". Int J Drug Deliv Technol. 2026;16(87s):1210-1212. DOI: 10.25258/ijddt.16.87s.125.

In organic synthesis functional group transformations have significant importance[1] amongst, nitriles, represent an important reagen. ts for organic synthesis, have been attracted great attention to chemist for long time because of their transformation into variety of heterocycles such as thiazoles[2a], 2-oxazolines[2b-c], tetrazoles[2d], imidazoles[2e], triazoloannulatedpyri- midines[2f] and others[2g-h]. Formations of nitriles from the corresponding aldehydes are an important functional group transformation. Methods known in the literature either involve the initial conversion of aldehydes to aldoximes followed by dehydration[3] to give nitriles or direct conversion of aldehydes into nitriles without isolation of nitrogen containing intermediates, by treating with variety of chemicals[4]. The use of ammonia combined with different oxidants such as oxygen[5], lead tetraacetate[6], iodine[7] and hydrogen peroxide[8], is considered amongst an appropriate method for the transformation of aliphatic and aromatic aldehydes to their corresponding nitriles. These observations inspired us to investigate the transformation of 3-quinolinylformaldehyde 1a to 3- quinolinylnitrile 2a (ref.9f) in aqueous ammonia with easily available oxidants.

In continuation to the studies in carbo/hetero annulatedquinolines synthesis[9] from 2-chloro-3-formylquinolines **1a**. 2-chloro-3-cyanoquinolines **2a** is required to explore analogous reactions on them for synthesis of carbo/hetero annulatedquinolines. Thus, in this paper one pot synthesis of 2-chloro-3-cyanoquinolines from 2-chloro-3-formylquinolines in aqueous ammonia using cheap and easily available iodine as oxidant and Fe_2O_3 nanoparticle as catalyst is described.

The reaction of 2-chloro-3-formylquinoline **1a** with iodine in aqueous ammonia along with Fe_2O_3 nanoparticle was chosen as model and the reaction condition was optimized. It is found that reaction proceeded well and completed in a short period at RT. The better yield of product, 2-chloro-3-cyanoquinoline **2a**, was obtained by using 6 mL of aqueous ammonia (30% solution, **Table I**). Thus, the reaction of 2-chloro-3-formylquinoline **1a** in THF (5 mL) was treated with aqueous ammonia (6 mL), Fe_2O_3 nanoparticle (0.5mmole) and iodine (1.1 mmole) at RT for 75 minutes to afford the desired nitrile **2a** in good yield. The structure of compound **2a** was characterized from its NMR and IR spectra. The ^{1}H NMR spectrum of compound **2a** showed absence of formyl peak at δ 10.5 and its IR spectrum showed absence of carbonyl absorption at 1687 $cm^{-1}$and presence of weak absorption of nitrile group at 2226 cm^{-1}. After optimizing reaction conditions for transformation of **1a** into **2a**, reaction of other substrates **1b-k** were carried out. These afforded the corresponding nitriles **2b-k** in a short time (20-85 min) in good yields (**Table I**).

In summary, a general and simple method for the transformation of carbaldehydes into nitriles in a better yield in aqueous medium is described using cheap and easily available reagents.











Experimental Section

Melting points were measured in an open capillary tube with a Buchi melting point apparatus and are uncorrected. Elemental analysis was obtained using Perkin-Elmer 240C CHN-analyzer. IR spectra were recorded on a FT/IR-5300 (JASCO) spectro- photometer; ^{1}H NMR spectra in $CDCl_3$/DMSO-d_6 at
300 MHz on a Jeol AL-300 spectrometer (chemical shifts in δ ppm) relative to TMS as an internal standard. Reactions were monitored by TLC, using silica gel $PF_{254+366}$ as an adsorbent and ethyl acetate- hexane in different ratios as eluent.

General method for conversion of carbaldehyde into nitrile

To a stirred solution of formyl compound (1 mmole) in MeOH (2 mL) was added aq. NH_3 (30%, 6 mL), Cat.Fe_2O_3 Nanoparticle followed by sublimed I_2 (1 mmole) and stirred the reaction- mixture at RT for the mentioned period

R = H, Me, MeO, Et

Entry	Substrate	Product	Time (min)	Yield (%)
1	 **1a**	**2a**	75	82
2	 **1b**	**2b**	72	80
3	 **1c**	**2c**	20	78
4	 **1d**	**2d**	68	80
5	 **1c**	**2e**	85	83
6	 **1f**	**2f**	72	79
7	 **1g**	**2g**	60	80
8	 **1h**	**2h**	75	73
9	 **1i**	**2i**	68	77
10	 **1j**	**2j**	50	67

(**Table I**). The dark violet/brown solution became colourless after completion of reaction (checked by TLC).The reaction-mixture was treated with aqueous $Na_2S_2O_3$ solution (5 mL of 5% solution) and extracted with ethyl acetate. The solution was dried, filtered and evaporated under vacuum to give pure
nitrile product.

2-Chloro-3-cyanoquinoline, 2a: Yield 82%, m.p. 144-45°C; IR (KBr): 2226 cm^{-1}; ^1H NMR ($CDCl_3$):
δ 7.70 (t, *J* = 7.8 Hz, 7-H), 7.90 (d, *J* = 8.1 Hz, 5-H), 7.92, (t, *J* = 8.1 Hz, 6-H), 8.09 (d, *J* = 8.1 Hz, 8-H), 8.57 (s, 4-H); Anal. Calcd for $C_{10}H_5N_2Cl$: C, 63.82; H, 2.68; N,14.90. Found: C, 63.75; H, 2.70; N, 14.82%.

6-Methyl-2-chloro-3-cyanoquinoline, 2b: Yield 80%, m.p. 142-45°C; IR (KBr): 2228 cm^{-1}; ^1H NMR ($CDCl_3$): δ 2.58 (s, CH_3), 7.65 (s, 5-H), 7.73 (d, *J* =8.7 Hz. 7-H), 7.97 (d, *J* = 8.1 Hz, 8-H), 8.46 (s, 4-H). Anal. Calcd for $C_{11}H_7N_2Cl$: C, 65.34; H, 3.49; N,13.86. Found: C, 65.21; H, 3.63; N, 13.48%.

6- Methoxy-2-chloro-3-cyanoquinoline, 2c: Yield 78%, m.p. 157-60°C; IR (KBr): 2231 cm^{-1}; ^1H NMR ($CDCl_3$): δ 3.96 (s, OCH_3), 7.10 (d, *J* = 2.4 Hz, 5-H), 7.54 (dd, *J* = 2.7, 9.0 Hz, 7-H), 7.97 (d, *J* = 9.3 Hz, 8- H), 8.43 (s, 4-H); Anal. Calcd for $C_{11}H_7N_2OCl$: C, 60.54; H, 3.24; N, 12.85. Found: C, 60.35; H, 3.18; N,
12.90%.

7- Methyl-2-chloro-3-cyanoquinoline, 2d: Yield 80%, m.p. 142-45°C; IR (KBr): 2222 cm^{-1}; ^1H NMR ($CDCl_3$): δ 2.62 (s, CH_3), 7.52 (d, *J* = 8.1 Hz, 5-H), 7.78 (d, *J* = 8.4 Hz, 6-H), 7.85 (s, 8-H), 8.50 (s, 4-H); Anal. Calcd for $C_{11}H_7N_2Cl$: C, 65.34; H, 3.49; N,
13.86. Found: C, 65.40; H, 3.41; N, 13.78%.

7- Methoxy-2-chloro-3-cyanoquinoline, 2e: Yield 83%, m.p. 147°C; IR (KBr): 2229 cm^{-1}; ^1H NMR ($CDCl_3$): δ 3.99 (s, OCH_3), 7.44 (s, 8-H), 7.46 (d, *J* =13.5 Hz, 6-H), 7.84 (d, *J* = 9.0 Hz, 5-H), 8.67 (s, 4- H); Anal. Calcd for $C_{11}H_7N_2OCl$: C, 60.54; H, 3.24;N, 12.85. Found: C, 60.46; H, 3.21; N, 12.78%.

8- Methyl-2-chloro-3-cyanoquinoline, 2f: Yield 79%, m.p. 95-97°C; IR (KBr): 2228 cm^{-1}; ^1H NMR ($CDCl_3$): δ 2.77 (s, CH_3), 7.59 (t, *J* = 7.5 Hz, 6-H),7.74 (t, *J* = 7.2 Hz, 5-H, 7-H), 8.53 (s, 4-H); Anal. Calcd for $C_{11}H_7N_2Cl$: C, 65.34; H, 3.49; N,13.86.

Found: C, 65.41; H, 3.42; N, 13.90%.

8- Ethyl-2-chloro-3-cyanoquinoline, 2g: Yield 80%, m.p. 124°C; IR (KBr): 2231 cm^{-1}; ^1H NMR ($CDCl_3$): δ 1.36 (t, 7.74, *J* = 7.5 Hz, CH_3), 3.25 (q, *J* = 7.5 Hz, CH_2), 7.63 (t, *J* = 7.5 Hz, 6-H), 7.75 (t, *J* = 7.2 Hz, 5-H, 7-H),8.53 (s, 4-H); Anal. Calcd for $C_{12}H_9N_2Cl$: C, 66.65; H, 4.20; N, 12.96. Found: C, 66.48; H, 4.28; N, 13.02%.

2-Chloro-3-cyano-5-phenylpyridine, 2h: Yield 73%, m.p. 150-52°C; IR (KBr): 223 cm^{-1}; ^1H NMR ($CDCl_3$): δ 7.50-7.54 (m, Ph-H), 8.15 (d, *J* = 2.7 Hz,), 8.79 (d, *J* = 2.7 Hz, 6-H); Anal. Calcd for $C_{12}H_7N_2Cl$: C, 67.28; H, 3.30; N, 13.09. Found: C,67.34; H, 3.22; N, 12.98%.

2-Chloro-3-cyano-4-methyl-N-phenylpyrazole, 2i: Yield 77%, m.p. 105-07°C; IR (KBr): 2227 cm^{-1};^1H NMR ($CDCl_3$): δ 2.57 (s, CH_3), 7.50 (m, Ph-H); Anal. Calcd for $C_{12}H_9N_2Cl$: C, 66.65; H, 4.20; N,12.96. Found: C, 66.48; H, 4.09; N, 13.04%.

3,4-Dimethoxy-benzonitrile, 2j: Yield 67%, m.p. 67-69°C; IR (KBr): 2226 cm^{-1}; ^1H NMR ($CDCl_3$): δ 3.90 (s, OCH_3), 3.94 (s, OCH_3), 6.90 (d, *J* = 8.4 Hz, 5-H), 7.26 (s, 2-H), 7.29 (d, *J* = 8.4 Hz, 6-H). Anal. Calcd for $C_9H_9NO_2$: C, 66.23; H, 5.56; N, 8.59. Found: C, 66.38; H, 5.48; N, 8.52%.

Acknowledgement

Authors thank Department of Chemistry, IIT Roorkee for providing spectral and analytical facilities. and Department of Chemistry, Swami Vivekananda Subharti University, Meerut, India, for providing laboratory and research facilities to carry out this research work.

Funding: The authors received no specific financial support for the research, authorship, and/or publication of this work.

Conflict of Interest: The authors declare that there is no conflict of interest regarding the publication of this manuscript.

Ethical Approval: This study did not involve human participants, animals, or clinical samples; therefore, ethical approval was not required.

Consent to Participate: Not applicable.

Consent for Publication: All authors have reviewed and approved the manuscript for publication.

Data Availability Statement: The data supporting the findings of this study are available within the article. Additional data may be provided by the corresponding author upon reasonable request.

References

1 Erman M B, Snow J W & Williams M J, *Tetrahedron Let* 41, **2000**, 6749.

2 Chichiro M, Nagamoto H, Takemura L, Kitano K, Komatsu H, Sekiguchi K, Tabusa F, Mori T, Tominnaga M & Yobuuchi Y, *J Med Chem*, *38*, **1995**, 353; (b) Serrano J L, Sierra T, Gonzalez Y, Bolm C, Weickhardt K & Magnus A, *J Am Chem Soc*, *117*, **1995**, 8312; (c) Jnaneshwara G K, Deshpande V H, Lalithambika, Ravindranathan T & Bedekar A V, *Tetrahedran Lett*, *39*, **1998**, 459; (d) Diana G D, Cutchiffe D, Volkots D L, Mallamo J P, Vescio N, Oglesby R C, Nitz T J, Wetzel J, Girand V, Pevear D C & Dutko F J, *J Med Chem*, *36*, **1993**, 3240; (e) Khanna I K, Weier R M, Yu Y, Xu X D, Koazyk F J, Collins PW, Koboldt C M, Veenhuizen A W, Perkins W E, Caster J J, Maslerrer J L, Zhang Y Y, Gregory S A, Seibert K & Isakson P C, *J Med Chem*, *40*, **1997**, 1634; (f) Medwid J B, Paul R, Baker J S, Brockman J A, Du M T, Hallett W A, Hanifin J W, Hardy R A,Tarrent M E, Torley I W & Wrenn S, *J Med Chem*, *33*, **1990**, 1230; (g) Judkins B D, Allen D G, Cook T A, Evans B & Sardharwala T E, *Synth Commun*, *26*, **1996**, 4351; (h) March J, Advanced Organic Chemistry, John Wiley & Sons, New York, **1992**, pp. 918-919 and literatures cited therein.

3 (a) March J, Advanced Organic Chemistry, John Wiley & Sons: New York, **1992**, pp. 1038-1039 and literatures cited therein; (b) Meshram H M, *Synthesis*, **1992**, 943; (c) Katritzky A R, Zhanmg G F & Fan W Q, *Org Prep Proc Int*, *25*, **1993**, 315.

4 (a) Olah G A, & Keumi T, *Synthesis* **1979**, 112 and references cited therein; (b) Saednya A, *Synthesis*, **1982**, 190; (c) Bose D S, & Narsaiah A V, *Tetrahedron Lett*, *39*, **1998**, 6533; (d) Kumar H M S, Reddy B V S, Reddy P T & Yadav J S, *Synthesis*, **1999**, 586.

5 Brackman W, & Smit P J, *Recl Trav Chim*, *82*, **1963**, 757.

6 Parameswaran K N & Friedman O M, *Chem Ind*, **1965**, 988.

7 (a) Misono A, Osa T & Koda S, *Bull Chem Soc Jpn*, *40*, **1967**, 287;. (b) Talukdar S, Hsu J-L, Chou T-C & Fang J-M, *Tetrahedron Lett*, *42,* **2001**, 1103.

8 Erman M B, Snow J W & Williams M J, *Tetrahedron Lett*, *41*, **2000**, 6749.

9 (a) Singh R M, Srivastava A & Chandra A, *Indian J Chem, 44B*, **2005**, 2077; (b) Singh R M, Srivastava A & Singh M K, *Indian J Chem, 45B*, **2006**, 292; (c) Singh R M, Srivastava A & Chandra A, *Indian J Chem, 46B*, **2007**, 303; (d) Singh R M, Chandra A, Singh M K, *Synth Commun*, *37*, **2007**, 1689; (e) Singh R M, Singh M K, Chandra A & Singh B, *Tetrahedron Lett*, *48*, **2007**, 5987; (f) Singh R M & Srivastava A, *Indian J Chem, 44B*, **2005**, 1868.