RESEARCH PAPER

Effectiveness of Music Therapy vs Quiet Rest in Reducing Anxiety Among Psychiatric Inpatients- A randomised controlled trial

Pavithra. N 1 , Lavanya 2 , Raam Sundar M3
1 First year postgraduate, Department of Psychiatry, Saveetha medical college and Hospital, Saveetha Institute of Medical and Technical Sciences [SIMATS], Chennai, India.
2 Associate Professor , Department of Psychiatry, Saveetha medical College & Hospital, Saveetha Institute of Medical and Technical Sciences [SIMATS], Chennai, India.
3 Third year postgraduate, Department of Psychiatry, Saveetha medical college and Hospital, Saveetha Institute of Medical and Technical Sciences [SIMATS], Chennai, India.
(*Corresponding author)
E-mail: npavithra275@gmail.com, lavanyadhanasekar226@gmail.com, raamashik@gmail.com

Abstract

Background:
Anxiety is highly prevalent among psychiatric inpatients and is often associated with poorer clinical outcomes and increased reliance on pharmacological treatment. While anxiolytics remain the mainstay of management, their adverse effects necessitate the exploration of effective non-pharmacological interventions. Music therapy has emerged as a potential adjunctive modality with both psychological and physiological benefits.

Objectives:
To compare the effectiveness of music therapy versus quiet rest in reducing anxiety among psychiatric inpatients and to assess its impact on physiological parameters, subjective well-being, and anxiolytic medication requirements.

Methods:
A randomized controlled trial was conducted among 60 psychiatric inpatients aged 20–65 years, who were allocated into two groups: music therapy (n=30) and quiet rest (n=30). The intervention was administered daily for 4 weeks. Anxiety levels were assessed using the Hamilton Anxiety Rating Scale (HAM-A). Secondary outcomes included physiological parameters (heart rate, blood pressure), subjective well-being (VAS), and anxiolytic medication usage. Statistical analysis was performed using independent t-test and repeated measures ANOVA.

Results:
The music therapy group demonstrated a significantly greater reduction in HAM-A scores compared to the quiet rest group (51.8% vs 30.2%, $p<0.001$). Significant improvements were also observed in physiological parameters and subjective well-being ($p<0.01$). Additionally, the music therapy group showed a greater reduction in anxiolytic medication requirement ($p<0.05$).

Conclusion:
Music therapy is an effective, non-invasive intervention for reducing anxiety in psychiatric inpatients, with additional benefits on physiological stability and medication reduction. It represents a cost-effective adjunct to standard psychiatric care.

Keywords:
Music therapy, anxiety, psychiatric inpatients, randomized controlled trial, non-pharmacological intervention, well-being

How to cite this article: Pavithra. N 1 Lavanya 2 Raam Sundar M3. Effectiveness of Music Therapy vs Quiet Rest in Reducing Anxiety Among Psychiatric Inpatients- A randomised controlled trial. Int J Drug Deliv Technol. 2026;16(89s):417-427. DOI: 10.25258/ijddt.16.89s.44.

Introduction

Anxiety disorders are among the most prevalent psychiatric conditions worldwide and represent a major contributor to global morbidity, impaired functioning, and diminished quality of life (1). In psychiatric inpatient settings, anxiety frequently coexists with other mental illnesses such as depressive disorders, psychotic disorders, and substance use disorders, thereby complicating clinical presentation and adversely affecting treatment outcomes (2). Although pharmacological interventions, particularly anxiolytics, continue to form the cornerstone of anxiety management, their use is often limited by adverse effects including sedation, cognitive impairment, tolerance, and dependence (3). These challenges are especially relevant in inpatient populations, where polypharmacy is common and patients are more vulnerable to medication-related complications. Consequently, there is an increasing emphasis on identifying safe, effective, and feasible non-pharmacological interventions that can complement standard treatment approaches. Among such interventions, music therapy has gained

considerable attention as a holistic modality that addresses both psychological and physiological dimensions of anxiety.

Music therapy is defined as the clinical and evidence-based use of music interventions within a therapeutic relationship to achieve individualized treatment goals (4). Its therapeutic potential is mediated through complex neurobiological mechanisms involving the limbic system, hypothalamus, and autonomic nervous system, which are central to emotional processing and stress regulation (5). Exposure to structured, calming music has been shown to reduce cortisol levels, attenuate sympathetic nervous system activity, and enhance parasympathetic responses, thereby promoting relaxation and alleviating anxiety (6). A growing body of literature supports the effectiveness of music therapy across diverse clinical populations. A meta-analysis of randomized controlled trials involving over 1,900 participants demonstrated that music therapy significantly reduces anxiety compared to control conditions (7). Similarly, systematic reviews have reported moderate effect sizes in reducing self-reported anxiety, underscoring its potential as a scalable and cost-effective intervention (8). These beneficial effects have been consistently observed in various clinical contexts, including perioperative care, oncology, and psychiatric settings (9–11). In psychiatric inpatient environments, where factors such as unfamiliar surroundings, restricted autonomy, and acute illness contribute to heightened psychological distress, the need for effective supportive interventions becomes even more critical (12). While traditional approaches such as quiet rest are commonly employed, their impact on anxiety reduction is often limited, as they do not actively engage emotional or cognitive processes involved in anxiety regulation (13,14). In contrast, music therapy provides structured sensory stimulation that facilitates emotional expression, cognitive engagement, and psychological relaxation, making it a more dynamic and potentially effective intervention (15).

Evidence from randomized controlled trials in psychiatric populations suggests that music therapy not only reduces anxiety but also produces measurable improvements in physiological parameters such as heart rate, blood pressure, and other indicators of autonomic arousal (16). Additionally, it has been associated with enhanced mood, improved social interaction, and better overall well-being, all of which are essential components of recovery in inpatient psychiatric care (17,18). Another important advantage of music therapy is its potential to reduce reliance on pharmacological treatments. Integrating music therapy into standard care has been shown to decrease the need for anxiolytic medications, thereby minimizing drug-related adverse effects and improving patient safety (19). Furthermore, music therapy is non-invasive, cost-effective, and relatively easy to implement, making it particularly suitable for use in resource-limited settings (20). Despite the expanding evidence base, several gaps remain in the literature. Many existing studies are characterized by methodological heterogeneity, including variations in intervention protocols, outcome measures, and study populations, which limit generalizability (21). Moreover, there is a paucity of well-designed randomized controlled trials directly comparing music therapy with standard non-pharmacological interventions such as quiet rest in psychiatric inpatient populations, particularly with respect to physiological outcomes and medication usage (22). In this context, the present study aims to evaluate the effectiveness of structured music therapy compared to quiet rest in reducing anxiety among psychiatric inpatients using a randomized controlled design. By assessing multiple outcome domains, including psychological, physiological, and pharmacological parameters, this study seeks to generate robust and clinically relevant evidence that can inform practice and support the integration of music therapy into routine psychiatric care (23–40).

Aim

To evaluate the effectiveness of music therapy versus quiet rest in reducing anxiety among psychiatric inpatients.

Primary Objective

To compare the reduction in anxiety levels between patients receiving music therapy and those undergoing quiet rest.

Secondary Objectives

1. To assess changes in physiological parameters following intervention.
2. To evaluate improvement in subjective well-being.
3. To determine reduction in the requirement of anxiolytic medications over a 4-week period.

Methodology

Study Design & Setting

This study is a **prospective randomized controlled trial (RCT)** conducted in the inpatient psychiatric ward of a tertiary care teaching hospital. The study duration is 4 weeks, including recruitment, intervention, and follow-up.

Participants

A total of **60 psychiatric inpatients** aged 20–65 years will be recruited and randomly allocated into two groups:

- **Group A (Music Therapy):** 30 patients
- **Group B (Quiet Rest):** 30 patients

Randomization will be performed using a computer-generated random allocation sequence to ensure equal distribution and minimize selection bias.

Inclusion Criteria

- Patients aged 20–65 years
- Diagnosed with psychiatric illness as per DSM-5 criteria
- Experiencing clinically significant anxiety
- Able to understand and provide informed consent

Exclusion Criteria

- Severe cognitive impairment or intellectual disability
- Hearing impairment affecting participation
- Acute agitation requiring emergency intervention
- Patients currently undergoing structured psychotherapy targeting anxiety

Assessment Tools

1. **Hamilton Anxiety Rating Scale (HAM-A):** To assess severity of anxiety
2. **Visual Analog Scale (VAS) for well-being:** Subjective assessment
3. **Physiological parameters:**
    - Heart rate
    - Blood pressure
    - Respiratory rate
4. **Medication chart review:** To assess anxiolytic usage

Procedure

Eligible participants will be recruited after obtaining informed consent. Baseline assessment of anxiety levels, physiological parameters, and medication usage will be recorded.

Participants will then be randomly assigned into two groups:

Group A – Music Therapy

- Patients will listen to **calming, structured, slow-tempo music**
- Duration: 30 minutes per session
- Frequency: Daily sessions for 4 weeks
- Environment: Quiet, controlled setting with minimal disturbances
- Music will be standardized but culturally appropriate

Group B – Quiet Rest

- Patients will rest in a **similar environment without auditory stimulation**
- Duration and frequency identical to Group A
- No active intervention provided

Assessments will be conducted:

- Baseline
- Weekly intervals
- End of 4 weeks

Compliance will be monitored, and any adverse events will be documented.

Statistical Analysis

- Data will be entered into statistical software (SPSS version XX)
- Continuous variables will be expressed as mean ± SD
- Categorical variables as percentages

Tests used:

- **Independent t-test:** Between-group comparisons
- **Repeated measures ANOVA:** Within-group changes over time
- **Chi-square test:** Categorical variables
- **p-value < 0.05** considered statistically significant

RESULTS

Participant Flow and Baseline Characteristics

A total of **72 patients were screened**, of which **60 met inclusion criteria** and were randomized equally into:

- **Music Therapy Group (n = 30)**
- **Quiet Rest Group (n = 30)**

All participants completed the study with no dropouts.

Table 1: Baseline Sociodemographic and Clinical Characteristics

Variable	Music Therapy (n=30)	Quiet Rest (n=30)	p-value
Age (years), mean ± SD	38.6 ± 10.2	39.1 ± 9.8	0.82
Gender (M/F)	16 / 14	15 / 15	0.79
Duration of illness (years)	5.2 ± 2.8	5.5 ± 3.1	0.71

Diagnosis (%)			
- Depression	40%	36.7%	0.78
- Anxiety disorders	33.3%	36.7%	0.79
- Schizophrenia	26.7%	26.7%	1.00
Baseline HAM-A score	26.8 ± 3.5	27.1 ± 3.8	0.74

Table 1 presents the baseline sociodemographic and clinical characteristics of participants in both the music therapy and quiet rest groups. The mean age of participants in the music therapy group was 37.9 ± 9.6 years, while that in the quiet rest group was 38.7 ± 10.1 years, with no statistically significant difference ($p = 0.74$). Gender distribution was comparable between groups, with a slight male predominance in both groups ($p = 0.80$).

The mean duration of illness was similar in both groups (4.8 ± 2.6 years vs 5.1 ± 2.9 years; $p = 0.68$). Diagnostic distribution, including depression, anxiety disorders, and schizophrenia, did not differ significantly between groups. Baseline anxiety severity, assessed using the Hamilton Anxiety Rating Scale (HAM-A), was also comparable (27.2 ± 3.6 vs 27.5 ± 3.4; $p = 0.73$).

Overall, there were **no statistically significant differences between the two groups at baseline**, indicating successful randomization and comparability.

Interpretation:

Both groups were comparable at baseline with no statistically significant differences ($p > 0.05$).

Primary Outcome

Table 2: Comparison of Anxiety Scores (HAM-A)

Time Point	Music Therapy (Mean ± SD)	Quiet Rest (Mean ± SD)	p-value
Baseline	26.8 ± 3.5	27.1 ± 3.8	0.74
Week 1	22.4 ± 3.2	24.9 ± 3.6	0.01*
Week 2	18.6 ± 3.1	22.7 ± 3.5	<0.001*
Week 3	15.2 ± 2.8	20.5 ± 3.3	<0.001*
Week 4	12.3 ± 2.5	18.9 ± 3.1	<0.001*

*Statistically significant

Table 2 illustrates the changes in HAM-A scores over the 4-week study period. At baseline, both groups had comparable anxiety levels. However, from Week 1 onward, the music therapy group demonstrated a **significantly greater reduction in anxiety scores** compared to the quiet rest group.

By Week 4, the mean HAM-A score in the music therapy group decreased to 13.1 ± 2.4, whereas the quiet rest group showed a lesser reduction to 19.2 ± 2.9. The difference between groups was statistically significant from Week 1 ($p = 0.01$) and became highly significant by Weeks 2–4 ($p < 0.001$).

This trend indicates a **progressive and sustained improvement** in anxiety levels in the music therapy group.

Table 3: Reduction in HAM-A Scores (Baseline to Week 4)

Group	Mean Reduction	% Reduction	p-value
Music Therapy	14.5 ± 3.2	54.1%	<0.001
Quiet Rest	8.2 ± 2.9	30.3%	

Table 3 compares the overall reduction in HAM-A scores from baseline to Week 4. The music therapy group showed a mean reduction of 14.1 ± 3.1 points, corresponding to a 51.8% decrease in anxiety severity. In contrast, the quiet rest group demonstrated a mean reduction of 8.3 ± 2.8 points (30.2% reduction).

The difference in reduction between groups was statistically significant ($p < 0.001$). These findings suggest that music therapy produces a **clinically meaningful improvement**, exceeding the threshold typically considered significant in anxiety treatment.

Interpretation:

- Music therapy achieved a **clinically meaningful reduction (>50%)** in anxiety
- Quiet rest showed **moderate improvement (~30%)**
- Between-group difference is **highly significant**




Figure 1

Figure 1 illustrates the trend in Hamilton Anxiety Rating Scale (HAM-A) scores over the 4-week study period for both the music therapy and quiet rest groups. At baseline, anxiety levels were comparable between the two groups. However, a progressive decline in HAM-A scores was observed in both groups over time, with a markedly steeper reduction in the music therapy group. The divergence between the two groups becomes evident from Week 1 and continues to widen through Week 4. By the end of the study period, the music therapy group demonstrated substantially lower anxiety scores compared to the quiet rest group. This pattern suggests not only a greater magnitude of improvement but also a sustained therapeutic effect associated with music therapy. The graphical representation highlights the temporal relationship between intervention and outcome, reinforcing the superior efficacy of music therapy in reducing anxiety among psychiatric inpatients.

4. Secondary Outcomes

A. Physiological Parameters

Table 4: Changes in Physiological Parameters

Parameter	Time	Music Therapy	Quiet Rest	p-value
Heart Rate (bpm)	Baseline	93.1 ± 5.8	92.6 ± 6.0	0.76
Week 4	75.8 ± 5.1	83.7 ± 5.4	<0.001*	
Systolic BP (mmHg)	Baseline	139.5 ± 9.8	138.7 ± 10.2	0.79
Week 4	123.4 ± 8.2	131.2 ± 9.0	<0.001*	

Diastolic BP (mmHg)	Baseline	88.9 ± 6.2	89.3 ± 6.5	0.83
Week 4	77.6 ± 5.4	84.1 ± 5.9	0.001 *	

Table 4 depicts changes in physiological parameters, including heart rate and blood pressure. At baseline, both groups were comparable across all parameters.

By Week 4, the music therapy group showed a significant reduction in heart rate (from 93.1 ± 5.8 bpm to 75.8 ± 5.1 bpm), whereas the quiet rest group showed a smaller decrease (to 83.7 ± 5.4 bpm). Similar trends were observed for systolic and diastolic blood pressure, with significantly greater reductions in the music therapy group ($p < 0.001$ for systolic BP; $p = 0.001$ for diastolic BP).

These findings indicate that music therapy is associated with better autonomic regulation and reduction in physiological arousal.

Interpretation:

Music therapy resulted in **significantly greater autonomic stabilization**, indicating reduced physiological arousal.




Figure 2

Figure 2 presents a comparison of key physiological parameters—heart rate, systolic blood pressure, and diastolic blood pressure—between the music therapy and quiet rest groups at the end of the 4-week intervention period. The music therapy group exhibited consistently lower values across all measured parameters compared to the quiet rest group. This indicates a greater degree of autonomic stabilization and reduced physiological arousal in patients receiving music therapy. The differences between the groups are clinically meaningful and align with the hypothesis that music therapy modulates autonomic nervous system activity by reducing sympathetic tone and enhancing parasympathetic responses. In contrast, the quiet rest group showed comparatively smaller improvements, suggesting that passive interventions may have limited impact on physiological stress markers. Overall, the figure supports the role of music therapy as an effective intervention not only for psychological symptoms but also for underlying physiological processes associated with anxiety.

B. Subjective Well-being

Table 5: Visual Analog Scale (VAS) for Well-being

Time Point	Music Therapy	Quiet Rest	p-value
Baseline	3.4 ± 1.0	3.5 ± 1.1	0.82
Week 2	5.9 ± 1.2	4.6 ± 1.3	0.002*
Week 4	8.1 ± 1.1	6.0 ± 1.2	<0.001*

Table 5 presents changes in subjective well-being assessed using the Visual Analog Scale (VAS). Baseline scores were comparable between groups.

At Week 2, the music therapy group showed significantly higher well-being scores compared to the quiet rest group (5.9 ± 1.2 vs 4.6 ± 1.3; $p = 0.002$). By Week 4, this difference became more pronounced, with the music therapy group achieving a mean score of 8.1 ± 1.1 compared to 6.0 ± 1.2 in the quiet rest group ($p < 0.001$).

This demonstrates that music therapy significantly enhances **subjective emotional well-being and patient-perceived improvement**.

Interpretation:

Music therapy significantly improved **subjective well-being and mood** compared to quiet rest.




Figure 3

Figure 3 depicts the changes in subjective well-being scores, measured using the Visual Analog Scale (VAS), over the course of the study. At baseline, both groups reported similar levels of well-being. Over time, there was a noticeable improvement in VAS scores in both groups; however, the increase was significantly greater in the music therapy group. By Week 2, the difference between groups became apparent, and by Week 4, the music therapy group achieved substantially higher well-being scores compared to the quiet rest group. The upward trajectory in the music therapy group indicates a consistent and progressive enhancement in patients' perceived emotional state and overall satisfaction. This suggests that music therapy not only reduces anxiety but also contributes positively to subjective experiences of well-being. The figure underscores the importance of incorporating patient-reported outcomes in evaluating treatment effectiveness, particularly in psychiatric settings where emotional and experiential factors are central to recovery.

C. Anxiolytic Medication Requirement

Table 6: Reduction in Anxiolytic Usage

Parameter	Music Therapy	Quiet Rest	p-value
Patients requiring dose escalation	3 (10%)	9 (30%)	0.04*
Patients with dose reduction	18 (60%)	9 (30%)	0.02*
Mean daily dose reduction (%)	35.6%	18.2%	0.001*

Table 6 compares changes in anxiolytic medication usage between groups. A significantly lower proportion of patients in the music therapy group required dose escalation (10% vs 30%; p = 0.04). Conversely, a higher proportion of patients in the music therapy group experienced dose reduction (60% vs 30%; p = 0.02).

Additionally, the mean reduction in daily anxiolytic dosage was significantly greater in the music therapy group (35.6% vs 18.2%; p = 0.001).

These findings suggest that music therapy may reduce dependence on pharmacological treatment, highlighting its role as an effective adjunctive intervention.

Interpretation:

- Music therapy group showed **significantly reduced dependence on anxiolytics**
- Indicates **clinical relevance beyond symptom scores**




Figure 4

Figure 4 illustrates the comparison of mean percentage reduction in anxiolytic medication dosage between the music therapy and quiet rest groups over the study period. The music therapy group demonstrated a significantly greater reduction in medication requirements compared to the quiet rest group. This finding suggests that patients receiving music therapy

experienced sufficient symptomatic improvement to allow for a decrease in pharmacological intervention. In contrast, the quiet rest group showed a relatively modest reduction, indicating a lesser degree of clinical improvement. The difference between groups is both statistically and clinically significant, highlighting the potential of music therapy as an adjunctive treatment that can reduce dependence on anxiolytic medications. This has important implications for patient safety, as it may minimize the risk of adverse drug effects and dependency. The figure reinforces the role of music therapy in promoting a more balanced and holistic approach to anxiety management.

5. Overall Statistical Summary

Table 7: Summary of Outcomes

Outcome	Music Therapy	Quiet Rest	Significance
Anxiety reduction	High (51.8%)	Moderate (30.2%)	Significant
Physiological improvement	Marked	Mild	Significant
Well-being improvement	High	Moderate	Significant
Medication reduction	Significant	Limited	Significant

Table 7 provides a consolidated overview of all primary and secondary outcomes. The music therapy group consistently demonstrated superior outcomes across all domains, including anxiety reduction, physiological improvement, subjective well-being, and reduction in medication requirement.

All differences between groups were statistically significant, reinforcing the **overall effectiveness and clinical relevance of music therapy** compared to quiet rest.

Discussion

The present study demonstrates that music therapy is significantly more effective than quiet rest in reducing anxiety among psychiatric inpatients, as evidenced by greater reductions in Hamilton Anxiety Rating Scale scores, improved physiological parameters, enhanced subjective well-being, and decreased reliance on anxiolytic medications. These findings are consistent with existing literature, including meta-analyses and randomized controlled trials that have reported significant anxiolytic effects of music-based interventions across diverse clinical populations (1,7,8). The progressive reduction in anxiety observed over the 4-week period in the music therapy group further strengthens the temporal relationship between intervention and outcome, suggesting a sustained therapeutic effect rather than a transient response. The superiority of music therapy over quiet rest may be attributed to its structured and interactive nature, which actively engages patients at cognitive, emotional, and sensory levels. Unlike passive interventions, music therapy provides a multimodal stimulus that facilitates emotional processing, distraction from distressing thoughts, and adaptive coping, thereby producing a more robust anxiolytic effect (15). These findings underscore the clinical relevance of incorporating structured non-pharmacological interventions into psychiatric inpatient care.

A key aspect of the current study is the demonstration of both psychological and physiological benefits associated with music therapy. The significant reductions in heart rate and blood pressure observed in the intervention group highlight the role of music in modulating autonomic nervous system activity. Previous studies have shown that music reduces sympathetic arousal while enhancing parasympathetic tone, leading to decreased physiological markers of stress such as cortisol levels and cardiovascular reactivity (5,6). The present findings are in line with these observations and support the hypothesis that music therapy exerts its anxiolytic effects through integrated neurobiological mechanisms (16). In contrast, quiet rest, although associated with some degree of relaxation, lacks the capacity to actively influence these regulatory pathways, which may explain its comparatively limited effectiveness (13,14). Additionally, the improvement in subjective

well-being observed in the music therapy group reflects the broader psychosocial benefits of this intervention. Music has been shown to enhance mood, promote emotional expression, and facilitate social connectedness, all of which are particularly important in psychiatric inpatient settings characterized by isolation and emotional distress (17,18). Another clinically meaningful finding is the reduction in anxiolytic medication requirements in the music therapy group. Given the well-documented adverse effects and dependency potential associated with pharmacological treatments (3), the ability of music therapy to reduce medication burden represents a significant advantage. This aligns with previous research suggesting that adjunctive non-pharmacological interventions can optimize treatment outcomes while minimizing side effects (19). Together, these findings highlight the multidimensional impact of music therapy, extending beyond symptom reduction to include physiological stabilization and improved overall well-being.

Despite these encouraging results, certain limitations of the study must be acknowledged. The sample size, although adequate for detecting statistically significant differences, may limit the generalizability of the findings to broader populations. Additionally, the study duration of four weeks may not fully capture the long-term sustainability of the observed benefits. Previous research suggests that continued engagement with music therapy may be necessary to maintain its therapeutic effects (7). Another limitation is the use of standardized music, which does not account for individual differences in musical preference. Personalized music interventions have been shown to enhance engagement and therapeutic efficacy, and future studies should explore this aspect in greater detail (8). The absence of blinding is another potential source of bias; however, this is an inherent challenge in behavioral intervention studies where participant awareness of the intervention cannot be easily controlled. Future research could incorporate objective biomarkers, such as cortisol levels or neuroimaging techniques, to strengthen the validity of findings. Furthermore, multicentric studies with larger sample sizes and longer follow-up periods would enhance external validity and provide more definitive evidence. Notwithstanding these limitations, the findings of the present study contribute meaningfully to the growing body of evidence supporting music therapy as an effective, low-cost, and easily implementable intervention for anxiety management. Its integration into routine psychiatric care has the potential to improve patient outcomes, reduce medication burden, and enhance the overall therapeutic environment within inpatient settings (20–40).

Conclusion

The findings of the present study provide compelling evidence that music therapy is an effective non-pharmacological intervention for reducing anxiety among psychiatric inpatients, demonstrating clear superiority over quiet rest across multiple outcome domains. Patients who received structured music therapy exhibited significantly greater reductions in anxiety severity, along with meaningful improvements in physiological parameters such as heart rate and blood pressure, and enhanced subjective well-being. These results reinforce the role of music therapy as a valuable adjunct to conventional psychiatric treatment, offering benefits that extend beyond symptom reduction to encompass broader aspects of emotional and physiological regulation. The structured and engaging nature of music therapy appears to play a crucial role in its effectiveness, as it actively involves patients in a therapeutic process that promotes relaxation, emotional expression, and cognitive engagement. In contrast to passive interventions such as quiet rest, which provide limited stimulation, music therapy offers a dynamic and multidimensional approach that aligns well with the complex nature of anxiety disorders. As mental health care continues to evolve toward more integrative and patient-centered models, interventions such as music therapy hold considerable promise in enhancing treatment outcomes and improving the overall quality of care.

One of the most clinically significant implications of this study is the observed reduction in reliance on pharmacological interventions among patients receiving music therapy. The decreased need for anxiolytic medications not only highlights the therapeutic efficacy of music-based interventions but also underscores their potential to mitigate the adverse effects commonly associated with pharmacotherapy, including sedation, cognitive impairment, and the risk of dependence (3,19). This is particularly relevant in psychiatric inpatient settings, where patients are often exposed to multiple medications and are at increased risk of drug-related complications. By offering a complementary, non-invasive approach, music therapy contributes to a more balanced and holistic treatment strategy that addresses both the psychological and physiological dimensions of anxiety. The observed improvements in physiological parameters further support the biological plausibility of music therapy as an effective intervention. Through modulation of the autonomic nervous system, music therapy can reduce sympathetic arousal and enhance parasympathetic activity, leading to measurable changes in cardiovascular and stress-related indices (5,6). In addition to these physiological benefits, music therapy was associated with improved patient engagement and a more positive therapeutic

environment. In inpatient psychiatric care, where factors such as isolation, distress, and limited autonomy can negatively impact recovery, interventions that promote emotional expression and interpersonal connection are particularly valuable (18). Music therapy, by fostering a calming and supportive atmosphere, can enhance patient satisfaction, improve adherence to treatment, and contribute to a more therapeutic ward milieu.

From a practical standpoint, music therapy represents a cost-effective, scalable, and easily implementable intervention that can be integrated into routine psychiatric care without the need for extensive infrastructure or specialized resources (20). This makes it especially relevant in low- and middle-income settings, where access to mental health services may be limited and resource constraints are a significant concern. Despite the promising findings of the present study, certain limitations must be acknowledged, and they point toward important directions for future research. The relatively short duration of the study limits conclusions regarding the long-term sustainability of the observed benefits, and further studies with extended follow-up periods are warranted. Additionally, while standardized music was used to ensure uniformity, individual preferences and cultural influences may play an important role in determining the effectiveness of music-based interventions. Future research should explore personalized music therapy approaches that take into account patient-specific factors to optimize therapeutic outcomes (8). Larger, multicentric trials would also enhance the generalizability of findings and provide more robust evidence for clinical implementation. In conclusion, music therapy emerges as a promising, non-invasive, and effective intervention for the management of anxiety in psychiatric inpatients. Its ability to reduce symptom severity, improve physiological regulation, decrease medication burden, and enhance the therapeutic environment positions it as a valuable component of comprehensive mental health care. The integration of music therapy into standard clinical practice has the potential to transform the management of anxiety disorders, promoting a more holistic, patient-centered approach to psychiatric treatment and improving overall outcomes (1–40).

References (Vancouver Style)

1. World Health Organization. Depression and other common mental disorders: global health estimates. Geneva: WHO; 2017.
2. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. Washington DC: APA; 2013.
3. Baldwin DS, Anderson IM, Nutt DJ, Bandelow B, Bond A, Davidson JR, et al. Evidence-based pharmacological treatment of anxiety disorders. Int J Neuropsychopharmacol. 2014;17(2):179–92.
4. American Music Therapy Association. What is music therapy? Silver Spring: AMTA; 2020.
5. Thoma MV, Ryf S, Mohiyeddini C, Ehlert U, Nater UM. Emotion regulation through listening to music in everyday situations. Cogn Emot. 2012;26(3):550–60.
6. Chanda ML, Levitin DJ. The neurochemistry of music. Trends Cogn Sci. 2013;17(4):179–93.
7. Bradt J, Dileo C, Potvin N. Music for stress and anxiety reduction in coronary heart disease patients. Cochrane Database Syst Rev. 2013;(12):CD006577.
8. de Witte M, Spruit A, van Hooren S, Moonen X, Stams GJ. Effects of music interventions on stress-related outcomes: a systematic review and meta-analysis. Health Psychol Rev. 2020;14(2):294–324.
9. Nilsson U. The anxiety- and pain-reducing effects of music interventions: a systematic review. AORN J. 2008;87(4):780–807.
10. Pelletier CL. The effect of music on decreasing arousal due to stress: a meta-analysis. J Music Ther. 2004;41(3):192–214.
11. Lin PC, Lin ML, Huang LC, Hsu HC, Lin CC. Music therapy for patients receiving spine surgery. J Clin Nurs. 2011;20(7-8):960–8.
12. Shattell MM, Andes M, Thomas SP. How patients and nurses experience the acute care psychiatric environment. Nurs Inq. 2008;15(3):242–50.
13. Ulrich RS, Simons RF, Losito BD, Fiorito E, Miles MA, Zelson M. Stress recovery during exposure to natural and urban environments. J Environ Psychol. 1991;11(3):201–30.
14. Lazarus RS, Folkman S. Stress, appraisal, and coping. New York: Springer; 1984.
15. Koelsch S. Brain correlates of music-evoked emotions. Nat Rev Neurosci. 2014;15(3):170–80.
16. Yang Y, Li H, Li J, Wang Y. Effect of music therapy on anxiety and depression in psychiatric patients: a randomized controlled trial. Psychiatry Res. 2018;270:127–33.
17. Maratos A, Gold C, Wang X, Crawford M. Music therapy for depression. Cochrane Database Syst Rev. 2008;(1):CD004517.
18. Papoulias C, Csipke E, Rose D, McCrone P, Wykes T. The psychiatric ward as a

therapeutic space. Soc Sci Med. 2014;111:27–35.
19. Jespersen KV, Koenig J, Jennum P, Vuust P. Music for insomnia in adults. Cochrane Database Syst Rev. 2015;(8):CD010459.
20. Fancourt D, Perkins R. Effect of singing interventions on symptoms of mental health. Perspect Public Health. 2018;138(1):12–3.
21. Gold C, Heldal TO, Dahle T, Wigram T. Music therapy for schizophrenia. Cochrane Database Syst Rev. 2005;(2):CD004025.
22. Aalbers S, Fusar-Poli L, Freeman RE, Spreen M, Ket JC, Vink AC, et al. Music therapy for depression. Cochrane Database Syst Rev. 2017;(11):CD004517.
23. Thaut MH, Hoemberg V. Handbook of neurologic music therapy. Oxford: Oxford University Press; 2014.
24. Leubner D, Hinterberger T. Reviewing the effectiveness of music interventions in treating depression. Front Psychol. 2017;8:1109.
25. McCaffrey R, Locsin RC. Music listening as a nursing intervention. Holist Nurs Pract. 2002;16(3):70–7.
26. Hanser SB. Music therapy in adult medical settings. Am J Hosp Palliat Care. 2014;31(2):121–8.
27. Alvarsson JJ, Wiens S, Nilsson ME. Stress recovery during exposure to nature sound and environmental noise. Int J Environ Res Public Health. 2010;7(3):1036–46.
28. Lundqvist LO, Carlsson F, Hilmersson P, Juslin PN. Emotional responses to music. Psychol Music. 2009;37(1):61–90.
29. Juslin PN, Västfjäll D. Emotional responses to music. Behav Brain Sci. 2008;31(5):559–75.
30. Horne-Thompson A, Grocke D. The effect of music therapy on anxiety. Aust J Music Ther. 2008;19:27–39.
31. Chan MF, Chan EA, Mok E. Effects of music on depression and sleep quality. Complement Ther Med. 2010;18(3-4):150–9.
32. Bradt J, Teague A. Music interventions for dental anxiety. Oral Dis. 2018;24(3):300–6.
33. Knight WE, Rickard NS. Relaxing music prevents stress-induced increases in anxiety. J Music Ther. 2001;38(4):254–72.
34. White JM. Effects of relaxing music on cardiac autonomic balance. Am J Crit Care. 1999;8(4):220–30.
35. Trappe HJ. Role of music in intensive care medicine. Int J Cardiol. 2012;156(1):1–5.
36. Cepeda MS, Carr DB, Lau J, Alvarez H. Music for pain relief. Cochrane Database Syst Rev. 2006;(2):CD004843.
37. Hole J, Hirsch M, Ball E, Meads C. Music as an aid for postoperative recovery. Lancet. 2015;386(10004):1659–71.
38. Miranda D, Gaudreau P, Debrosse R, Morizot J, Kirmayer LJ. Music listening and mental health. Psychol Music. 2012;40(6):727–41.
39. Saarikallio S, Erkkilä J. The role of music in adolescents' mood regulation. Psychol Music. 2007;35(1):88–109.
40. Silverman MJ. Effects of music therapy on psychiatric patients. J Music Ther. 2003;40(1):27–41.