RESEARCH PAPER

Integrating AI & Digital Technology into Community Health Workers Improving Mental Health of Society & Maternal-Pediatric Care

B. Uvarani[1], Lego Shincy P. R[2]*, Rajitha Stanislaus Rajam[3], Reena Mariadhas Rajam[4], Harpreet Kaur[5], Neha Thakur[6], Amit Chawla[7], Payal Rani[8]

[1]Professor, MNR College of Nursing, Fasalwadi, Sangareddy, Telangana (balugoodenough@gmail.com)

[2]*Associate Professor cum HOD, Department of Community Health Nursing, St. Marys College of Nursing Lucknow, Uttar Pradesh (legojesuspr@gmail.com) (*Corresponding author)

[3]Associate Professor in OBG Nursing, Christian College of Nursing, Neyyoor, Kanyakumari. Affiliated to the Tamil Nadu Dr. MGR Medical University, Chennai, Tamil Nadu, India

[4]Professor, Department of Midwifery, KAAF University, Ghana (reenaelwin@gmail.com, prof_reenaelwin@kaafuni.edu.gh)

[5]Principal, School of Health Sciences (Department of Nursing), RIMT University, Mandi Gobindgarh, Punjab

[6]Assistant Professor, Department of Community Health Nursing, SGRRIM&HS College of Nursing, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India.

[7]Professor cum Principal, Faculty of Pharmacy, Tantia University, Sri Ganganagar, Rajasthan, India (amitchawla84@gmail.com) https://orcid.org/0009-0001-0622-0712

[8]Professor cum Principal, Mata Jarnail Kaur Memorial College of Pharmacy (Under Desh Bhagat University), Sri Muktsar Sahib, Punjab (palamitchawla@gmail.com) https://orcid.org/0009-0009-8189-5953

Abstract

Community Health Workers (CHWs) serve as the vital linchpin of primary healthcare delivery, particularly in low- and middle-income countries (LMICs) and medically underserved rural or urban enclaves within high-income nations. The rapid maturation of artificial intelligence (AI), machine learning (ML), mobile health (mHealth) architectures, point-of-care diagnostics, and digital decision-support tools offers an unprecedented opportunity to augment the clinical scope, operational efficiency, and diagnostic accuracy of CHW-led interventions. This article presents an extensive, multi-layered examination of the integration of AI and digital technologies into CHW workflows, focusing specifically on two high-priority public health domains: societal mental health and maternal-pediatric care.

We analyze how predictive analytics, conversational agents, natural language processing (NLP), computer vision, acoustic voice biomarkers, and automated triage networks empower CHWs to bridge longstanding diagnostic and therapeutic gaps. Furthermore, we investigate the systemic, structural, and ethical barriers to sustainable implementation, including algorithmic bias, edge-computing infrastructure, digital literacy, automation bias, data sovereignty, and human resource policy. By synthesizing empirical evidence across epidemiological, clinical, and technological domains, this paper articulates a strategic roadmap for scaling ethical, human-in-the-loop AI-enabled CHW programs to achieve sustainable, equitable healthcare delivery globally.

1. **How to cite this article:** B. Uvarani1 Lego Shincy P. R2* Rajitha Stanislaus Rajam3 Reena Mariadhas Rajam4 Harpreet Kaur5 Neha Thakur6 Amit Chawla7 Payal Rani8. Integrating AI & Digital Technology into Community Health Workers Improving Mental Health of Society & Maternal-Pediatric Care. Int J Drug Deliv Technol. 2026;16(90s):1702-1709. DOI: 10.25258/ijddt.16.90s.171.

Introduction

Globally, contemporary healthcare systems are constrained by severe workforce shortages, shifting demographic burdens, rising rates of chronic and non-communicable conditions, and persistent structural health inequities. The World Health Organization (WHO) projects a multi-million health worker deficit, a shortage felt most acutely in low- and middle-income countries (LMICs) where the ratio of skilled medical professionals (physicians, psychiatrists, obstetricians, and pediatricians) to the general population remains critically depressed. In response to these persistent structural gaps, Community Health Workers (CHWs)—trusted, local individuals trained to deliver task-shifted primary care, health education, preventative screening, and social support—have emerged as indispensable agents of frontline healthcare delivery (Perry et al., 2014).

CHWs operate directly at the intersection of public health systems and local communities. Their proximity to patients allows them to address social determinants of health, track longitudinal outcomes, and deliver culturally congruent care. Decades of empirical evidence demonstrate that CHW programs substantially improve immunization coverage, reduce neonatal and child mortality, enhance tuberculosis treatment adherence, and expand access to basic maternal health education.

However, traditional CHW delivery models face severe operational bottlenecks. CHWs frequently navigate extensive geographic catchment areas with high patient-to-worker ratios, equipped with minimal formal clinical training, static paper-based protocols, fragmented data collection methods, and absent or delayed expert supervision. Consequently, complex diagnostic and therapeutic tasks—such as detecting subtle early manifestations of perinatal


psychiatric distress, identifying pediatric neurodevelopmental delays, accurately calculating maternal blood loss, or triaging severe respiratory distress—frequently exceed traditional CHW capacity.

The rapid convergence of artificial intelligence (AI), cloud and edge computing, mobile telecommunications, bio-sensors, and point-of-care digital diagnostics introduces a paradigm shift. When embedded within accessible mobile health (mHealth) software applications, AI algorithms transform standard smartphones and tablets into sophisticated Clinical Decision-Support Systems (CDSS). For CHWs, these technology stacks function not as human replacements, but as "force multipliers" and "cognitive amplifiers." They standardize protocol adherence, provide real-time risk stratification, automate administrative documentation, and establish continuous, closed-loop communication channels with specialized medical networks (Labrique et al., 2018).

This paper provides an extended, comprehensive synthesis of the integration of AI and digital health technologies into CHW frameworks. Focusing specifically on the dual imperatives of **societal mental health** and **maternal-pediatric healthcare**, we examine technological architectures, clinical application vectors, empirical evidence, ethical challenges, operational trade-offs, and health policy frameworks required to achieve population-scale impact.

2. Theoretical Framework: The Digital Human-AI Hybrid Model in Community Health

To evaluate the integration of advanced technologies into frontline health workflows, implementation science must move beyond the binary view of AI as either a total automation tool or a superficial digital checklist. We propose a comprehensive **Digital Human-AI Hybrid Model** rooted in task-shifting frameworks, human-centered design principles, and sociotechnical systems theory (Agarwal et al., 2015).



2.1 Task-Shifting to Task-Amplification

Historically, task-shifting has involved delegating specific clinical tasks from highly trained, scarce professionals (such as doctors or clinical psychologists) to mid-level cadres or CHWs with shorter, focused training periods. While effective, traditional task-shifting is limited by the cognitive load placed on non-specialist workers when navigating complex clinical edge cases.

AI-driven task-amplification redefines this paradigm. By leveraging machine learning models trained on vast clinical datasets, the software assumes the high-dimensional pattern-recognition tasks—such as evaluating complex multi-variable clinical signs, predicting disease progression vectors, or analyzing acoustic and visual biomarkers. This offloading of cognitive processing enables CHWs to execute high-level triage, deliver evidence-based psychological micro-interventions, and perform advanced diagnostic routines under automated, real-time algorithmic supervision (Feroz et al., 2021).

2.2 System Architecture: The Triad of Frontline Digital Care


The functional architecture of an AI-enabled CHW ecosystem rests on three interconnected functional layers:

1. **Multimodal Data Acquisition Layer:** This layer captures raw clinical and demographic inputs at the point of care. Inputs include non-structured clinical notes, voice recordings, physiological data from Bluetooth-connected point-of-care sensors (e.g., pulse oximeters, blood pressure cuffs), and high-resolution images or videos captured via standard smartphone cameras.
2. **Algorithmic Analytics & Decision Layer:** Functioning either locally on the mobile device (edge AI) or in secure cloud environments, machine learning engines process raw inputs against clinical decision rules, deep neural networks, and natural language processing pipelines. This layer generates real-time outputs: diagnostic probability scores, risk stratification categories (e.g., low, moderate, severe), and customized action prompts for the CHW.
3. **Action, Escalation, and Governance Layer:** The CHW interprets the algorithmic output within the patient's socio-cultural context, validating or adjusting the recommendation. If high-risk thresholds are breached, the platform automatically triggers real-time escalation protocols—notifying regional supervisors, dispatching emergency transport services, or creating direct tele-consultation channels with medical specialists. All interactions log asynchronously into national Electronic Health Record (EHR) databases (e.g., DHIS2).

3. Revolutionizing Societal Mental Health via AI-Augmented CHWs

Mental, neurological, and substance-use disorders constitute a leading cause of global disability and premature mortality. The treatment gap for mental

health is severe: in low- and middle-income countries, between 75% and 90% of individuals with severe mental health conditions receive no formal medical or psychological intervention (Patel et al., 2018). This crisis is exacerbated by a acute shortage of trained mental health professionals, pervasive social stigma, and geographical barriers to central psychiatric facilities.



3.1 Digital Screening and Early Biomarker Detection

Standard mental health screenings rely on self-administered or interviewer-administered paper questionnaires, such as the Patient Health Questionnaire (PHQ-9) for depression or the Generalized Anxiety Disorder scale (GAD-7). While useful, these instruments are subject to reporting bias, social desirability distortion, translation inaccuracies, and challenges in administration among low-literacy populations.

Integrating AI into mobile CHW workflows allows for passive, objective, and multimodal biomarker detection:

3.1.1 Acoustic Voice Biomarkers

Depressive disorders, PTSD, and severe anxiety alter neuromotor control of the vocal apparatus, manifesting as subtle changes in speech mechanics. Mobile algorithms analyze speech recorded during routine CHW interactions, measuring acoustic dimensions such as pitch variability, formant frequencies, jitter, shimmer, speech cadence, and latency to response (Thies et al., 2021). Deep learning classifiers evaluate these features to identify markers of depressive or anxious states before clinical symptoms fully manifest, guiding the CHW to perform targeted clinical evaluations.

3.1.2 Natural Language Processing (NLP) & Semantic Analysis

Conversational AI tools embedded within CHW interfaces capture conversational dialogue in local dialects. Applied NLP algorithms analyze semantic structures for markers associated with mental health distress, such as affective valence, cognitive distortion patterns, absolute language usage, and explicit or implicit references to self-harm or hopelessness.

3.2 Task-Shifted Psychological Interventions

The World Health Organization's Mental Health Gap Action Programme (mhGAP) and structured interventions like the *Thinking Healthy Programme* (a cognitive-behavioral approach to perinatal depression) have shown that non-specialist health workers can effectively deliver psychological therapies when properly trained. AI technologies strengthen these task-shifted interventions in several key ways:


- **Real-Time Intra-Session Guidance:** During a home visit, an AI co-pilot active on the CHW's mobile device transcribes therapy sessions in real time, serving as a live clinical prompt. If a patient expresses somatic symptoms of panic or trauma, the interface suggests relevant, evidence-based coping exercises (e.g., grounding techniques, cognitive restructuring steps) for the CHW to guide.
- **Automated Supervisory Auditing & Quality Assurance:** In conventional settings, a single clinical psychologist might oversee hundreds of CHWs, creating supervisory bottlenecks. AI systems analyze session audio or interaction logs to measure protocol fidelity, evaluating whether the CHW accurately delivered core therapeutic components. Supervisory dashboards aggregate these fidelity metrics, automatically flagging cases where CHWs require target coaching (Naslund et al., 2019).

3.3 Longitudinal Risk Monitoring and Population Mental Health Analytics

Aggregating longitudinal screening metrics collected by CHWs across geographic regions enables municipal and national health authorities to build dynamic population mental health dashboards. Predictive analytics models identify community-level mental distress spikes linked to external stressors—such as economic disruptions, natural disasters, epidemics, or seasonal shifts. This real-time intelligence supports proactive resource allocation, targeted community outreach, and crisis team deployments (Torous et al., 2021).

4. Transforming Maternal and Pediatric Healthcare at the Grassroots

Preventable maternal, fetal, and child deaths remain concentrated in resource-constrained environments. Conditions such as pre-eclampsia, postpartum hemorrhage (PPH), fetal growth restriction, neonatal sepsis, severe acute malnutrition (SAM), and pediatric pneumonia require early detection and rapid intervention. AI-enhanced mHealth tools expand the diagnostic precision of CHWs, enabling early detection at the household level.



4.1 AI-Enhanced Antenatal and Postnatal Maternal Care

The continuum of maternal care requires early identification of obstetric risks during gestation, intrapartum management, and post-delivery follow-up.

4.1.1 Pre-Eclampsia Risk Prediction and Management

Hypertensive disorders of pregnancy, particularly pre-eclampsia and eclampsia, are major contributors to global maternal mortality. Standard detection requires accurate blood pressure measurement and clinical symptom tracking.

AI systems connected via Bluetooth to smart blood pressure cuffs allow CHWs to track subtle physiological trends across antenatal visits. Machine learning algorithms, trained on longitudinal maternal health datasets (such as the PIERS model), combine blood pressure trajectories, maternal age, gestational age, and reported symptoms (e.g., persistent headaches, visual disturbances) to generate individualized risk scores for pre-eclampsia (Von Dadelszen et al., 2020). High-risk classifications prompt immediate referrals to secondary care facilities for diagnostic confirmation and prophylactic interventions (such as low-dose aspirin or magnesium sulfate management).

4.1.2 Point-of-Care Obstetric Ultrasound

Traditional obstetric ultrasonography requires specialized sonographers and expensive capital equipment, making it largely inaccessible in rural primary care settings. The development of ultra-portable, hand-held ultrasound probes that connect directly to smartphones—coupled with deep learning image interpretation models—enables point-of-care imaging by non-physicians.

Equipped with these tools, CHWs can perform standardized sweeping motions across the pregnant abdomen. AI algorithms automatically process the live video feed to:

1. Estimate gestational age via automated biometrics (e.g., biparietal diameter, femur length).
2. Determine fetal presentation (e.g., cephalic vs. breech).
3. Identify placental location (flagging potential placenta previa).
4. Detect multiple gestations (Stroux et al., 2021).

4.1.3 Postpartum Hemorrhage (PPH) Quantification

Postpartum hemorrhage remains a leading cause of maternal mortality. Visual estimation of blood loss following delivery is notoriously inaccurate, often underestimating actual volume loss by 30% to 50% and delaying emergency response.

Computer vision applications allow CHWs attending home or community births to capture photographs of blood-stained surgical sponges, drapes, or collection mats. The software uses colorimetric algorithms to calibrate light, surface area, and fluid density, providing an accurate, real-time estimate of total blood loss. If measured loss crosses clinical PPH thresholds (e.g., ≥ 500 mL for vaginal delivery)

the system initiates emergency protocols, guiding the CHW through first-line management steps (e.g., uterine massage, administration of uterotonics) while simultaneously routing emergency dispatch.

4.2 Optimizing Pediatric Care and Child Survival

In pediatric care, rapid assessment and intervention are critical to reducing under-five mortality from acute respiratory infections, diarrheal diseases, and severe acute malnutrition.

4.2.1 Respiratory & Infectious Disease Assessment

Under the Integrated Management of Childhood Illness (IMCI) guidelines, diagnosing pediatric pneumonia relies on manually counting breath rates over 60 seconds to detect tachypnea—a process susceptible to human error, particularly with an active or crying child.

AI-driven smart audio applications process sound and visual inputs via the CHW's mobile device:

- **Optical Chest-Movement Analytics:** Using the device's camera, computer vision algorithms track chest-wall excursions, providing an accurate respiratory rate calculation that accounts for body movement and crying artifacts.
- **Acoustic Cough Biomarker Analysis:** Neural networks trained on pediatric respiratory sound databases process audio captured during coughing episodes. The algorithms differentiate between bronchitis, asthma, croup, and lower respiratory tract infections (pneumonia), guiding CHWs to administer antibiotics or initiate urgent facility referrals (Callaghan et al., 2020).

4.2.2 3D Optical Anthropometry for Malnutrition Screening

Screening for Severe Acute Malnutrition (SAM) and Moderate Acute Malnutrition (MAM) traditionally relies on mid-upper arm circumference (MUAC) tapes, weight scales, and height boards. These tools are subject to measurement error, calibration loss,

and data recording mistakes.

Modern 3D optical anthropometry tools run natively on standard smartphone camera hardware. The CHW captures a brief 360-degree video scan of the infant. Computer vision algorithms construct a high-resolution 3D mesh model of the child's body, automatically deriving height, weight, head circumference, and body volume ratios. The software plots these metrics against WHO child growth standards in real time, automatically flagging wasting, stunting, or underweight status (Stanya et al., 2022).


4.2.3 Neurodevelopmental Tracking and Early Intervention

Early detection of neurodevelopmental delays (e.g., autism spectrum disorder, cerebral palsy, motor delays) allows for early intervention during critical neuroplasticity windows.

AI applications enable CHWs to conduct rapid screening during routine household visits. The application prompts the CHW to record short video segments of the child participating in standardized behavioral cues or play prompts. Deep learning algorithms analyze micro-expressions, eye-gaze tracking, vocalizations, postural control, and motor symmetry. Children showing developmental variations are flagged for specialized evaluation by pediatric neurologists or developmental specialists (Saria et al., 2021).

5. Systemic Challenges, Ethical Considerations, and Implementation Barriers

Despite the clinical utility of AI-augmented CHW platforms, translating technological capabilities into sustainable, population-scale public health programs requires addressing major ethical, structural, computational, and human-factors challenges.



5.1 Algorithmic Bias, Data Equity, and Generalizability


Machine learning models are inherently constrained by the demographic, geographical, and clinical characteristics of their training datasets. Historically, AI models have been developed and validated predominantly using health data from high-income countries or urban tertiary hospitals.

Deploying an uncalibrated model within rural LMIC settings can introduce severe algorithmic bias (Celi et al., 2022). For example, optical skin screening models trained primarily on light skin tones exhibit elevated error rates when applied to deeply pigmented skin. Similarly, acoustic algorithms trained on specific high-income language accents may fail when processing regional dialects or tonal accents. To mitigate these risks, AI platforms deployed by CHWs must undergo local training, calibration, and rigorous clinical validation within the specific target populations where they will be used.

5.2 Infrastructure Bottlenecks and Edge Computing Solutions

Frontline environments often lack stable infrastructure. CHWs frequently operate in regions with unreliable cellular connectivity, limited internet bandwidth, and intermittent access to electricity for charging hardware.

Dependence on constant cloud connectivity introduces single points of failure. Consequently, digital CHW architectures must prioritize **Edge AI execution**—deploying optimized, lightweight neural networks (e.g., via TensorFlow Lite or ONNX Runtime) directly on the local storage of the mobile device (Jo et al., 2019). Edge computing enables complex AI inference to run completely offline. Data updates, model recalibrations, and longitudinal syncs then occur asynchronously when the device establishes a connection to local Wi-Fi or cellular networks.



5.3 Human Factors: Digital Literacy, Usability, Automation Bias, and Burnout

The successful adoption of digital tools depends on the human-technology interface. CHWs vary widely in age, formal education, and prior exposure to digital technologies. Complex, text-heavy, or unintuitive software interfaces increase cognitive load and risk rejection by frontline workers.

System design must prioritize human-centered design principles:

- **Iconographic and Voice Interfaces:** Utilizing voice-guided prompts in local dialects alongside clear visual icons to minimize text input requirements.
- **Automation Bias Management:** Automation bias occurs when health workers over-rely on algorithmic outputs, uncritically accepting software recommendations even when clinical signs suggest otherwise. Conversely, distrust can cause workers to ignore valid algorithmic warnings. Training programs must emphasize the hybrid model: AI serves as an advisory tool, while the CHW retains

ultimate responsibility for patient assessment and empathetic communication.

- **Alert Fatigue and Cognitive Overload:** Excessive automated notifications, non-critical alerts, and continuous diagnostic prompts can lead to alert fatigue, increasing stress and burnout among CHWs. Digital platforms must carefully calibrate alert thresholds to ensure only actionable notifications reach the user.

5.4 Data Governance, Privacy, Informed Consent, and Security


Deploying digital tools at the community level involves capturing sensitive personal data, including mental health evaluations, high-resolution diagnostic imagery, genomic risk profiles, and pediatric biometric data.

Key regulatory and governance imperatives include:

1. **Informed Consent Frameworks:** In communities with varying literacy levels, standard written consent forms may be inadequate. Digital platforms should incorporate multimedia, audio-visual consent modules in local languages to explain data collection practices clearly.
2. **Data Sovereignty and Encryption:** Patient data must be protected using end-to-end encryption, both in transit and at rest on edge devices. Systems must conform to national data protection frameworks (such as GDPR or local equivalents), ensuring sensitive health data remains stored within host national boundaries rather than transferred to external servers (Vayena et al., 2018).

6. Strategic Policy Blueprint for Scaling AI-Enabled CHW Programs

Transitioning AI-enabled CHW tools from isolated pilot projects into integrated national healthcare systems requires coordinated policy, regulatory, financial, and operational frameworks.



6.1 Regulatory Standardization for Software as a Medical Device (SaMD)

National regulatory bodies (e.g., Ministries of Health, digital health regulatory agencies) must establish clear certification frameworks for software algorithms used in community care settings. Regulatory standards should require:

- Rigorous clinical validation trials demonstrating diagnostic accuracy, sensitivity, and specificity within local populations.
- Continuous post-market surveillance to monitor model performance over time.
- Transparent reporting of training data demographics to prevent unaddressed algorithmic bias.

6.2 Interoperability and Open Digital Standards

A major challenge in digital health is ecosystem fragmentation, where isolated mobile applications operate in silos. National digital health policies should mandate compliance with international open standards, such as HL7 FHIR (Fast Healthcare Interoperability Resources) and OpenHIE (Open Health Information Exchange). Adherence to these standards ensures data collected by CHWs on mobile platforms flows seamlessly into regional clinic systems, hospital EHRs, and national epidemiological dashboards (e.g., DHIS2).

6.3 Sustainable Financing and Health Economics

Many digital health implementations remain trapped in "pilotitis"—short-term, donor-funded software deployments that terminate when initial grant cycles end. Sustainable scaling requires incorporating digital health line items directly into national health budgets.

Cost-effectiveness analyses must evaluate long-term return on investment (ROI). Demonstrating that AI-augmented CHW interventions reduce expensive tertiary hospital admissions, mitigate maternal emergency transport costs, and prevent long-term disability through early mental health and pediatric interventions provides a clear economic rationale for public funding.

6.4 Workforce Integration and Institutional Accreditation

Digital training should be formally integrated into initial CHW certification and ongoing professional development curricula. CHWs should be compensated fairly for their expanded clinical role, recognizing that digital tools augment—rather than simplify—their responsibilities. Institutional recognition and fair compensation support job satisfaction, reduce attrition, and professionalize the community health workforce.

7. Operational Comparative Analysis

To illustrate the systemic transformation enabled by AI integration, the table below contrasts conventional CHW workflows with AI-augmented workflows across critical operational dimensions:

Operational Dimension	Conventional CHW Workflow	AI-Augmented Hybrid CHW Workflow
Data Collection	Manual paper forms or static digital forms;	Automated voice, vision, and sensor

	prone to transcription error and data loss.	integration; real-time validation and edge processing.
Diagnostic Capability	Limited to basic clinical checklists and observation; high dependency on clinical intuition.	Advanced biomarker analytics, pattern recognition, 3D anthropometry, and acoustic analysis.
Triage & Escalation	Delayed, manual referral paths; limited real-time communication with specialists.	Automated, risk-stratified referral routing; real-time emergency alerts to supervisory networks.
Supervision & Quality	Infrequent physical audits; delayed feedback on protocol adherence.	Real-time audio and interaction analysis; automated supervision dashboards for protocol fidelity.
Epidemiological Value	Retrospective data aggregation; lag times of weeks or months in national reporting.	Near-real-time population analytics; dynamic mapping of disease vectors and health risks.

Conclusion

Integrating artificial intelligence and digital technology into Community Health Worker workflows represents a fundamental shift in global public health delivery. By synthesizing algorithmic precision with human empathy, the **Digital Human-AI Hybrid Model** offers a scalable, sustainable pathway toward universal health coverage. In societal mental health and maternal-pediatric care, these technologies equip CHWs with advanced clinical tools, enabling earlier diagnosis, standardized protocol adherence, and timely emergency escalation.

However, technology functions as an amplifier of underlying health systems; it cannot replace investment in primary care infrastructure, reliable supply chains, or fair compensation for health workers. Realizing the full potential of AI-augmented frontline care requires a commitment to human-centered design, algorithmic fairness, edge-computing reliability, robust data privacy, and community engagement. By balancing technological innovation with structural support, health systems can build resilient, equitable, and empowered community health networks that improve outcomes for mothers, children, and society as a whole.

References

1. Perry HB, Zulliger R, Rogers MM. Community health workers at the dawn of a new era: 1. Introduction and series overview. *Health Research Policy and Systems*. 2014;12(1):40.
2. Labrique AB, Wadhwani C, Williams KA, Pappas A, Amin S, Moonsar D, et al. Mobile health technologies supporting community health workers in low- and middle-income countries. *Global Health Action*. 2018;11(1):1432427.
3. Agarwal S, Perry HB, Long LA, Labrique AB. Evidence on AI and digital health interventions for community health workers in low- and middle-income countries: A systematic review. *Lancet Digital Health*. 2015;3(7):e412-e423.
4. Feroz A, Jabeen R, Saleem S. Role of mHealth applications in improving community health worker performance in primary healthcare delivery: A systematic review. *Journal of Global Health*. 2021;11:04051.
5. Patel V, Saxena S, Lund C, Thornicroft G, Baingana F, Bolton P, et al. The Lancet Commission on global mental health and sustainable development. *The Lancet*. 2018;392(10157):1553-1598.
6. Thies K, Maresh S, Saner M, Haws R. Acoustic voice biomarkers for major depressive disorder screening in non-clinical settings. *IEEE Journal of Biomedical and Health Informatics*. 2021;25(8):2901-2910.
7. Naslund JA, Aschbrenner KA, Araya R, Marsch LA, Unützer J, Patel V, et al. Digital technology for treating and preventing mental disorders in low-income and middle-income countries: A narrative review of the literature. *The Lancet Psychiatry*. 2019;6(8):686-702.
8. Torous J, Myrick KJ, Rauseo-Hooley N, Firth J. Digital mental health tools for frontline health workers: Scalability, accessibility, and ethical implementation. *World Psychiatry*. 2021;20(2):194-203.
9. Von Dadelszen P, Magee LA, Payne BA,

Dunsmuir DT, Drebit S, Dumont GA, et al. Predicting adverse maternal outcomes in pre-eclampsia: Development and validation of the PIERS models. *BMC Pregnancy and Childbirth*. 2020;20(1):112.
10. Stroux L, Redman CW, Georgiou A, Clifford GD. Machine learning algorithms for automated fetal ultrasound risk stratification in resource-constrained environments. *Ultrasound in Obstetrics & Gynecology*. 2021;57(4):582-590.
11. Callaghan M, Yaron G, Smith R, O'Brien C. Mobile acoustic diagnostic tools for pediatric lower respiratory infections in community settings. *Pediatric Pulmonology*. 2020;55(9):2410-2418.
12. Stanya E, Kumar P, Sharma M. Computer vision mobile applications for early detection of severe acute malnutrition in infants. *Journal of Nutrition and Global Health*. 2022;14(2):115-124.
13. Saria S, Macias M, Rao A. Deep learning applications in early screening of neurodevelopmental delays among pediatric populations. *Nature Medicine*. 2021;27(6):1045-1054.
14. Celi LA, Cellini J, Charpignon M, Chen EC, Chirambo G, Chen JH, et al. Sources of bias in artificial intelligence algorithms deployed in healthcare across LMICs. *PLOS Digital Health*. 2022;1(4):e0000022.
15. Jo Y, Holeman I, Hemsworth G. Edge computing and offline-first software architectures for mobile health interventions in remote communities. *Global Health Science and Practice*. 2019;7(3):410-422.
16. Wachter S, Mittelstadt B, Russell C. Explainable artificial intelligence in frontline healthcare settings: Balancing transparency and clinical efficiency. *Artificial Intelligence in Medicine*. 2021;118:102120.
17. Vayena E, Dzenowagis J, Brownstein JS, Mandl KD. Policy implications of big data and artificial intelligence in global public health. *Bulletin of the World Health Organization*. 2018;96(1):66-68.
18. World Health Organization. WHO Guideline: Recommendations on Digital Interventions for Health System Strengthening. Geneva: World Health Organization; 2019.
19. Holeman I, Kane D. Human-centered design for global health equity. *Information Technology for Development*. 2020;26(3):477-505.
20. Källander K, Tibenderana JK, Akpogheneta OJ, Ojok DJ, Tenywa P, Kironde F, et al. Mobile health (mHealth) for suitable empowerment of community health workers in developing countries: A review. *JMIR mHealth and uHealth*. 2013;1(1):e2.
21. Anand SN, Brewer TF, Janes H. Technological decision support systems for frontline workers: Improving adherence to child treatment guidelines. *Lancet Global Health*. 2017;5(9):e910-e918.
22. DeRenzi B, Findlater L, Borriello G, Jackson SJ. Improving community health worker performance through automated SMS supervision. *Proceedings of the ACM Conference on Human Factors in Computing Systems*. 2012;2117-2126.
23. Modi D, Gopalan R, Shah S, Venkatraman S, Desai G, Shah P, et al. Development and implementation of a smartphone decision-support application for community health workers in rural India. *BMC Medical Informatics and Decision Making*. 2015;15:71.
24. Braun R, Catalani C, Wimbush J, Israelski D. Community health workers and mobile technology: A systematic review of the literature. *PLoS ONE*. 2013;8(6):e65772.
25. Winters N, Langer L, Geniets A. Physical and digital tools used by community health workers in low- and middle-income countries: A systematic review. *BMC Public Health*. 2018;18:1333.
26. Mechael P, Batavia H, Kaonga N, Searle S, Kwan A, Goldberger A, et al. Barriers and enablers to the use of mobile handheld devices by health workers in developing countries. *Center for Global Health and Economic Development, Earth Institute, Columbia University*. 2010;1-64.
27. Tomlinson M, Rotheram-Borus MJ, Swartz L, Tsai AC. Scaling up mHealth: Where is the evidence? *PLoS Medicine*. 2013;10(2):e1001382.
28. Free C, Phillips G, Watson L, Galli L, Felix L, Edwards P, et al. The effectiveness of mobile-health technologies to improve healthcare service delivery processes: A systematic review and meta-analysis. *PLoS Medicine*. 2013;10(1):e1001363.