RESEARCH PAPER

Forensic Evaluation of Burned Female Bodies: Distinguishing Antemortem Abuse from Postmortem Thermal Effects

Abdulaziz Adnan Aljuraysan[1], Faris Albrahim[1], Faris Alsabt[1], Habeebuddin Shaji Mohammed[2]

[1]*Student, College of Medicine, King Faisal University, Al-Ahsa, Saudi Arabia*
[2]*Faculty Member, College of Medicine, King Faisal University, Al-Ahsa, Saudi Arabia*

Received: 17 Sep, 2026; Revised: 10th Oct 2026; Accepted: 18th Oct, 2026; Available Online: 10 Nov, 2026

ABSTRACT

Background: When a burned body is found, forensic investigators must decide whether the person was harmed before the fire or whether the fire itself caused death. Burns destroy evidence and conceal injuries, and the task is harder in communities where religious or cultural beliefs limit traditional autopsy.

Objective: To review the forensic methods used to distinguish antemortem injury and abuse from postmortem thermal effects in burned female bodies, with emphasis on signs of vitality, biological evidence of sexual assault, and situations in which autopsy is limited.

Methods: A narrative review of the published literature on fire deaths, vital reactions to burning, post-mortem imaging, heat-related DNA degradation and sexual assault evidence was carried out. Thirty-four sources were included and synthesised qualitatively under five themes.

Results: Soot deep in the airway and raised carboxyhaemoglobin were the most consistently reported indicators that the victim was alive during the fire; blistering with an inflammatory response, a line of redness, reparative changes and stomach ulcers linked to burn shock were supportive. Published autopsy series show that fire is used to conceal homicide. Post-mortem computed tomography separates heat-related from traumatic fractures and guides sampling. Heat progressively reduces recoverable DNA in blood and semen, and complete burning usually yields no profile, although partial profiles are sometimes obtained. Physical genital findings and semen detectability decline as the depth of genital burning increases.

Conclusion: No single method is sufficient. Only by combining scene findings, external examination, imaging, toxicology, histology and DNA analysis can investigators piece together what truly happened and deliver justice for the victim; a negative finding in a burned body does not exclude prior assault.

Keywords: Burned bodies, antemortem burns, postmortem burns, forensic DNA analysis, seminal stains, sexual assault, post-mortem computed tomography

How to cite this article: Abdulaziz Adnan Aljuraysan, Faris Albrahim, Faris Alsabt, Habeebuddin Shaji Mohammed, Forensic Evaluation of Burned Female Bodies: Distinguishing Antemortem Abuse from Postmortem Thermal Effects. Int J Drug Deliv Technol. 2026;16(91s):28-37. DOI: 10.25258/ijddt.16.91s.5

Source of support: This research was supported by the Deanship of Scientific Research and the Vice Presidency for Graduate Studies and Scientific Research at King Faisal University, Al-Ahsa, Saudi Arabia, under Grant No. KFU26434.

Conflict of interest: None

INTRODUCTION

Burned bodies represent one of the most challenging scenarios in forensic investigations due to the extensive destruction caused by fire and the ethical, legal, and cultural constraints that may limit traditional postmortem examinations. In many communities, autopsy procedures may be restricted or avoided for religious and societal reasons, placing an obligation on forensic scientists to adopt alternative approaches that respect human dignity while still ensuring justice. Protecting the rights of deceased individuals and accurately determining the circumstances of death are fundamental responsibilities within forensic medicine.

Fire deaths are also a public-health problem that, in some regions, falls disproportionately on women. A national analysis from India estimated more than 163,000 fire-related deaths in 2001, about 2% of all deaths and six times the number reported by the police. About 106,000 of these deaths were in women, mostly between 15 and 34 years of age, a pattern the authors attributed to kitchen accidents, self-immolation and different forms of domestic violence.[1]

An important point in the examination of burned corpses is determining whether the individual was subjected to abuse or fatal injury prior to the fire or whether death resulted directly from thermal injury and inhalation of combustion products. Differentiating between antemortem and postmortem burning is essential, as it directly influences the interpretation of injuries, the determination of the cause and manner of death, and subsequent legal outcomes. The identification of vital reactions to heat, smoke

**Author for Correspondence:* Abdulaziz Adnan Aljuraysan, azaljurysan@gmail.com

inhalation, and associated trauma can provide crucial evidence in the absence of a full autopsy.

This distinction matters because fire is sometimes used deliberately. In a French autopsy series, criminal deaths were the second most frequent category of deaths by burning, after accidents and ahead of suicide, and criminal burning appeared mainly to be a means of covering up a homicide.[2] Series of homicide victims who were burned after death describe the same intention to destroy evidence and hinder identification.[3,4]

However, fire can obscure or destroy critical forensic evidence, making it difficult to assess pre-burn injuries or establish abuse. In such cases, investigators often face limited and fragmented evidence, necessitating the use of multiple complementary forensic techniques. These may include toxicological analysis, histopathological examination, radiological imaging, DNA profiling, and detailed scene investigation, all of which contribute to reconstructing events surrounding death. Where dissection is restricted, a review of autopsy in Islam and its practice in Arab Muslim countries recommended wider application of non-invasive autopsy,[5] and cross-sectional imaging has been applied to charred bodies for this purpose.[6,7]

This research aims to evaluate the forensic methods used to determine whether a burned body was subjected to abuse before burning, with particular focus on distinguishing antemortem from postmortem injuries when autopsy options are limited. By examining current forensic practices and challenges, this study seeks to contribute to more accurate death investigations while maintaining ethical, legal, and societal considerations.

MATERIALS AND METHODS

Study Design

This study is a narrative (non-systematic) review of published literature. It was designed to bring together evidence from forensic pathology, forensic radiology, forensic genetics and clinical forensic medicine that bears on a single practical question: how can antemortem injury and abuse be distinguished from postmortem thermal effects in a burned female body?

Literature Search

The literature was identified through electronic searches of PubMed/MEDLINE-indexed journals, supplemented by publisher websites and open-access repositories, and by screening the reference lists of the retrieved articles. No lower date limit was applied, and the last search was run in November 2026. Search terms were used alone and in combination and included: burned bodies, charred body, fire death, antemortem burn, postmortem burn, vitality, soot, carboxyhaemoglobin, homicide, criminal burning, post-mortem computed tomography, virtual autopsy, autopsy and religion, DNA degradation, temperature, short tandem repeat, burnt bone, seminal stain, spermatozoa persistence, and sexual assault.

Eligibility Criteria

Sources were eligible if they were published in English and reported on (i) autopsy or case series of fire deaths or burned bodies, (ii) signs used to diagnose vitality during burning, (iii) post-mortem imaging of burned bodies or alternatives to conventional autopsy, (iv) the effect of heat or burning on blood, semen, bone or DNA typing, or (v) the examination of, and evidence collection from, victims of sexual assault. Original studies, case series, reviews, an international guideline and a standard textbook were all accepted. Sources unrelated to forensic practice were excluded.

Data Extraction and Synthesis

For each source, the study design, the material examined and the main findings relevant to the review question were extracted. Because the sources differed widely in design, material and outcome measures, no pooled statistical analysis was attempted. The findings were instead grouped and synthesised qualitatively under five themes: burn severity and heat-related changes; antemortem versus postmortem burning; burning used to conceal homicide; post-mortem imaging when autopsy is restricted; and biological evidence of sexual assault after exposure to heat.

Ethical Considerations

The study is based entirely on previously published material. No human participants, human tissue or identifiable personal data were involved, and ethical approval was therefore not required.

REVIEW OF LITERATURE

Classification and Severity of Burns

Burns are injuries that occur to the body by dry heat, heated material or radiation.

For the classification of burn severity, there are many criteria. Dupuytren's classification includes six degrees of burns.

First degree: This is a simple erythema or redness produced by heat. It is superficial, with only inflammation seen; the pain disappears in 2 to 3 days. It involves the epidermis, which is superficially affected.

Second degree: Blisters are found and redness is produced on application of heat. The epidermis is totally affected; healing takes up to one week.

Third degree: The epidermis is destroyed, and the dermis is involved. This burn is quite painful as nerve endings are totally exposed.

Fourth degree: The whole skin is destroyed; it is not painful, and the skin comes away as sloughs after several days.

Fifth degree: The dermis is completely destroyed. In this stage, muscles are also affected, which can lead to nerve deformities.

Sixth degree: Charring of the whole limb may be seen, and the bones are also charred.

The effect of burning differs based on many factors, including:

1. **Intensity of heat:** Burns may result from high or low heat.
2. **Duration of exposure:** Longer exposure results in more severe burn damage.
3. **Area involved:** According to the "Rule of Nines" (Table 1), 60 to 80% of superficial burns can be fatal, while 40% of deep burns can also be fatal. However, there are some cases where patients survived despite higher percentages of burned body surface area.
4. **Site of burns:** Burns on the trunk, especially over the abdomen, are more severe compared to burns on the limbs. Burns to the genitalia carry a higher risk of fatality.
5. **Age:** Children can tolerate burns better, while elderly people have higher mortality compared to adults.
6. **Sex:** Males tend to tolerate burns better than females.

Table 1: Body surface area according to the Rule of Nines

Area	Percentage
Head and neck	9%
Right upper limb	9%
Left upper limb	9%
Right lower limb	18%
Left lower limb	18%
Anterior trunk	18%
Posterior trunk	18%
Genitalia	1%
Total	100%

Although the genitalia represent a small percentage of total burn injuries, burns in this area are associated with higher mortality rates.[8]

Heat-related Changes of the Body

Several changes seen in a burned body are produced by heat alone and must not be mistaken for injury inflicted in life. With burning, dehydration and denaturation of protein shorten the muscles, so that the limbs flex and the body takes up the position of a boxer.[4,9] This posture develops quickly after exposure to the fire and is a thermal effect; it does not by itself show that the person was alive.[4]

The degree of destruction of the body is related to the duration of the fire, as shown by observations made during cremation,[10] and prolonged burning leads to fragmentation of bone that complicates both recovery at the scene and later interpretation.[9] Heat also produces fractures and collections of blood within the skull. On post-mortem computed tomography, heat-related fractures can be differentiated from traumatic fractures, and epidural collections are described as typical heat-related head lesions rather than evidence of a blow.[7]

DNA Evidence and Environmental Conditions

DNA fingerprinting has revolutionised forensic science by providing investigators with a reliable way to identify individuals and solve crimes. One of its greatest advantages is its high sensitivity: even a very small sample can provide enough genetic information to link someone to a crime scene or confirm their identity.[11,12] Advances in forensic DNA technologies have further improved the accuracy and efficiency of this process.[13]

In practice, however, evidence is not always collected under ideal conditions. The success of DNA analysis depends not only on the techniques themselves but also on how well samples are handled from the moment they are discovered. Factors such as heat, humidity, and ultraviolet light can damage DNA over time, reducing its usefulness.[14,15] In more extreme situations, fire may be used intentionally to destroy evidence, making forensic analysis even more challenging.[16] Studies have shown that high temperatures can negatively affect STR typing, although partial DNA profiles can still be recovered.[17]

Blood and seminal stains are among the most important types of biological evidence, especially in violent and sexual crimes. Therefore, understanding how environmental conditions affect them is essential. Previous studies have demonstrated that temperature, humidity, and pH significantly influence both the quantity and quality of DNA in biological samples.[14,18] This study examines how different levels of temperature and burning affect DNA recovery, providing insight into how forensic experts can manage damaged evidence and still reach reliable conclusions.

When soft tissue has been consumed, bone may be the only material left for identification. In a study of 71 bone fragments from 13 burned individuals, identification by DNA analysis was reliable and reproducible from well-preserved and semi-burnt bone. In black burnt bone the DNA was highly degraded, sometimes leaving mitochondrial DNA analysis as the only option, and blue-grey and blue-grey-white burnt bone only sporadically or barely gave authentic profiles; heavily burnt bone was also very prone to contamination with external DNA.[19]

Seminal Stains and Evidence of Sexual Assault

The analysis of seminal stains is an important part of forensic investigations, especially in cases involving sexual violence. However, the reliability of this evidence depends mostly on how it is collected, preserved, and examined. Proper collection is the first essential step; samples must be handled carefully and allowed to air-dry in order to reduce contamination and

protect the quality of the evidence. In addition, the type of material on which the semen is deposited plays a role, since different fabrics can affect how well the stain is retained and how easily DNA can be recovered.[20]

Environmental conditions can strongly affect the quality of semen evidence. Factors such as temperature, humidity, and pH have been shown to significantly impact the quality of semen evidence. High temperatures and prolonged exposure to unsuitable conditions can accelerate DNA degradation, while extreme pH levels may damage the cellular components within the sample.[14,18] Despite these challenges, studies suggest that usable DNA can often still be recovered, although the resulting profiles may be partial or less reliable, especially when the samples are old or degraded.[21] Enzymatic activity and biochemical processes may also contribute to DNA breakdown over time.[22]

The examination of seminal stains usually begins with presumptive tests, which help indicate the possible presence of semen. These are followed by confirmatory methods, such as microscopic identification of sperm cells or detection of semen-specific biomarkers. In situations where spermatozoa are absent or have been degraded, molecular techniques provide another valuable method of analysis.

DNA extraction followed by Short Tandem Repeat (STR) analysis remains the most reliable method for identifying individuals from seminal stains. This technique is highly sensitive and can produce results even from small or partially degraded samples.[12,23] However, the success of STR analysis depends heavily on the condition of the evidence before testing.

Time also limits what can be found. In about 5,581 swabs from some 1,450 sexual assault cases examined at one national laboratory, the proportion of vaginal and anal swabs containing sperm declined significantly after 48 hours from intercourse, and 23% of sperm-positive swabs gave a negative acid phosphatase reaction, which led the authors to question that presumptive test as a predictor of sperm.[24] Evidence from the dead has been studied less often; one medical examiner's office reviewed 199 cases that were both autopsied and examined for sexual assault over ten years, using oral, vaginal and anorectal swabs to relate spermatozoa and prostatic acid phosphatase to the postmortem interval.[25]

Where male and female cells are mixed, as on an intimate swab, typing of STRs on the Y chromosome identifies male-specific DNA and is therefore especially useful in sexual assault cases. Its limitation is that male relatives share the same Y-STR profile over several generations, which must be allowed for when a match is interpreted.[26]

Overall, advances in forensic science have made it possible to obtain valuable information even from compromised seminal stains. However, these findings also highlight the importance of careful evidence handling and storage to ensure accurate and reliable results.

Antemortem versus Postmortem Burns

Another subject that needs to be addressed is whether the burn is antemortem or postmortem. There are many features that help identify whether the burn occurred before or after death. The first feature is the line of redness. This line only appears if the burn is antemortem, which occurs because of a reaction that happens between the normal skin and the burnt part of the skin. This line may still be present after the death of the person, and it also causes redness because of the widening of capillaries beyond the burned area.

Another important feature is vesication. These are simply blisters that form if the burn is antemortem and have specific features that distinguish them from blisters that form postmortem. Features of antemortem blisters include a red, inflamed base containing a lot of serous fluid and white blood cells (WBC), unlike blisters that form postmortem, which have features like a hard, dry, and horny base, much less fluid and no WBC.

Another important feature of antemortem burns is the reparative process. Antemortem burns have granulations and other reparative processes that are present if the burn occurred before death. Simply put, any sign of a recovery process is an indication of an antemortem burn. And since recovery processes can happen while the person is alive, we can also notice another feature, which is enzyme activity. For example, redness and tissue cathepsin may be present immediately after the burn, along with many other enzymes that appear over the course of a week or even longer.

Another feature that can be present in antemortem burns is Curling's ulcer. It is an ulcer that forms inside the stomach and/or duodenum with severe burns while the victim is alive. The cause is that a severe burn leads to hypovolemia and shock, which decrease gastric blood flow and lead to ischemia. Eventually, an ulcer forms due to the increased gastric acid secretions. On the other hand, it will be absent in postmortem burns.

Another feature is the blood color, which will present as cherry red due to carbon monoxide inhalation. In postmortem burns, blood will be normal in color since there was no breathing of carbon monoxide or inhalation of any other fumes or smoke.

The last, and a very important, feature is the presence of soot particles in the upper respiratory tract. It is an easily distinguishable feature: if the victim inhales smoke coming from the fire, soot will be present in the respiratory passages. In postmortem burns it will be absent.[8]

The autopsy literature supports the weight given to these last two signs. Soot deposits in the respiratory tract and a raised carboxyhaemoglobin (COHb) concentration in the blood are regarded as the

characteristic parameters of vitality in burned bodies, although the diagnosis remains difficult in a proportion of cases in which these signs are weak or absent.[27] One recent series described soot in the respiratory tract as one of the main antemortem signs and noted that a COHb concentration above 50% indicates that the person died in the fire.[4]

Burning Used to Conceal Homicide

Fire may be the means of killing or a means of hiding a killing. In the French series already mentioned, burning of a body after a homicide was more common than killing by fire itself, and the authors considered it essential to take X-rays, to look for injuries due to trauma and to carry out systematic toxicological analysis in every victim of burning.[2] A series of homicides followed by postmortem burning found that such bodies are typically discovered in an unfamiliar, outdoor or abandoned place, with findings of a violent death, use of an accelerant and negative signs of vitality.[3] Four homicide cases reported from Lithuania illustrated how difficult the distinction can be on autopsy findings alone and how histological examination, toxicological analysis and post-mortem computed tomography add to the autopsy.[4]

Post-mortem Imaging When Autopsy is Restricted

In an early report, multi-slice computed tomography and magnetic resonance imaging of a charred body from a vehicle fire documented both the injuries caused by burning and forensically relevant vital reactions, namely air embolism and blood aspiration, before the conventional autopsy was performed.[6] Post-mortem computed tomography (PMCT) is now described as an accessible tool for the investigation of burned bodies. It helps to find features that allow identification, such as medical devices and prostheses; it depicts metallic foreign bodies of ballistic origin; it shows whether liquid blood or urine remains available for toxicological analysis; and it can show protected sites, such as the spinal canal, from which tissue can be taken for DNA sampling.[7]

These properties are directly relevant where families or religious authorities object to dissection. Attitudes towards the dead body vary with religious belief and cultural background, and the review of practice in Arab Muslim countries called both for more published experience from the region and for more use of non-invasive autopsy.[5]

Effect of Temperature and Burning on DNA Recovery

Blood and sperm samples can be successfully DNA-typed only when they are preserved under suitable temperature conditions. When the samples are exposed to high temperatures, such as 50°C or 100°C, the amount of extractable DNA decreases sharply. This occurs because heat damages the cells, breaks down internal proteins, and causes DNA strands to degrade.

Although the DNA quantity may be decreased after heat exposure, the remaining DNA can still be useful for forensic analysis. In certain heated samples, scientists were able to obtain DNA profiles, showing that DNA can survive even under challenging conditions. However, lower temperatures such as 37°C, 4°C and −20°C are good for DNA preservation, as there is no significant loss in DNA concentration, or at least the loss is minimal.

The most destructive condition was burning. In burned samples, the DNA was heavily damaged and no usable DNA profile could be obtained. Overall, these findings show a clear relationship between temperature and DNA survival: higher temperatures reduce DNA survival, while lower temperatures preserve it. This highlights the significance of storing samples correctly for forensic work.[28]

RESULTS

Overview of the Included Literature

Thirty-four sources published between 1998 and 2023 were included. They comprised autopsy and case series of fire deaths, experimental studies of heat and storage effects on blood, semen, bone and DNA typing, casework series on sexual assault evidence, imaging reports, narrative reviews, one international guideline and one textbook. The main findings of the studies that carried most weight in the synthesis are set out in Table 2.

Signs of Vitality

The sources agreed that soot in the airways and raised COHb are the principal indicators that a victim breathed during the fire.[4,8,27] In 115 unselected autopsy cases of death from thermal effects or fire, 85 victims died at the scene and 23 died during initial medical care or clinical treatment; of those who died at the scene, 77 showed intensive thermal or fire effects, while eight showed no thermal effects at all, seven of them having died of smoke poisoning and one of polytrauma after a leap from a height.[29] The external and internal features that separate antemortem from postmortem burns are summarised in Table 3.

Manner of Death and Concealment

Among deaths by burning in the French series, accidents accounted for 52%, criminal deaths for 31% and suicides for 16%.[2] Homicides followed by postmortem burning shared a recognisable profile: an unfamiliar outdoor scene, a traumatic cause of death, absence of vital reaction to the fire and the presence of an accelerant.[3]

Imaging

PMCT and magnetic resonance imaging documented burn injuries and vital reactions without dissection[6] and,

in burned bodies, distinguished heat-related from traumatic fractures, located ballistic foreign bodies and showed where fluid and tissue remained for toxicology and DNA sampling.[7]

Biological Evidence

The amount of extractable DNA in blood and seminal stains fell sharply at 50°C and 100°C, was preserved at 37°C, 4°C and −20°C, and gave no usable profile after burning.[28] In burnt bone, reliable typing was confined to well-preserved and semi-burnt fragments.[19] In living complainants, the proportion of sperm-positive vaginal and anal swabs declined significantly after 48 hours.[24] Bringing these findings together with the literature on genital injury, Table 4 relates the type of genital burn to the physical findings that may remain, the detectability of semen and the forensic points that follow.

Table 2: Main findings of the principal studies included in the review

Source	Material or design	Main finding
Sanghavi et al.[1]	National mortality data, India, 2001	More than 163,000 fire-related deaths (about 2% of all deaths); about 106,000 in women, mostly aged 15 to 34 years.
Gerling et al.[29]	115 autopsy cases of death from thermal effects or fire, 1990 to 1999	85 deaths at the scene and 23 during medical care; 8 scene deaths showed no thermal effects (7 smoke poisoning, 1 polytrauma).
Bohnert et al.[27]	Autopsy study of vitality in burned bodies	Soot in the respiratory tract and raised COHb are the characteristic parameters of vitality; diagnosis remains problematic in some cases.
Fanton et al.[2]	Autopsy series of deaths by burning, France	Accidents 52%, criminal deaths 31%, suicides 16%; criminal burning mainly used to cover up homicide. X-rays, search for trauma and systematic toxicology advised.
Tümer et al.[3]	Series of homicides followed by postmortem burning	Bodies typically found in unfamiliar outdoor places, with traumatic cause of death, no vital reaction and use of an accelerant.
Žiūkaitė et al.[4]	Four burnt homicide victims and literature review	Soot in the airway is a main antemortem sign; COHb above 50% indicates death in the fire; histology, toxicology and PMCT add to autopsy.
Thali et al.[6]	Charred body examined by MSCT and MRI before autopsy	Imaging documented burn injuries and vital reactions (air embolism, blood aspiration).
Coty et al.[7]	Pictorial review of PMCT in burned bodies	Heat-related fractures distinguishable from traumatic fractures; epidural collections typical of heat; imaging guides identification, toxicology and DNA sampling.
Schwark et al.[19]	71 bone fragments from 13 burned individuals	Reliable DNA identification from well-preserved and semi-burnt bone only; black, blue-grey and white burnt bone highly degraded.
Abdel Hady et al.[28]	Experimental heating and burning of blood and seminal stains	DNA yield falls sharply at 50°C and 100°C; no usable profile after burning; cold storage preserves DNA.
Casey et al.[24]	About 5,581 swabs from about 1,450 sexual assault cases	Sperm on vaginal and anal swabs declines significantly after 48 hours; 23% of sperm-positive swabs were acid phosphatase negative.
Roewer[26]	Review of Y-chromosomal STR typing	Male-specific typing is especially useful in male and female cell mixtures; male relatives share the same profile.

Table 3: Features distinguishing antemortem from postmortem burns[4,8,27]

Feature	Antemortem burn	Postmortem burn
Line of redness	Present at the margin between burnt and normal skin	Absent
Vesication (blisters)	Red, inflamed base; abundant serous fluid containing white blood cells	Hard, dry, horny base; little fluid; no white blood cells

Reparative process	Granulation and other signs of repair if the victim survived for a time	Absent
Enzyme activity	Present; tissue cathepsin early, other enzymes over the following days	Absent
Curling's ulcer	May be present in the stomach or duodenum after severe burns	Absent
Blood colour and COHb	Cherry red; raised COHb	Normal colour; COHb not raised
Soot in the respiratory tract	Present	Absent

Table 4: Physical findings, semen detectability and forensic points according to the type of genital burn

Type of genital burn	Physical findings	Is semen detectable?	Forensic points
Vagina is intact	No vaginal burn is present. Possible bruising of the hymen may be found. If the hymen is intact, abrasion may be found.[30]	Yes, by DNA.[28]	Genital injuries can be detected in the anus and the vagina and at extra-genital sites.[31] The type of injuries sustained (particularly to the mouth, breasts, vagina, and rectum) can help forensic assessment.[32] An important point is that an intact vagina does not rule out assault.[31]
Superficial burn of the vagina	At this stage, blistering, redness, thickening and swelling can be seen. However, if the cervix remains intact, small tears can be observed.[8]	Possible if the burn is limited, but it cannot be confirmed if the female genitalia are fully burned.[28]	Usually it is not reliable.[30] A deep vaginal swab should be taken because, if the sample has been exposed to excessive heat, the DNA may not be detectable.[33]
Charring of the vagina	There will be destruction of the vaginal tissue and shrinking of the vagina. The hymen and vagina may no longer be anatomically recognisable; true trauma may not be easily found.[30,34]	Not detectable, but not impossible.[28,34]	A negative DNA result does not rule out assault.[33]

DNA can usually be detected at normal temperatures, but extreme burning usually reduces the physical findings available to forensic scientists.[33]

DISCUSSION

Principal Findings

The literature reviewed gives a consistent answer to the question posed: no single sign or test separates antemortem abuse from postmortem thermal effects, and each line of evidence fails in a predictable way. Skin signs depend on survival for long enough to mount a reaction; airway soot and COHb depend on breathing in a smoke-filled atmosphere; physical genital findings depend on tissue that fire destroys early; and DNA depends on temperature and duration of exposure. A sound conclusion therefore rests on agreement between several independent findings, not on any one of them.[4,27]

Interpreting Signs of Vitality

Soot in the airway and raised COHb remain the strongest evidence that a victim was alive in the fire.[8,27] Their absence is harder to interpret. It is expected when a body is burned after death, and in the homicide series it was one of the features that raised suspicion,[3] but the diagnosis of vitality is acknowledged to be problematic in some genuine fire deaths.[27] The series of Gerling and colleagues is a reminder that the reverse also occurs: some victims die at a fire scene without any thermal injury, from smoke or from trauma.[29] Absent vital signs should therefore prompt a search for another cause of death, and should not on their own be read as proof of homicide.

Heat-related changes are a second source of error. The pugilistic posture, heat fractures and epidural collections are produced by fire regardless of how the person died,[4,7,9] and an examiner who is not familiar with them may report violence that did not occur. Equally, real injuries may be dismissed as heat effects. Radiological examination before any dissection helps with both problems.[2,7]

The Burned Female Victim

The excess of fire deaths among young women in the Indian data, and the link drawn by its authors to domestic violence,[1] mean that the death of a woman by burning should be approached with abuse in mind from the outset. Published homicide series show that fire is chosen precisely because it is expected to remove the evidence.[2,3] For the female victim the evidence at greatest risk is that of sexual assault, because the genital tissues are thin, superficial and quickly altered by heat (Table 4).

Two points follow. First, physical findings were never a dependable test of assault even in the unburned body: an intact hymen or vagina does not exclude it,[30,31] and burning only widens that uncertainty. Second, the biological evidence is fragile for reasons that add together. Semen is lost with time, with sperm becoming much less frequent on swabs after 48 hours in living complainants,[24] and its DNA is degraded by heat and destroyed by direct burning.[28] A negative swab from a burned body is therefore to be expected in many true cases and must be reported as inconclusive.[33]

Making the Most of Limited Evidence

The same literature points to practical ways of improving the yield. Deep vaginal swabs sample the site best shielded from heat.[33] Y-chromosomal STR typing can detect a small male component in a predominantly female sample, with the caution that it identifies a paternal lineage and not an individual.[26] Stains on clothing, bedding or other items that lay away from the seat of the fire may have been exposed to lower temperatures than the body, at which DNA yield is reduced but typing is often still possible.[17,28] Where soft tissue is lost, bone can still identify the victim provided that well-preserved or semi-burnt fragments are selected.[19]

When Autopsy is Restricted

In settings where dissection is limited on religious or cultural grounds, the reviewed evidence supports a stepwise, minimally invasive approach: careful scene investigation; full external examination with attention to the features in Table 3; whole-body PMCT to look for trauma, foreign bodies and heat-related change and to locate fluid and tissue for sampling;[6,7] toxicological analysis including COHb; and targeted swabs and tissue samples for DNA. Such an approach is in keeping with the recommendation that non-invasive autopsy be applied more widely in Arab Muslim countries.[5] Imaging cannot replace histological confirmation of a vital reaction in the skin, and the case for a full or partial autopsy should be made to the authorities when the findings point to homicide.

Evaluation of the Sexually Assaulted Victim

The definition of rape or sexual assault differs from one source to another. However, it is typically defined as penetration (whether deep or superficial) of the vagina or anus with any part of the body or an object, or even oral penetration. All of these actions are considered rape. Sexual assault is often accompanied by emotions such as anger and aggression, and these emotions can result in injuries in approximately 40% of rape victims, either in the genitalia or any other part of the body. For example, extra-genital injuries, such as bruises or lacerations, may occur due to the victim's resistance during the assault.

Other potential consequences of rape include infections such as hepatitis, HPV, HIV, or any other sexually transmitted infections (STIs). Psychological symptoms may also cause long-term effects such as post-traumatic stress disorder (PTSD).

Evaluation of a sexually assaulted victim should follow the principles applied to patients with trauma. The goals of the evaluation include medical assessment and treatment of injuries, prevention of unwanted pregnancy, and the usual forensic evidence collection. Psychological assessment and appropriate treatment are also important.

History taking and physical examination should be performed with the patient's consent; some victims may feel embarrassed or frightened. The examiner should be reassuring and empathetic, conducting the history and examination slowly and carefully. Important details about the sexual assault must be documented, including the types of injuries, their location on the body, any bleeding or abrasions, a description of how the attack occurred, whether the assailant used any weapons, and finally a description of the assailant.[32] These principles apply equally to the woman who survives a burn injury, in whom extra-genital injuries of resistance may be the only surviving physical evidence.

Limitations

This is a narrative review and not a systematic one: the search was not exhaustive, study quality was not formally appraised, and selection of sources may have introduced bias. The included studies were heterogeneous and could not be pooled. Much of the evidence on DNA comes from experimental stains heated under controlled conditions, which may not reproduce the conditions inside a burning body. Finally, few of the sources addressed genital findings in burned bodies directly, so Table 4 rests largely on extrapolation from the literature on sexual assault examination and on heat effects on DNA, and should be tested against casework.

CONCLUSION

Distinguishing antemortem abuse from postmortem thermal effects in a burned female body is possible but

depends on combining methods. Soot in the airway, raised carboxyhaemoglobin and vital skin reactions indicate life during the fire; their absence, together with trauma, an unusual scene or an accelerant, should raise suspicion of a concealed homicide. Heat-related changes must be recognised for what they are. Evidence of sexual assault becomes harder to demonstrate as genital burning deepens, and a negative result never excludes assault. Where conventional autopsy is restricted, external examination, post-mortem computed tomography, toxicology and targeted DNA sampling offer a respectful alternative that still serves justice. Prospective casework studies of burned victims, particularly from regions where autopsy is limited, are needed.

ACKNOWLEDGMENTS

The authors thank the Deanship of Scientific Research and the Vice Presidency for Graduate Studies and Scientific Research at King Faisal University, Al-Ahsa, Saudi Arabia, for supporting this work.

DECLARATIONS

Funding: This research was supported by the Deanship of Scientific Research and the Vice Presidency for Graduate Studies and Scientific Research at King Faisal University, Al-Ahsa, Saudi Arabia, under Grant No. KFU26434.

Conflict of interest: The authors have no competing interest.

Ethical approval: Not required; the study is based on published literature and involved no human participants or identifiable data.

Data availability: No new data were generated; all material analysed is available in the cited sources.

Author contributions: A.A.A. conceived the study, led the literature search and wrote the first draft. F.Alb. and F.Als. contributed to the literature search, data extraction and drafting of the manuscript. H.S.M. supervised the work and critically revised the manuscript for important intellectual content. All authors read and approved the final version and agree to be accountable for all aspects of the work.

Use of artificial intelligence: An AI-assisted tool (Claude, Anthropic) was used to support the literature search, language editing and formatting of the manuscript. The authors reviewed and verified the content and take full responsibility for it.

REFERENCES

1. Sanghavi P, Bhalla K, Das V. Fire-related deaths in India in 2001: a retrospective analysis of data. Lancet. 2009;373(9671):1282-8.
2. Fanton L, Jdeed K, Tilhet-Coartet S, Malicier D. Criminal burning. Forensic Sci Int. 2006;158(2-3):87-93.
3. Tümer AR, Akçan R, Karacaoğlu E, Balseven-Odabaşı A, Keten A, Kanburoğlu Ç, et al. Postmortem burning of the corpses following homicide. J Forensic Leg Med. 2012;19(4):223-8.
4. Žiūkaitė G, Jasaitė M, Chmieliauskas S, Vasiljevaitė D, Laima S, Banionis D, et al. Homicides disguised as fire deaths. Acta Med Litu. 2023;30(1):86-96.
5. Mohammed M, Kharoshah MA. Autopsy in Islam and current practice in Arab Muslim countries. J Forensic Leg Med. 2014;23:80-3.
6. Thali MJ, Yen K, Plattner T, Schweitzer W, Vock P, Ozdoba C, et al. Charred body: virtual autopsy with multi-slice computed tomography and magnetic resonance imaging. J Forensic Sci. 2002;47(6):1326-31.
7. Coty JB, Nedelcu C, Yahya S, Dupont V, Rougé-Maillart C, Verschoore M, et al. Burned bodies: post-mortem computed tomography, an essential tool for modern forensic medicine. Insights Imaging. 2018;9(5):731-43.
8. Sharma K. Concise textbook of forensic medicine and toxicology. 2011.
9. Ubelaker DH. The forensic evaluation of burned skeletal remains: a synthesis. Forensic Sci Int. 2009;183(1-3):1-5.
10. Bohnert M, Rost T, Pollak S. The degree of destruction of human bodies in relation to the duration of the fire. Forensic Sci Int. 1998;95(1):11-21.
11. Aronson JD, Cole SA. Science and the death penalty: DNA, innocence, and the debate over capital punishment in the United States. Law Soc Inq. 2009;34(3):603-33.
12. Jobling MA, Gill P. Encoded evidence: DNA in forensic analysis. Nat Rev Genet. 2004;5(10):739-51.
13. Horsman KM, Bienvenue JM, Blasier KR, Landers JP. Forensic DNA analysis on microfluidic devices: a review. J Forensic Sci. 2007;52(4):784-99.
14. Al-Kandari N, Singh J, Sangar V. Time dependent effects of temperature and humidity on quantity of DNA in samples of human saliva, blood and semen in Kuwait. Int J Pharma Sci Res. 2016;7(7):2852-73.
15. Khushbu K, Shalika N, Rashmi K. Identification of blood stains under different environmental conditions. Int J Biomed Res. 2017;8(12):707-10.
16. Tontarski KL, Hoskins KA, Watkins TG, Brun-Conti L, Michaud AL. Chemical enhancement techniques of bloodstain patterns and DNA recovery after fire exposure. J Forensic Sci. 2009;54(1):37-48.

17. Barbaro A, Cormaci P. Study about the effect of high temperatures on STR typing. Forensic Sci Int Genet. 2008;1(1):92-4.
18. Kannika S, Panvisavas N, Chaikum N. Effect of temperature and pH on bloodstain evidence. Forensic Sci Int. 2009;8(4):21-6.
19. Schwark T, Heinrich A, Preusse-Prange A, von Wurmb-Schwark N. Reliable genetic identification of burnt human remains. Forensic Sci Int Genet. 2011;5(5):393-9.
20. Raina A, Pramanik P, Dogra T. Effect of storage conditions of seminal stains on different textures of cloths in relation to DNA yield. Indian Internet J Forensic Med Toxicol. 2005;3(2):384-6.
21. Nakanishi H, Hara M, Takahashi S, Takada A, Saito K. Evaluation of forensic examination of extremely aged seminal stains. Leg Med (Tokyo). 2014;16(5):303-7.
22. Antrim RL, Buchert J, Burrows H, et al. Industrial enzymes: enzymes in nonfood applications. Enzymes Ind Prod Appl. 2004;2:155-244.
23. Somiari RI, Adebiyi E, Ukachukwu L, et al. STR analysis of human DNA samples after dry-state ambient temperature storage in GenPlates. Forensic Sci J. 2011;4:30-5.
24. Casey DG, Domijan K, MacNeill S, Rizet D, O'Connell D, Ryan J. The persistence of sperm and the development of time since intercourse (TSI) guidelines in sexual assault cases at Forensic Science Ireland, Dublin, Ireland. J Forensic Sci. 2017;62(3):585-92.
25. Collins KA, Bennett AT. Persistence of spermatozoa and prostatic acid phosphatase in specimens from deceased individuals during varied postmortem intervals. Am J Forensic Med Pathol. 2001;22(3):228-32.
26. Roewer L. Y chromosome STR typing in crime casework. Forensic Sci Med Pathol. 2009;5(2):77-84.
27. Bohnert M, Werner CR, Pollak S. Problems associated with the diagnosis of vitality in burned bodies. Forensic Sci Int. 2003;135(3):197-205.
28. Abdel Hady RH, Thabet HZ, Ebrahem NE, Yassa HA. Thermal effects on DNA degradation in blood and seminal stains: forensic view. Acad Forensic Pathol. 2021;11(1):7-23. doi:10.1177/1925362121998547
29. Gerling I, Meissner C, Reiter A, Oehmichen M. Death from thermal effects and burns. Forensic Sci Int. 2001;115(1-2):33-41.
30. Mishori R, Ferdowsian H, Naimer K, Volpellier M, McHale T. The little tissue that couldn't – dispelling myths about the hymen's role in determining sexual history and assault. Reprod Health [Internet]. 2019;16(1). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6547601/
31. Sawyer Sommers M. Defining patterns of genital injury from sexual assault. Trauma Violence Abuse. 2007;8(3):270-80.
32. Basile KC, Smith SG, Chen J, Zwald M. Chronic diseases, health conditions, and other impacts associated with rape victimization of U.S. women. J Interpers Violence. 2020;36(23-24):088626051990033.
33. World Health Organization. Guidelines for medico-legal care for victims of sexual violence [Internet]. Geneva: World Health Organization; 2003. Available from: https://iris.who.int/bitstream/handle/10665/42788/924154628X.pdf
34. Magalhães T, Dinis-Oliveira RJ, Silva B, Corte-Real F, Nuno Vieira D. Biological evidence management for DNA analysis in cases of sexual assault. ScientificWorldJournal. 2015;2015(1):1-11.