

A Case-Control Investigation to Determine the Amount of Serum Zinc in Children Who had an Acute Lower Respiratory Tract Infection

Sonali Suman¹, Sanjeev Kumar Sinha², Ashutosh Kumar Sinha³, Hemant Kumar⁴

¹Senior Resident, Upgraded Department of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India

²Assistant Professor, Upgraded Department of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India

³Assistant Professor, Department of Cardiology, Patna Medical College and Hospital, Patna, Bihar, India

⁴Associate Professor, Upgraded Department of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India

Received: 02-04-2021 / Revised: 03-05-2021 / Accepted: 27-06-2021

Corresponding author: Dr. Sanjeev Kumar Sinha

Conflict of interest: Nil

Abstract

Aim: The aim of this study to evaluate the serum zinc level in children with acute lower respiratory tract infection. **Methods:** This Case-Control study was done the Upgraded Department of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India for 14 months. Among 210 patients divided into two equal groups. 105 Children between the age of 2 months to 59 months and children with Acute Lower Respiratory Tract infection were include in case group and 105 children for control group. The details of blood investigations and imaging for confirmation of clinical diagnosis were also noted during the stay of the patient in the hospital. The serum zinc estimation was done by using colorimetric test. **Results:** The mean age of cases was 1.83 ± 1.39 yrs and that of controls was 1.89 ± 1.64 yrs. The Sex wise distribution of the cases and controls consisted of 74(59.20%) of cases being male and 51(40.80%) being female as compared to 69(55.20%) of controls being male and 56(44.80%) being female. The Mean serum zinc levels in the cases and controls, after comparison, were found to be significantly different [$p=0.0001$], with mean value for the cases being 56.06 ± 10.11 ug/dl as compared to 82.36 ± 11.26 ug/dl for the controls. A total of 25 cases and controls (23.81%) were found to have deficiency of zinc, of which majority (85.71%) were cases (normal range of 60 to 150ug/dl). Severe Pneumonia group (Mean= 41.23 ± 5.63 ug/dl) having significantly lower value than that of Pneumonia group (Mean= 60.22 ± 5.96 ug/dl). This is also reflected when we see serum zinc levels according to oxygen requirements, with cases managed on room air having mean of 63.63 ± 8.06 ug/dl, cases requiring supplemental oxygen by nasal prongs having mean of 60.74 ± 9.11 ug/dl and cases requiring mechanical ventilation having mean of 39.49 ± 8.67 ug/dl. **Conclusion:** we concluded that the zinc deficiency occurs in the majority of recurrent respiratory infection in children and therefore a decreased serum zinc level is considered an additional risk factor for recurrent respiratory infection.

Keywords: Zinc Level, Children, Lower Respiratory Tract Infection.

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

According to WHO, annually 4.1 million deaths occur worldwide due to acute respiratory infections (ARIs), with 90% being due to acute pneumonia. Specifically, 1.9 million of them are children younger than five years old[1], mostly related to developing countries because of malnutrition.[2] Meanwhile, lower respiratory tract infections and especially pneumonia constitute around 20% of the causes of pediatric mortality; per every 1000 live children born in developing countries, 12-20 children die before the age of five because of pneumonia.[1,3] By definition, lung inflammation is called pneumonitis, and if the cause of this inflammation is a microbial agent, it is called pneumonia. Microbial agents can include bacterial, viral, or parasitic. According to WHO, clinically, pneumonia involves acute cough attacks with or without fever associated with respiratory problems or tachypnea.[4,5] According to this definition, fever and tachypnea have high sensitivity and low specificity, while auscultating pulmonary crackles or pleural pain have high specificity and low sensitivity in diagnosing pneumonia.[6]

Zinc is an essential mineral that is involved in numerous aspects of cellular metabolism. It is required for the catalytic activity of approximately 100 enzymes[7,8] and it plays a role in immune function,[9,10] protein synthesis,[10] wound healing,[11] DNA synthesis,[8,10] and cell division.[10] It is required for maintaining intestinal cells, bone growth and immune function. It is second to iron as the most abundant trace element in the body. Zinc deficient children are at increased risk of restricted growth and developing diarrheal diseases and respiratory tract infections. Zinc is thought to decrease susceptibility to Acute Lower

Respiratory Tract Infection (ALRTI) by regulating various immune functions including protecting the health and integrity of respiratory cells during lung inflammation and injury. Supplementation of zinc could reduce the risk of pneumonia and the risk and duration of diarrhea, dysentery and malaria deaths among all infectious diseases, and they accounted for 3.9 million deaths worldwide.[11] According WHO estimates respiratory infection cause about 987,000 deaths in India of which 969,000 are LRTI.[12] ALRTI are the leading cause of mortality and a common cause of morbidity in children below five years of age. Most of these deaths are caused by pneumonia and bronchiolitis. Pneumonia kills more children each year than AIDS, malaria or measles combined with more than 2 million deaths per year.[13] The need for the study was to establish that zinc deficiency may lead to LRTI. The study was Investigating the effect of prescribing zinc sulphate on improving the clinical symptoms of pneumonia in 2 to 59 months of children's.

Material and methods

This Case-Control study was done the Upgraded Department of Paediatrics, Patna Medical College and Hospital, Patna, Bihar, India for 14 months, after taking the approval of the protocol review committee and institutional ethics committee. Among 210 patients divided into two equal groups (105 cases and 105 controls) were included in the study. 105 Children between the age of 2 months to 59 months and children with Acute Lower Respiratory Tract infection were include in case group. Children suffering from Acute Gastroenteritis or diarrheal illness, reactive airway disease/asthma or with underlying

chronic illnesses and congenital heart disease were excluded from this study.

Methodology

The detailed demographic information, history, clinical findings, laboratory findings and details of clinical course of cases and controls included in the study were entered in predesigned and validated proforma. Socio-economic status was assessed according to the Modified Kuppaswamy scale updated in 2017.[14] Detailed General examination was carried out in the patients along with Respiratory system and other systemic examination and a clinical diagnosis was made and entered into the proforma. The details of blood investigations and imaging for confirmation of clinical diagnosis were also noted during the stay of the patient in the hospital. The serum zinc estimation was done by using colorimetric test. The kit used for this study was manufactured by Centromic GMBH, Germany. The Sample used was serum obtained by centrifugation of 2 ml of blood sample collected at 3000 rpm for 3 to 5 minutes. The blood sample was obtained at Day 1 of admission of cases and controls. In two different ependoff tubes, 1000 ul of reagent in both along with 50 ul of serum in one tube and standard solution in other were mixed and incubated at 37⁰ for 5 minutes. Absorption of the standard A (Standard) and the sample A (Sample) was measured against the reagent blank A

(Blank) via the spectrophotometer at 560 nm wavelength, which was directly proportional to the concentration of total zinc in the sample.[15] Apart from measuring the serum zinc levels, the details of clinical course of the cases were also documented in terms of the duration of stay, oxygen requirements, severity of disease according to WHO IMNCI grading 2014 and outcome of the cases.

Statistical analysis

The data obtained from the cases and controls was compiled and entered into Microsoft Excel case sheet. Statistical analysis was done by using descriptive and inferential statistics using chi square test, students unpaired t-test one way ANOVA and Pearson's correlation coefficient and software used in the analysis were SPSS 21. version and p<0.05 were considered as level of significance.

Results

The mean age of cases was 1.83±1.39yrs and that of controls was 1.89±1.64 yrs. The Sex wise distribution of the cases and controls consisted of 74(59.20%) of cases being male and 51(40.80%) being female as compared to 69(55.20%) of controls being male and 56(44.80%) being female. On comparison, the distribution of cases and controls in this study according to age, sex, nutritional status and socioeconomic status was statistically not significant. (table.1)

Table 1: Demographic profile of children

Parameter	Cases=105	Controls=105
Age(mean)	1.83±1.39yrs	1.89±1.64 yrs
Sex		
Male	65(61.90%)	57(54.29%)
Female	40(38.10%)	48(45.71%)

The Mean serum zinc levels in the cases and controls, after comparison, were found to be significantly different [p=0.0001], with mean value for the cases being 56.06 ± 10.11 ug/dl as compared to 82.36 ± 11.26 ug/dl for the controls (Table 1). A total of 25 cases and controls (23.81%) were found to have deficiency of zinc, of which majority (85.71%) were cases (normal range of 60 to 150 ug/dl) (Table 2).

Table 2: Comparison of Zinc level in cases and controls

Group	N	Mean(ug/ dl)	Std. Deviation (ug/ dl)	Std. Error Mean	t-value
Cases	105	56.06	10.11	1.69	10.59, p=0.0001,S
Controls	105	82.36	11.26	2.55	

Table 3 shows comparison of serum zinc levels according to the clinical characteristics of cases. Here, the difference in mean serum zinc levels of cases according to WHO IMNCI grading was statistically significant (p value=0.0001) with cases belonging to Severe Pneumonia group (Mean=41.23 ± 5.63 ug/dl) having significantly lower value than that of Pneumonia group (Mean=60.22 ± 5.96 ug/dl). This is also reflected when we see serum zinc levels according to oxygen requirements, with

cases managed on room air having mean of 63.63 ± 8.06 ug/dl, cases requiring supplemental oxygen by nasal prongs having mean of 60.74 ± 9.11 ug/dl and cases requiring mechanical ventilation having mean of 39.49 ± 8.67 ug/dl (Table 3). The serum zinc analysis of patients according to outcome shows significantly lower zinc values (p value=0.0001) in cases who eventually died due to the ALRTI and its complications (n=11) as compared to those who got discharged after treatment (n=94) (Table 3).

Table 3: Zinc level according to clinical characteristics in cases

IMNCI Grading	No of cases	Mean(ug/dl)	SD	t-value
Pneumonia	77(73.33%)	60.22	5.96	8.98 p=0.0001,S
Severe Pneumonia	28(26.67%)	41.23	5.63	
Total	105(100%)	55.77	10.69	
O2 Requirement				
Room Air	43(40.95%)	63.63	8.06	33.98 p=0.0001,S
Supplemental Oxygen	23(21.90%)	60.74	9.11	
Mechanical Ventilation	39(37.14%)	39.49	8.67	
Outcome				
Discharge	94(89.52%)	59.87	10.87	38.97 p=0.0001,S
Death	11(10.48%)	40.26	6.39	

The mean serum zinc level was found to have a negative correlation ($r' = -0.055$) with the duration of stay of cases, however, this correlation was statistically not significant (p value = 0.598).

Discussion

Zinc plays an important role in maintaining a normal immune function and participates in all major biochemical pathways. It plays multiple roles in the perpetuation of genetic material and cellular division. Studies have suggested that zinc deficiency impairs immunocompetence with reduced cell-mediated immune responses, decreased T-

lymphocytes, abnormal T-helper and/or suppressor functions, impaired macrophage function, reduced killer cells and antibody dependent cytotoxicity.[16] Zinc supplementation in children causes an increase in the levels of complement in the blood that modulate the function of monocytes, macrophages and neutrophils polymorphs. It also helps in the development and activation of T-lymphocytes. When zinc supplements are given to individuals with low levels of zinc, the numbers of T-cell lymphocytes circulating in the blood increase and the ability of lymphocytes to fight against infection improves.[17] Meeks-Gardner J

et al have shown a positive Zinc supplementation in these patients.[18] The mean serum zinc levels were comparable to that found in the study by Hussain et al.[19] A study in Bangladesh by Shakur et al. and a study in Egypt by Rady et al. showed mean serum zinc levels in cases to be higher than this study.[20,21] On the other hand, a study done by Ibraheem et al. in Nigeria showed mean serum zinc levels of cases to be lower than this study[22] This variation of mean zinc values can be ascribed to the dietary habits of the country and nutritional status of the subjects of the study as a whole.[20-22] The difference in serum zinc levels of the cases and controls in this study, as well as in the above mentioned studies is statistically significant (p value=0.0001). Study by Kumar et al. in India and Arica et al in Greece also showed similar results.[23,24] These finding could be explained by the fact that serum zinc level is decreased by interleukins and tumour necrosis factor alfa as a part of a cute phase reaction in response to inflammatory stimulus.²³ The difference in mean serum zinc levels of cases according to WHO IMNCI grading was statistically significant (p value = 0.0001) with cases belonging to Severe Pneumonia group (Mean=41.23 \pm 5.63 ug/dl) having significantly lower value than that of Pneumonia group (Mean=60.22 \pm 5.96 ug/dl) and similar findings were seen in study by Rady et al., Hussain et al. and Brooks et.al.[19,21,25] This may be due to the fact that in zinc deficiency, there is loss of immunomodulatory effect of zinc causing unregulated immune response in the respiratory tract, leading to increased airway injury.[25] However, evidence to the contrary was found in studies by Bose et al. and Valentinier - Branth et al.[26,27] Argument has been put by the above studies that as zinc is required to mount a better immune response by the host against infection, there will be increased damage to the respiratory epithelium due to the increased immune response and thus leading to worsening of symptoms.[27]

Regarding the duration of stay of cases, Basnet et al. also found lower duration of stay in zinc supplemented group as compared to placebo, but similar to our study, the difference was statistically not significant. However, Brooks et al., Singh et al. and Malik et al. found significant reduction in duration of stay of patients of ALRTI after supplementation of Zinc.[25,28,29] Meanwhile, Bose et al., Valentinier-Branth et al. and Yuan et al found the supplementation of zinc either had no benefit or increased the duration of stay of patients of ALRTI.[26,27,30] A similar trend is also seen while evaluating the patients in terms of oxygen requirement during treatment. In this study with cases managed on room air having mean of 63.63 \pm 8.06 ug/dl, cases requiring supplemental oxygen by nasal prongs having mean of 60.74 \pm 9.11 ug/dl and cases requiring mechanical ventilation having mean of 39.49 \pm 8.67 ug/dl. While studies by Rady et al. and Brooks et al. concur with the findings of our study, studies by Bose et al. and Valentinier-Branth et al. have found no significant reduction of oxygen requirement.[21,25-27] When comparing the outcome of cases according to serum zinc levels, the findings of our study were in concordance with Rady et al., Brooks et al. and Basnet et al.[21,25,31]Also, a large systematic review of zinc supplementation by Mayo-Wilson et al. found that giving children zinc supplements might reduce their risk of death in general, and their risk of death due to lower respiratory tract infection.

Conclusion

The present investigation concluded that the Zinc supplementation had no overall effect on the duration of hospitalization or of clinical signs associated with severe infection in young children hospitalized for severe pneumonia in southern India. This finding differs from the results of 2 previously reported trials wherein zinc supplementation was associated with a shorter period of recovery from severe pneumonia. Given the conflicting results,

further research in representative settings is required to help clarify the role of zinc in the treatment of severe pneumonia.

Reference

- Williams, B.G., Gouws, E., Boschi-Pinto, C, Bryce, J, and Dye, C. Estimates of world-wide distribution of child deaths from acute respiratory infections *Lancet Infect Dis* 2002;2:25–32.
- Zar HJ Pneumonia in HIV-infected and HIVuninfected children in developing countries: Epidemiology, clinical features, and management. *Curr Opin Pulm Med* 2004;10: 176–82.
- UNICEF State of the World's Children New York: 2007, United Nations Children's Fund.
- Karimi A , Kadivar MR , Fararoe M , Alborzi A Active case finding of communicable disease in the south of the Islamic republic of Iran Eastern Mediterranean Health Journal 2000(6): 2&3 .
- WHO Program for Control of Acute Respiratory Infections. 1990 Acute respiratory infections in children: case management in small hospitals in developing countries A manual for doctors and other senior health workers WHO Geneva, Switzerland.p. 74.
- McIntosh K Community-acquired pneumonia in children *N Engl J Med* 2002; 346:4.
- Institute of Medicine, Food and Nutrition Board. Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Washington, DC: National Academy Press, 2001.
- Solomons NW. Mild human zinc deficiency produces an imbalance between cell-mediated and humoral immunity. *Nutr Rev.* 1998; 56: 27-28.
- Prasad AS. Zinc: an overview. *Nutrition.* 1995; 11: 93-99.
- Burtis CA. Tietz fundamentals of clinical chemistry.5th ed. Philadelphia: W.B. Saunders; 2008. chapter30; 1137-1141.
- Park K. Park's Textbook of Preventive and Social Medicine. 20th ed. M/s Banarsidas Bhanot Publishers, India; 2005. Epidemiology of communicable disease; chapter-5; 131-313.
- Chakama T, Singh SB, Tiwary RS. Acute Lower respiratory tract infections incidence and magnitude. *Indian pediatric.*1999; 28: 42-44.
- Singh T et al. Socio-economic status scales updated for 2017. *Int J Res Med Sci.* 2017 ; 5(7):3264-3267
- Johnsen Ø, Eliasson R. Evaluation of a commercially available kit for the colorimetric determination of zinc in human seminal plasma. *International Journal of Andrology.* 1987; 10(2):435 40.
- De Raeve HR, Thunnissen FB, Kaneko FT, et al. Decreased Cu, Zn-SOD activity in asthmatic airway epithelium correction by inhaled corticosteroid in vivo. *Am J Physiol Lung Cell Mol Physiol.* 1997; 272: 148-154.
- Ravaglia G, Forti P, Maioli F, Bastagli L, Facchini A, Mariani E, et al. Effect of micronutrient status on natural killer cell immune function in healthy free-living subjects aged ≥ 90 y. *American Journal of Clinical Nutrition* 2000;71(2):590-8
- Fraker PG, King LE, Gravy BA. The immunopathology of zinc deficiency in humans and rodents: a possible role for programmed cell death. *Nutrition and Immunology.* Klurfeld DM. New York, NY, 1993.
- Hussain AM, Saldanha PR, Sharma D et al. Estimation of Zinc Levels in Children with Lower Respiratory Tract Infections: A Prospective Observational Study from India.

- Pediatrics and Neonatal Nursing - Open Journal. 2016; 2(3):91-8.
19. Shakur S, Malek MA, Bano N, Islam K. Zinc Status in Well Nourished Bangladeshi Children Suffering from Acute Lower Respiratory Infection. *INDIAN PEDIATRICS*. 2004; 4.
 20. Rady HI, Rabie WA, Rasslan HA, El Ayadi AA. Blood zinc levels in children hospitalized with pneumonia: A cross sectional study. *Egyptian Journal of Chest Diseases and Tuberculosis*. 2013; 62(4):697-700.
 21. Rasheedat Mobolaji Ibraheem, AbdulWahab Babatunde Rotimi Johnson, Aishatu Ahmed Abdulkarim, Sikiru A. Biliaminu. Serum zinc levels in hospitalized children with acute lower respiratory infections in the north-central region of Nigeria. *African Health Sciences* 2014; 14(1): 136-142.
 22. Kumar S, Awasthi S, Jain A, Srivastava R.C. et al. Blood Zinc Levels in Children Hospitalized with Severe Pneumonia: A Case Control Study. *Indian Pediatrics* Volume 41, 2004
 23. Arica S, Arica V, Dag H, Kaya A, Hatipoglu S, Fenercioglu A, et al. serum zinc levels in children of 0-24 months diagnosed with pneumonia admitted to our clinic. *Int J Clin Exp Med*. 2011; 4(3):227-33.
 24. Brooks WA, Yunus M, Santosham M, Wahed M, Nahar K, Yeasmin S, et al. Zinc for severe pneumonia in very young children: double-blind placebo-controlled trial. *The Lancet*. 2004; 363(9422):1683-8
 25. Bose A, Coles CL, Gunavathi, John H, Moses P, Raghupathy P, et al. Efficacy of zinc in the treatment of severe pneumonia in hospitalized children <2 y old. *The American Journal of Clinical Nutrition*. 2006; 83(5):1089-96.
 26. Valentiner-Branth P, Shrestha PS, Chandyo RK, Mathisen M, Basnet S, Bhandari N, et al. A randomized controlled trial of the effect of zinc as adjuvant therapy in children 2-35 mo of age with severe or nonsevere pneumonia in Bhaktapur, Nepal. *The American Journal of Clinical Nutrition*. 2010 Jun 1; 91(6):1667-74.
 27. Singh AK, Sultan MA. Comparing the Effects of Zinc Supplementation as Adjunct to the Conventional Therapy and Placebo on Morbidity in Children with Pneumonia between Ages 1 Year to 5 Years. *Journal of Pediatric Care* 2017 May 31
 28. Malik A, Taneja DK, Devasenapathy N, Rajeshwari K. Zinc Supplementation for Prevention of Acute Respiratory Infections in Infants: A Randomized Controlled Trial. *Indian Pediatrics*. 2014;51
 29. Yuan X, Qian S-Y, Li Z, Zhang Z-Z. Effect of zinc supplementation on infants with severe pneumonia. *World Journal of Pediatrics*. 2016; 12(2):166-9.
 30. Basnet S, Shrestha PS, Sharma A, Mathisen M, Prasai R, Bhandari N, et al. A Randomized Controlled Trial of Zinc as Adjuvant Therapy for Severe Pneumonia in Young Children. *Pediatrics*. 2012; 129(4):701-8.
 31. Mayo-Wilson E, Junior JA, Imdad A, Dean S, Chan XHS, Chan ES, Jaswal A, Bhutta ZA. Zinc supplementation for preventing mortality, morbidity, and growth failure in children aged 6 months to 12 years of age. *Cochrane Database of Systematic Reviews* 2014:5. Art. No.: CD009384