

A Retrospective Research to Determine the Incidence of Hypocalcaemia in Patients Receiving Complete Thyroidectomy for Papillary Cancer

Pankaj Kumar¹, Kumar Ratnesh², Anamica Arya³

¹Assistant Professor, Department of General Surgery, J.L.N.M.C.H, Bhagalpur, Bihar, India

²Associate Professor, Department of General Surgery, J.L.N.M.C.H, Bhagalpur, Bihar, India

³Medical Officer, Department of General Medicine, J.L.N.M.C.H, Bhagalpur, Bihar, India

Received: 07-05-2021 / Revised: 10-06-2021 / Accepted: 20-09-2021

Corresponding author: Dr. Pankaj Kumar

Conflict of interest: Nil

Abstract

Aim: Study of incidence of hypocalcaemia in patients undergoing total thyroidectomy for papillary carcinoma. **Methods:** The prospective study was conducted in the Department of general surgery J.L.N.M.C.H. Bhagalpur, Bihar, India for 12 months. A total 30 patients who underwent bilateral thyroidectomy for papillary carcinoma were analysed. Data was collected from the patients undergoing total thyroidectomy by meticulous history taking, careful clinical examination, appropriate radiological, haematological investigations including serum calcium and serum albumin, operative findings and follow-up of the cases will be done after surgery for post-operative hypocalcaemia. **Results:** In this study 30 patients underwent total thyroidectomy for papillary carcinoma were studied. Patients with altered calcium before surgery were excluded. 27 female patients and 3 males' patients were included. According to age distribution higher incidence of hypocalcemia occurred in patients between 20 to 35 years. During surgery, all precautions were taken recurrent laryngeal nerve and parathyroid glands. Recurrent laryngeal nerve was identified in 12 (93.33%) patients on right side and 13 (96.67%) patients on left side. In post-operative period, all the patients were thoroughly examined and evaluated for signs and symptoms of hypocalcemia like paresthesia's, carpopedal spasm, convulsions, Chvostek's sign and Trousseau's sign. Post-operative progress of the patients showed that paresthesia was observed in 18 (60.0%) patients, Chvostek's sign in 10 (33.33%), Trousseau's sign in 3 (10%) and convulsion in one patient. In none of the patients, authors found carpopedal spasm. **Conclusion:** Post-operative hypocalcemia remains a frequent complication which can be avoided by identification of risk factors, by following proper meticulous surgical technique.

Keywords: hypocalcemia, risk factors, thyroidectomy

This is an Open Access article that uses a fund-ing model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Papillary thyroid carcinoma (PTC) is a type of well-differentiated thyroid carcinoma (wDTC), along with follicular and Hürthle cell carcinoma. However, PTC is much more common than the other two types of wDTC, accounting for 70% to 90% of all cases of thyroid malignant neoplasms.[1–2]

Patients can have seizures, laryngeal stridor, bronchospasm in acute severe condition. Cardiac arrhythmias, refractory congestive heart failure, cataract and xeroderma are the other manifestations described. Hypocalcemia caused by transient or definitive hyperparathyroidism was the most frequent complication after thyroidectomy, occurring in 63% of the cases and the nadir of hypocalcemia often appears within 48 hours after thyroidectomy it generally present on second to fifth postoperative day. The clinical feature includes circumoral paraesthesia, carpopedal spasm, and in severe cases it may lead to tetany.

However, in a few patients, hypoparathyroidism persists after 1 year and may be considered permanent. The occurrence of permanent hypocalcemia is mainly attributed to postoperative hypoparathyroidism when parathyroid glands are devascularized, injured, or dissected during the surgery present study was done to see the incidence of hypocalcemia after total thyroidectomy and to analyse the factors which might play role in its development. Post-operative hypocalcemia requires administration of calcium and vitamin D supplementation along with monitoring of blood calcium levels.[3-5]

Material and Methods

The prospective study was conducted in the Department of general surgery J.L.N.M.C.H. Bhagalpur, Bihar, India for 12 months, after taking the approval of the protocol review committee and institutional ethics committee.

Methodology

A total 30 patients who underwent bilateral thyroidectomy for papillary carcinoma were analysed. Thirty patients were enrolled in study who qualified the inclusion criteria i.e. patients more than 13 years of age including both genders admitted and diagnosed as having thyroid swelling requiring thyroid surgery. Patients less than 12 years of age, previous thyroid operation or irradiation, concomitant parathyroid disease, hypocalcaemia patients, patients already on calcium supplementation were excluded from the study. An informed consent was taken from all patients/guardians for inclusion in study. Data was collected from the patients undergoing total thyroidectomy by meticulous history taking, careful clinical examination, appropriate radiological, haematological investigations including serum calcium and serum albumin, operative findings and follow-up of the cases will be done after surgery for post-operative hypocalcaemia. Duration of operation and detailed operative findings like site and size of specimen, anatomy of parathyroid glands, anatomy of laryngeal nerves and relation of goitre with regional structures like trachea, esophagus etc. were noted. In post-operative period, all the patients were assessed for any symptoms and signs of hypocalcemia. Serial monitoring of serum calcium levels in pre-op period, immediate and early post-operative period i.e.; immediately after the surgery, 4 hours after surgery and 24 hours after surgery and on 5th post-operative day was done. All these findings were entered into predesigned patient proforma and results were analysed

Results

In this study 30 patients underwent total thyroidectomy for papillary carcinoma were studied. Patients with altered calcium before surgery were excluded. 27 female patients and 3 males' patients were included (Table 1).

Table 1: Description of study patients

Sex distribution	Number of patients
Male	3
Female	27
Total	30

Table 2: Age distribution of study patients

Age distribution (years)	Number of patients
12-30	5
31-40	13
41-50	7
>50	5
Total	30

According to age distribution higher incidence of hypocalcemia occurred in patients between 20 to 35 years. During surgery, all precautions were taken recurrent laryngeal nerve and parathyroid glands. Recurrent laryngeal nerve was identified in 12 (93.33%) patients on right side and 13 (96.67%) patients on left side. In post-operative period, all the patients were thoroughly examined and evaluated

for signs and symptoms of hypocalcemia like paresthesia's, carpopedal spasm, convulsions, Chvostek's sign and Trousseau's sign.[7] Post-operative progress of the patients showed that paresthesia was observed in 18 (60.0%) patients, Chvostek's sign in 10 (33.33%), Trousseau's sign in 3 (10%) and convulsion in one patient. In none of the patients, authors found carpopedal spasm.

Table 3: Post-operative evaluation of hypocalcemia

Post-operative progress	No. of patients	Percentage (%)
Paraesthesia	18	60
Carpopedal spasm	0	0
Convulsions	1	3.33
Chevostek's sign	10	33.33

Table 4: Incidence of hypocalcemia in different thyroid conditions

Severity of hypocalcemia	No. of patients	Incidence (%)
Severe	6	20
Mild to moderate	15	50
Normocalcemic	9	30

Discussion

In the present era mortality after thyroid surgery has decreased drastically because of expertise of surgeons and better surgical techniques. But post-operative complications specially hypocalcemia remains a fairly prevalent complication in patients undergoing total thyroidectomy and central lymph node dissection.

In the present study, recurrent laryngeal nerve (RLN) was tried to identify in all the patients. In 12 patients (93.33%) it was identified on left side and in 13 patients (96.67%) it was identified on right side. Zakaria et al in their study identified recurrent laryngeal nerve in 67.35% patients.[8]

The identification of recurrent laryngeal nerve by the surgeon during the surgery

minimizes the risk of damage to the nerve during surgical dissection. Recurrent laryngeal nerve injury after thyroidectomy can jeopardize the quality of life. In addition to the hoarseness of voice that occurs with unilateral recurrent laryngeal nerve injury, bilateral nerve injury leads to dyspnea and often life-threatening glottal obstruction. In this surgeon identified all parathyroid glands in 46.6% patients, three parathyroid glands in 23.3% patients, two parathyroid glands in 23.3% patients and one parathyroid gland in 6.8% patients. Nair et al in their study could identify all parathyroid gland in 80.14% patients and three parathyroid glands in 19.8% patients.[9] Eismontas et al in their study identified all parathyroid glands in 21.3% patients, three parathyroid in 52% patients, two parathyroid in 24.2% and single parathyroid gland in only 2% patients. The greater the number of PGs found during the surgery, lower the chance of hypocalcemia. The study by Thomusch et al determined that permanent post-operative hypocalcemia was more likely to developed if less than two PGs were found during surgery.[10] On other hand, some studies have found that more PGs found during the surgery may be associated with temporary hypocalcemia.

The prevalence of post-operative hypocalcemia following thyroidectomy which may be temporary or permanent ranges from 0% to 83%, with the highest incidence seen in patients undergoing total thyroidectomy for papillary carcinoma (70%). Decreased serum calcium, secondary to hypoparathyroidism, may present clinically with muscle cramps, perioral and peripheral paresthesia's, tetany and confusion.

Management protocol was as follows- (a) every total thyroidectomy patient or completion thyroidectomy patient is started on 3 g of elemental calcium, p.o., per day. This should begin as soon as the patient can take p.o. unless there is a specific contraindication to oral calcium in the

patient; (b) check ionized calcium q8 hours post-op; (c) if two consecutive calcium values (within the normal range) are stable or increasing, discontinue checking and patient is tapered off calcium supplementation. Regimen as follows: 1 g elemental calcium TID for 1 week, 1 g elemental calcium BID for 1 week, 1 g elemental calcium Q-day until RTC attending. Of note 2.5 g of TUMS (calcium carbonate) is equivalent to 1 g of elemental calcium. If calcium is decreasing, increase oral calcium to 4 g elemental per day. If this stabilizes the calcium, arrangements for the above taper regimen beginning at 1 g QID and tapering down by one gm per week are made at discharge. Check and correct abnormal magnesium. The patient should have calcium checked 1 week after discharge (locally or at UIHC follow-up); (d) every total thyroidectomy/completion thyroidectomy patient started on Os- Cal with vitamin D (500 mg CaCO₃- 200 IU vitamin D₃ per tab), 2 tabs (total of 1000 mg CaCO₃) TID taper over 3 weeks- 2 tabs TID for one week, then 2 tabs BID for one week, then 2 tabs daily for one week, then off; (e) if the patient requires 1,25-dihydroxy vitamin D, the patient should be sent home on the dose of oral calcium and vitamin D that stabilized the ionized calcium; (f) for patients that are severely symptomatic with last evaluated ionized calcium below 3.8 or for those with a calcium below 3.2, IV calcium may be administered ONE amp of calcium gluconate (10 ml calcium gluconate 10% contains 1 g calcium gluconate) in 500 ml of D5W is given IV over 5 hours. Be sure that the IV is functioning well before administration of the calcium. This infusion will usually stop the symptoms. Recheck calcium after the administration. If necessary, this may be repeated. Recheck or check the magnesium level and correct if needed.

Conclusion

Post-operative hypocalcemia remains a frequent complication which can be

avoided by identification of risk factors, by following proper meticulous surgical technique. The estimation of serial serum calcium levels is fairly predictable and cost-effective method for diagnosis of early post-operative hypocalcemia and prediction of patients who are at higher risk of developing long-term hypocalcemia after thyroid surgery. It aids in early management with calcium and vitamin supplements reducing the morbidity involving total thyroidectomy.

Reference

1. Haymart MR, Esfandiari NH, Stang MT, Sosa JA. Controversies in the management of low-risk differentiated thyroid cancer. *Endocr Rev.* 2017;38(4):351–78. doi:10.1210/er.2017-00067
2. Dralle H, Musholt TJ, Schabram J, Steinmüller T, Frilling A, Simon D, et al. German Association of Endocrine Surgeons practice guideline for the surgical management of malignant thyroid tumors. *Langenbeck's Arch Surg.* 2013;398(3):347–75.
3. Roh JL, Park CI. Routine oral calcium and vitamin D supplements for prevention of hypocalcemia after total thyroidectomy. *Am J Surg.* 2006;192(5):675-8.
4. Abboud B, Sargi Z, Akkam M, Sleilaty F. Risk factors for postthyroidectomy hypocalcemia. *J Am Coll Surg.* 2002;195(4):456-61.
5. Thomusch O, Machens A, Sekulla C, Ukkat J, Lippert H, Gastinger I, Dralle H. Multivariate analysis of risk factors for postoperative complications in benign goiter surgery: prospective multicenter study in Germany. *World J Surg.* 2000;24(11):1335-41.
6. Henry CR, Speroff T, Wentworth D, Murphy T. Risk factors for postthyroidectomy hypocalcemia. *Surgery.* 1994;116(4):641-7.
7. Baldassarre RL, Chang DC, Brumund KT, Bouvet M. Predictors of hypocalcemia after thyroidectomy: results from the nationwide inpatient sample. *ISRN Surg.* 2012;838614.
8. Zakaria HM, Awad NA, Kreedes AS, Mulhim AM, Sharway MA, Hadi MA, et al. Recurrent laryngeal nerve injury in thyroid surgery. *Oman Med J.* 2011;26(1):34-8.
9. Eismontas V, Slepavicius A, Janusonis V, Zeromskas P, Beisa V, Strupas K, et al. Predictors of postoperative hypocalcemia occurring after a total thyroidectomy: results of prospective multicenter study. *BMC Surg.* 2018;18(1):55.
10. Thomusch O, Machens A, Sekulla C, Ukkat J, Brauckhoff M, Dralle H. The impact of surgical technique on postoperative hypoparathyroidism in bilateral thyroid surgery: a multivariate analysis of 5846 consecutive patients. *Surgery.* 2003;133:180-5.
11. Arumugam S, Mohankumar A, Muthukumaraswamy A, Anandan H. Clinical study of hypocalcemia following thyroid surgery. *Int J Sci Stud.* 2017;4:37-41.