

A Comparative Study of Laparoscopic Appendicectomy Specimen Retrieval Using Plastic Endo-Bag and Direct Removal Through 10mm Umbilical Port

S. Mamatha¹, G. Bharath², N. Venkata Harish³, V. Mahidhar Reddy⁴

¹Junior Resident, Department of General Surgery, Narayana Medical College, Nellore, Andhra Pradesh India

²Assistant Professor, Department of General Surgery, Narayana Medical College, Nellore, Andhra Pradesh India

³Associate Professor, Department of General Surgery, Narayana Medical College, Nellore, Andhra Pradesh India

⁴Professor, Department of General Surgery, Narayana Medical College, Nellore, Andhra Pradesh India

Received: 25-07-2025 / Revised: 23-08-2025 / Accepted: 26-09-2025

Corresponding Author: Dr. Mamatha Suraneni

Conflict of interest: Nil

Abstract:

Background: Laparoscopic appendectomy is the preferred approach for treating acute appendicitis. Specimen retrieval techniques may influence operative times and postoperative outcomes.

Objective: To compare the operative outcomes between Plastic Endo-Bag retrieval and Direct 10mm Umbilical Port retrieval during laparoscopic appendectomy.

Materials and Methods: In this prospective observational study conducted over 18 months, 50 patients diagnosed with acute appendicitis were alternatively allocated into two groups: Group A (Endo-Bag retrieval) and Group B (Direct retrieval). Mean operative time, specimen retrieval time, first passage of flatus, length of hospital stay, and port-site infections were evaluated.

Results: Mean operation and specimen retrieval times were significantly longer in Group A compared to Group B ($p < 0.05$). Port-site infections were slightly more common in Group B but without statistical significance. No differences were observed regarding the first passage of flatus and hospital stay.

Conclusion: Direct retrieval through a 10mm umbilical port reduces operative time and specimen retrieval time without significantly affecting infection rates. Larger studies are recommended to confirm these findings.

Keywords: Laparoscopic Appendicectomy, Endobag, Specimen Retrieval.

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

Acute appendicitis is a common surgical emergency worldwide.

Laparoscopic appendicectomy has become the standard treatment due to advantages such as reduced postoperative pain, shorter hospital stay, and quicker recovery. Specimen retrieval during laparoscopic appendicectomy can be challenging, espe-

cially with inflamed appendices. Retrieval via a plastic endo-bag is thought to reduce contamination, but direct retrieval through a 10mm umbilical port is quicker and cost-effective.

This study aims to compare these two retrieval techniques in terms of operative outcomes and postoperative complications.




Figure 1:

Materials and Methods

Study Design and Setting: A prospective observational study was conducted at the Department of General Surgery, Narayana Medical College & Hospital, Nellore, Andhra Pradesh, India, from October 2022 to March 2024.

Study Population: Fifty patients clinically diagnosed with acute appendicitis and fit for laparoscopic surgery were enrolled. Patients were alternatively allocated into two groups:

- Group A: Specimen retrieval using Plastic Endo-Bag.
- Group B: Direct specimen retrieval through 10mm umbilical port.

Inclusion Criteria

- Patients with clinical diagnosis of acute appendicitis.
- Patients willing to undergo laparoscopic appendectomy.

Exclusion Criteria

- Patients diagnosed with other causes of acute

abdomen.

- Patients refusing surgery.
- Patients medically unfit for surgery.

Outcome Measures

- Mean operation time.
- Specimen retrieval time.
- First passage of flatus.
- Length of hospital stay.
- Port-site infections.

Statistical Analysis: Data were analyzed using independent samples t-test, chi-square test, and one-way ANOVA. A p-value <0.05 was considered statistically significant.

Results

Demographic Profile: The mean age of patients was 22.7 ± 9.65 years. Male and female patients were nearly equally distributed across both groups.

Operative Outcomes

Mean Operation Time: Group A: 46.25 ± 11.2 minutes; Group B: 42.5 ± 10.5 minutes ($p = 0.026$).




Figure 2:

Specimen Retrieval Time: Group A: 3.5 minutes; Group B: 3.05 minutes ($p < 0.001$).




Figure 3:

Hospital Stay: 90% of patients were discharged on day 3; 10% required longer stay due to port- site infections.




Figure 4:

First Passage of Flatus: Majority of patients in both groups passed flatus by postoperative day 1 ($p > 0.05$).




Figure 5:

Port-Site Infection: Group A: 4%; Group B: 16% ($p = 0.157$, not statistically significant).




Figure 6:

Discussion

The findings indicate that while using a Plastic Endo-Bag adds a few extra minutes to the operation and retrieval times, it does not significantly reduce port-site infections compared to direct retrieval through a 10mm umbilical port. Similar findings have been reported in previous studies, suggesting that although retrieval bags reduce contamination theoretically, their clinical advantage regarding wound infection remains debatable.

The study by Mustafa Sami Bostan et al. (2022) found that the use of specimen retrieval bags was not statistically significant in reducing surgical site infections, supporting our results.

Furthermore, most patients experienced early return of bowel function and had a short hospital stay, consistent with standard outcomes for laparoscopic appendectomy.

Conclusion

Direct specimen retrieval through a 10mm umbilical port during laparoscopic appendectomy reduces operative and extraction times without significantly increasing port-site infections.

The chances of port site infections were found more in direct retrieval technique through 10mm port, but the data was not statistically significant.

However, larger, multicentric randomized studies are warranted to validate these results.

References

1. Moore KL, Dalley II AF, Agur AMR. MOORE Clinically Oriented Anatomy. 7th ed. Lip-pincott Williams & Wilkins; 2014.
2. Standring S. Gray's Anatomy: The Anatomical Basis of Clinical Practice. 41st ed. Elsevier Limited; 2016.
3. Richmond B. Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice. 21st ed. Elsevier Inc; 2022.
4. Liang MK, Andersson RE, Jaffe BM, Berger DH. Schwartz's Principles of Surgery. 10th ed. McGraw-Hill Education; 2015.
5. Bostan MS, Ugurlu C. Is it necessary to use specimen retrieval bag for reducing surgical site infection in laparoscopic appendectomy? A randomized controlled trial. Turkish Journal of Trauma and Emergency Surgery 2022.