

A Prospective Study on Role of Plasma Fibrinogen in Diagnosis of Neonatal Sepsis and it's Prognosis

Aaisha Anjum Risaldar¹, Aryak Banerjee², Ayesha Fatima³

¹Consultant Pathologist, Shoaib Clinical Laboratory

²Assistant Professor, Dept. of Pathology, Raniganj Institute of Medical Sciences, West Bengal, India

³Associate Professor, Dept. of Pathology, Raniganj Institute of Medical Sciences, West Bengal, India

Received: 26-04-2026 / Revised: 25-05-2026 / Accepted: 21-06-2026

Corresponding Author: Dr. Ayesha Fatima

Conflict of interest: Nil

Abstract:

Purpose: To study the relation between plasma level of fibrinogen and neonatal sepsis and its outcome.

Methods: A prospective case control study was done including 48 neonatal sepsis cases in study group and 42 cases in control group. For all cases Plasma fibrinogen, prothrombin time (PT), activated plasma thrombin time (APTT), C-reactive protein (CRP) and blood culture were performed. Mean Plasma fibrinogen levels for both groups and cut off value of plasma fibrinogen for diagnosis was calculated using receiver operating characteristic curve.

Results: Plasma fibrinogen levels were found to be higher in study group with than in control group ($p < 0.050$). Cutoff value of plasma fibrinogen for diagnosing neonatal sepsis was found to be 305.5mg/dl with a sensitivity and specificity of 80% and 72.8% respectively.

Conclusion: In our study, we found that plasma fibrinogen act as an acute phase reactant and can be used as an immediate marker for detection of early onset neonatal sepsis.

Keywords: Plasma Fibrinogen, Neonatal Sepsis, Acute Phase Reactant, Complication.

DOI: 10.25258/ijpqa.17.6.53

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>) and the Budapest Open Access Initiative (<http://www.budapestopenaccessinitiative.org/read>), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

“Neonatal sepsis is a clinical syndrome resulting from the pathophysiological effect of severe bacterial infection in the first month of life.” [1] Neonatal sepsis is the most common cause of morbidity and second most common primary cause of neonatal mortality next to perinatal asphyxia. Neonatal mortality rate was 25.3 per 1000 live births as per National Neonatal Perinatal Database (NNPD) 2002-2003 Report.

Since neonatal sepsis is a preventable cause of mortality, early detection of onset of sepsis is helpful for treatment and also in preventing complications. Neonatal sepsis symptoms and signs include refusal to feed, fever, excessive cry, hypoglycemia, hypothermia, tachycardia, tachypnea, convulsions. These are not specific to neonatal sepsis and also seen in various other conditions, which pose difficulty in diagnosis. [2],[3],[4]

Blood culture, CSF, urine or other body fluids culture are gold standard methods for diagnosis of sepsis but they need at least 48 to 72 hours for reporting. [4, 5] These delays crucial periods during which antibiotics should be administered for treatment. Also there are few false negative blood culture cases present which need to be diagnosed. This necessi-

tates other alternate methods for diagnosis or other indicators for sepsis which requires less time and hence reducing the time to start the treatment.

There are few other method which are used to support the diagnosis like, leukocyte indices, absolute neutrophil count, toxic granules, nRBC's, Acute phase reactants like micro ESR, C-reactive protein, procalcitonin.

Plasma fibrinogen is also one such an acute phase reactant. Mean value of plasma fibrinogen level in healthy preterm and full-term neonates aged up to 30days range from 2.43 to 2.54g/L and 2.70 to 2.83g/L respectively[2],[6],[7]. It is seen to increase several fold in inflammatory conditions as a positive acute phase reactant.[2] It is produced in the liver as one of the coagulation pathway agent, which is needed for haemostasis. This feature is used for prediction of neonatal sepsis in this study.

This study was undertaken to study the relation between plasma levels of fibrinogen and neonatal sepsis and it's outcomes like prediction of complications in neonatal sepsis.

Materials and Methods

Neonatal sepsis cases and hyperbilirubinemia cases as controls which fulfill inclusion and exclusion criteria were collected for case control study, from hospital NICU, in department of pediatrics and department of pathology.

Data of the both study group like clinical history, antenatal history, clinical examination and demographic data were collected. Blood samples were collected in plain, EDTA and Citrate anticoagulated vacutainers from umbilical cord blood or arterial or venous blood for blood culture, CRP, analysis of hematologic parameters and coagulation profile consisting PT, aPTT and plasma fibrinogen respectively. For blood culture, sample from plain vacutainer was used. Sample blood was first inoculated in glucose broth and incubated for 48 hours and then streaked into McConkey agar for further growth. Also, blood from plain sample was used for estimation of ESR.

Blood samples collected in Citrate anticoagulated vacutainers were used for estimation of PT, aPTT and plasma fibrinogen. Sample was centrifuged for 15 minutes at 1500 rpm to obtain plasma and was run on 'Automated coagulometer ACL Elite Pro' to estimate PT and aPTT values.

Plasma fibrinogen was estimated by automated Erba Mannheim ECL 105 coagulometer by using 'Erba Thrombin reagent for fibrinogen determination' which is made up of approximately 100 NIH units/ml bovine thrombin with stabilizers and Erba Owren's veronal buffer.

Statistical Analysis: The data obtained were recorded in a Microsoft Excel sheet, and statistical analysis was performed using statistical package for the social sciences (Version 17). Results are presented as drawings, Mean $\pm$ SD, counts and percentages. Optimal cut off values of Fibrinogen and CRP were defined by Youden's index. Sensitivity and specificity of these variables were performed using ROC curves. Categorical variables were compared using chi square test, quantitative variables were compared using independent t test & Mann Whitney U test. For all tests, significant value was achieved at $p < 0.05$. All statistical tests performed were two tailed.

Inclusion Criteria: Neonatal cases with early signs of sepsis and neonates born to mothers with premature rupture of membranes and fever.

Exclusion Criteria: Neonates with congenital anomalies.

Results

In this study we found out that among 48 suspected cases of neonatal sepsis, blood culture was positive for 12 cases and positive for viral infection in 6 cases and 2 cases having co-infection of both bacteria and

virus. Out of 12 culture positive cases, 3 (25%) cases were of *E. coli*, 2 (16.66%) cases each were of *Klebsiella pneumoniae*, *Staphylococcus* and Coagulase negative *Staphylococcus*. 1 (8.34%) case each of *Staphylococcus Aureus*, *Staphylococcus pneumoniae* and *Streptococci*. Among this 1 case of CON *Staphylococcus* with CMV positive was obtained from CSF culture. 1 case was coinfection with *Staphylococcus pneumoniae*, CMV, Rubella & *Toxoplasma*. 2 cases were CMV positive, 2 cases were positive for Herpes IgG and 1 case was positive for CMV and Rubella.

There was no significant difference between gestation period and age of neonates. However, plasma fibrinogen level (p value = 0.010), CRP (p value = 0.005) and total leucocyte counts (p value = 0.008) were significantly higher in study group than control group. Values of parameters are given in table 1.

With the help of ROC curve, cut off value for plasma fibrinogen was obtained. In the ROC Curve for Blood culture, the AUROC of Fibrinogen, is 76% (95% of Confidence Interval = 62 % to 90%) and the optimal cut-off value is 305.5. Using this cut-off value, the Sensitivity and specificity are 80% and 72.8 % respectively. The ROC curve is shown in figure 1.

Discussion

The current study was undertaken to evaluate the efficacy of plasma fibrinogen in the diagnosis of neonatal sepsis and its outcome as in complications. Along with it other haematological indices and CRP are also evaluated for their efficacy.

In current study we evaluated role of plasma fibrinogen levels on neonatal sepsis and its outcome and we found out that plasma fibrinogen levels were significantly elevated in study group containing suspected cases of neonatal sepsis compared to control group. In this study mean plasma fibrinogen was $307.25 \pm \text{SD } 90.015$ for study group and $273.21 \pm \text{SD } 80.884$, which was higher statistically significant with p value of 0.010.

It was also observed that 4 cases which develop complications showed lower level of plasma fibrinogen, but PT and APTT were prolonged whereas in cases which did not develop complications, plasma levels of fibrinogen were elevated but PT and APTT were within normal range.

In current study after comparing proven neonatal sepsis cases with our control population, on analysis of ROC curves we got cut off value of 305.5 mg/dl with sensitivity and specificity of 80% and 72.8%, which was comparable to study conducted by Mitra P et al.[2] who got cut off value of 301.9 mg/dl with sensitivity and specificity of 70.8% and 82.7%.

According to study conducted by Guibourdenche J et al[4] concluded that Plasma fibrinogen acts as an

acute phase reactant. They got cut off value of 300mg/dl for Plasma fibrinogen in diagnosing neonatal sepsis which was comparable to present study.

In present study the most common organism isolated was E coli followed by Klebsiella pneumonia, staphylococcus and coagulase negative staphylococcus. Similar findings were seen in Cortese F et al[8] , Manroe et al[9] Chandna A et al[10] and Renolder B et al[11]. Most common viral infection was CMV.

In this study mean total leucocyte count was higher than control group (p value of 0.008). As per study done by Ahirrao BM et al[12]. Total WBC count has lower sensitivity but higher specificity in detecting neonatal sepsis. It is because of wide normal range of WBC count between 5000-34000cells/cumm.

Conclusion:

Plasma fibrinogen is normally present in blood as a factor of coagulation system but will be increased in concentration in any infection or inflammation and acts as positive acute phase reactant. It is a rapid, simple and cost effective test, can be used along with other hematological parameters & biochemical markers like CRP in diagnosis of neonatal sepsis. The present study concluded that plasma fibrinogen is not only immediate reliable marker in diagnosing neonatal sepsis but also aids in predicting the complications thus improving the diagnostic efficacy. So this complimentary test also helps the clinicians to start the treatment as early as possible & thereby reducing neonatal mortality, bringing a significant impact in the neonatal health care.

References:

1. Mondal S K, Nag D N, Bandyopadhyay R, Chakraborty D, Sinha S K. Neonatal sepsis: Role of a battery of immunohematological tests in early diagnosis. *Int J Appl Basic Med Res*. 2012 Jan-June 2(1):43-7.
2. Mitra P, Guha D, Nag S S, Mondal B C, Dasgupta S. Role of Plasma Fibrinogen in Diagnosis and Prediction of Short-Term Outcome of Neonatal Sepsis. *Indian J Hematol Blood Transfus*. 2017 Apr-June 33(2):195-9.
3. G Vandana, S Lokesh Rao Magar, Praveen, Kavithadevi, Sandhya Rani, Sandhya Anil. Haematological profile in neonatal sepsis. *IOSR Journal of Dental and Medical sciences*. 2017 Apr; Vol 16(4):11-17.doi:10.9790/0853-1604091117.
4. Guibourdenche J, Bedu A, Petzold L, Marchand M, Mariani-Kurdjian P, Hurtaud-Roux M F, Aujard Y, Porquet D. Biochemical markers of neonatal sepsis: value of procalcitonin in the emergency setting. *Ann Clin Biochem* 2002; 39: 130- 5.
5. Das B, Das A, Saikia M. A study of haematological parameters in neonatal septicemia. *Indian Journal of Basic and Applied medical Research*. 2016 March; vol- 5(2): 491-500.
6. Andrew M, Paes B, Milner R, Johanson M, Mitchell L, Tollefsen DM, Powers P (1987). Development of the human coagulation system in the full-term infant. *Blood*.1987 july 70(1):165-72.
7. Andrew M, Paes B, Milner R, Johanson M, Mitchell L, Tollefsen DM, Castle V, Powers P (1988). Development of the human coagulation system in the healthy preterm infant. *Blood*. 1988 nov 72(5): 1651-57.
8. Cortese F, Scicchitano P, Gesualdo M, Filaninno A, De Giorgi E, Schettini F, et al. Early and Late Infections in Newborns: Where Do We Stand? A Review. *Pediatr Neonatol*. 2016;57(4):265-73.
9. Manroe BL, Rosenfeld CR, Browne R, Weinberg AG. The differential leucocyte count assessment and outcome of early-onset neonatal Group B streptococcal disease. *J Pediatr* 1977; 91: 632- 7
10. Chandna A, Rao MN, Srinivas, Shyamala S. Rapid diagnostic tests in neonatal septicemia. *Indian J Pediatr*.1988;55:947-53
11. Renolder B, Hofer N, Resch B. Early-Onset Neonatal Sepsis: Group B Streptococcal Compared to E. coli Disease. *J Neonatal Biol*.2015; 04:78-92.
12. Ahirrao B et al. Diagnostic utility of Hematological Scoring System (HSS) with clinicopathological and bacteriological evaluation in early diagnosis of neonatal sepsis. *Annals of pathol and laboratory medicine*.2017;4(6):721-6
13. Sankar MJ, Agarwal R, Deorari AK, Paul VK. Sepsis in the newborn. *Indian Journal of Paediatrics* 2008;75(3):261-6.
14. Abhishek MG, Sanjay M. Diagnostic efficacy of Nucleated Red Cell count in the early diagnosis of neonatal sepsis. *Indian Journal of Pathology and Oncology* 2015;2(4):182-5.
15. Mishra UK, Jacobs SE, Doyle LW, Garland SM. Newer approaches to the diagnosis of early onset neonatal sepsis. *Arch Dis Child Fetal Neonatal Ed*.2006;91:F208-12
16. Chiesa C, Panero A, Rossi N, Stegagno M, De Giusti M, Osborn JF, et al. Reliability of Procalcitonin Concentrations for the Diagnosis of Sepsis in Critically Ill Neonates. *Clin Infect Dis*. 1998;26(3):664-72.
17. Chiesa C, Panero A, Osborn JF, Simonetti AF, Pacifico L. Diagnosis of neonatal sepsis: a clinical and laboratory challenge. *Clin Chem*. 2004;50: 279-87.
18. Cant A J, Gennery A R. Immunodeficiency. Rennie and Robertson's Textbook of Neonatology 5th Ed: Churchill Livingstone Elsevier 2012; p994-1011

19. Haque KN. Neonatal Sepsis in the Very Low Birth Weight Preterm Infants : Part 1: Review of Pathophysiology. 2010;3:1–10.
20. Zawar MP, Tambekar RG, Deshpande NM, Gadgil PA, Kalekar SM. Early diagnosis of neonatal septicemia by sepsis screen. Indian J Pathol Microbiol. 2003; 46(4):610-12.
21. Bhat RY, Lewis LE, Vandana KE. Bacterial isolates of early-onset neonatal sepsis and their antibiotic susceptibility pattern between 1998 and 2004: an audit from a center in India. Italian Journal of Pediatrics. 2011;32:37
22. Buch AC, Srivastava V, Kumar H and Jadhav PS. Evaluation of haematological profile in early diagnosis of clinically suspected cases of neonatal sepsis. International Journal of Basic and Applied Medical Sciences. 2011;1(1):1-6
23. Agarwal AM, Rodgers GM. Miscellaneous causes of thrombocytopenia. In Greer JP, Arber DA, Glader B, List AF, Means RT, Paraskevas S, Rodgers GM (eds) Wintrobe's Clinical Hematology. 13th Ed. Philadelphia. Williams and Wilkins.2014:1097-105.
24. Hermansen MC. Nucleated red blood cells in the fetus and newborn. Arch Dis Child Fetal Neonatal Ed. 2001;84(3):F211–5.
25. Constantino BT, Cogionis B. Nucleated RBCs - Significance in the peripheral blood film. Lab Med. 2000;31(4):223–9.
26. Kwatakar SM, Erythrocyte Sedimentation Rate, Essentials of Clinical Parhology; 1st Ed, JP; 2010; 215-9.
27. Diwakar KK. Revised look at Micro erythrocyte sedimentation rate in neonates. Indian Pediatrics 1999;36:703-5
28. Pal GK. Textbook of Medical Physiology; 3rd Ed, JP Medicals Ltd; 2016;54, 64,108,131
29. Oaei-Bimpong A, Burthem J, Supplementary Techniques Including Blood Parasites Diagnosis, Dacie and Lweis Practical haematology; 12th Ed, Elsevier; 2017;p 63-111
30. Kumar V, Abbas KA, Aster CJ, Diseases of the immune system, Robbins and Cotran: Pathologic Basis of Disease; 9th Ed. Elsevier; 2015: p 99-106.
31. Vasudevan PM, Sreekumari S, Vaidyanathan K. Textbook of biochemistry 7th Ed: JP Medicals; 2013; p 383-6
32. Devlin TM, textbook of Biochemistry with clinical correlation; 7th Ed. Wiley-Liss;2006: p987-8