e-ISSN: 0975-9506, p-ISSN: 2961-6093

Available online on www.ijpqa.com

International Journal of Pharmaceutical Quality Assurance 2026; 17(6); 346-348

Original Research Article

A Study on Effect of BMI on VO2max Assessed by Medical College Step Test in Healthy Adults Males

Amera Anjum Saifee[1], Mujtaba Nausheen[2], Banda Naveen[3]

[1]Senior Resident, Dept. of Physiology, ESIC, Medical College & Hospital, Sanathnagar, Hyderabad, Telangana, India

[2]Assistant Professor, Dept. of Physiology, Dr. Shankarrao Chavan Government Medical College, Nanded, Maharashtra, India

[3]Senior Resident, Dept of General Medicine, Raichur Institute of Medical Sciences, Raichur, Karnataka, India

Received: 20-04-2026 / Revised: 11-05-2026 / Accepted: 18-06-2026

Corresponding Author: Dr. Amera Anjum Saifee

Conflict of interest: Nil

Abstract:

Background: Cardiorespiratory fitness (CRF) is not only objective measure of habitual physical activity, but also a useful diagnostic and prognostic health indicator for patients in clinical settings. CRF is a strong and independent predictor of cardiovascular disease mortality. Low CRF in young adults has emerged as an important factor for developing cardiovascular comorbidities later in middle age.

Aims: To determine the peak oxygen consumption (VO2max) by Queen's college step test in young male subjects. To study the relation between body mass index and VO2max.

Methods: One hundred young healthy male subjects in the age group of 18 to 25 years were included in this study group. Body mass index (BMI) was measured using standard calibrated instruments. VO2max was assessed by following the protocol of Queen's College Step Test (QCT).

Results: Data was statistically analysed by using Pearson's correlation. There was a significant negative correlation between Body mass index and VO2max (ml/kg/min) ($r= -0.418$, $p<0.01$).

Conclusion: The results suggest excessive amount of body fat exerts an unfavourable burden on cardiac function and oxygen uptake by working muscles. In our study BMI has significant negative correlation with VO2max and can be used in clinical settings to estimate cardiorespiratory fitness as it is a rapid and inexpensive method.

Keywords: VO2max, QCT, BMI, CRF.

DOI: 10.25258/ijpqa.17.6.83

This is an Open Access article that uses a funding model which does not charge readers or their institutions for access and distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0) and the Budapest Open Access Initiative (http://www.budapestopenaccessinitiative.org/read), which permit unrestricted use, distribution, and reproduction in any medium, provided original work is properly credited.

Introduction

The prevalence of cardiovascular disease has increased substantially over the past few decades in younger population [1]. VO2max is internationally accepted parameter to evaluate cardiovascular fitness [2]. CRF is an important component of physical fitness, representing the body's ability to take oxygen in and deliver this oxygen to muscle cells throughout the body during physical activity. Previous studies have emphasized the importance of CRF, providing convincing evidence that CRF is closely related to all-cause mortality [3, 4]. Earlier studies have established the use of Queen's college step test to predict VO2max indirectly [5].

Obesity is considered one of the most important medical and public health problems of our time. Excess body fat has been identified as a major risk factor for several common disorders including diabetes and cardiovascular diseases and imposes a substantial burden on health care systems [6]. Obesity can be measured in various ways. Body mass index (BMI) is the most commonly used measure of relative weight. It can be used both at individual level to asses body weight in a clinical setting and at population level where it would be impractical or too expensive to measure (excess) body fat accurately and consistently [7].

Earlier studies have demonstrated the importance of low VO2max in young adulthood as a factor for developing cardiovascular comorbidities later in middle age [8]. This study is designed to evaluate VO2max and its relation with body mass index in young healthy male subjects.

Methods

The study group comprises of 100 young healthy males in the age group of 18-25 yrs. subject's clinical history and details were taken according to the standard proforma. Informed written consent was

taken from all subjects in the study. The study involved noninvasive procedures and was performed at room temperature.

Subjects with history of cardiac disease, lung disease, Smoking history, those on regular medications affecting cardiovascular and respiratory system and those who are undergoing any physical conditioning programme were excluded from the study. The experimental protocol was fully explained to the participants to allay apprehension. The subjects were refrained from any energetic physical activity for 3 hours before the test.

Results

VO2max was evaluated and then the effect of BMI on cardiorespiratory fitness was studied in 100 young healthy males. Data was statistically analysed by Pearson's correlation and results are shown in tables and graphs. The data profile of subjects including name, age, height, weight, body mass index, QCT pulse rate and VO2max with their mean and standard deviation are shown in table 1.There was a significant negative correlation between BMI and VO2max (ml/kg/min)) r= -0.626, p<0.01 (figure 1 and table 2).

Table 1: Profile of the study group

Parameters	No	Mean	Standard deviation
Age	100	19.06	1.269852
Weight	100	63.04	11.78762
Height	100	1.7047	0.062465
BMI	100	21.660	3.719306
QCT Pulse rate	100	170.04	16.08558
VO2 Max	100	39.913	6.755943

Table 2: Correlation between BMI and VO2 Max

Variable	VO2 Max
BMI (r)	**-0.626****
BMI (P)	**< 0.01***

***--Significant**

****--the correlation between BMI and VO2max is negative .i.e., as BMI increases the VO2max is decreasing and vice versa**

Discussion

VO2max is a measure of the functional limit of cardiorespiratory system and single most valid index of maximal exercise capacity. The absolute value of VO2max is one of the indices of individuals CRF to transport oxygen to working muscles. Earlier studies have used VO2max values in ml/kg/min to examine the performance of CRF.

Excessive amount of body fat exerts an unfavorable burden as well as hindering action towards cardiac function, particularly during exhaustive exercise, when excessive hyperactive body musculature fails to uptake sufficient amount of oxygen due to deposition of proportionately high amount of fat mass. Loss of weight during weight reduction program of obese increased their VO2max (ml/kg/min) due to withdrawal of fat induced inhibitory action towards oxygen utilization by body musculature [11].

Elevated myocardial oxidative stress has been reported in patients with obesity. In obese individuals there is increase in type II muscle fibers and decrease in type I muscle fibers which may have important effect on reduced oxygen uptake. Overweight subjects have much greater increase in sympathetic nerve firing rate than normal subjects. Obesity is said to produce a state of chronic volume overload because heart is required to continuously circulate blood through the large and relatively low resistance depot of adipose tissue. Increased preload and stroke volume are associated with hypertension. Combination of overweight and hypertension leads to thickening of ventricular wall and larger heart volume and thus a greater likelihood of cardiac failure [12].

Body mass index (BMI) has traditionally been used to identify individuals who are the most likely to be overweight or obese. It was developed by Belgian polymath Adolphe Quetelet in course of working out his systems of "Social Physics" between 1830 and 1850 (also known as Quetelet index). Generally, a high value indicates excessive body fat and consistently relates to increased health risks and mortality. Unusually large muscle mass, as in trained athletes, can increase BMI to 30, but rarely above 32. BMI categories and cut-offs are commonly used to guide patient management [9].

Welch et al. reported that VO2max was significantly decreased in overweight individuals when fat mass was taken into account which suggests the possibility of deconditioning and changes in cardiorespiratory function in severely overweight individuals [13]. Watanabe K et al. reported that obesity accentuates exercise intolerance and low aerobic capacity [14]. Similar results were observed by Rowland TW et al. [15] Norman et al. studied influence of excess adiposity on exercise fitness and performance in overweight children and adolescents and found that

overweight and non-over weight adolescents had similar absolute CRF but the functional impairment was significantly associated with increased energy demands needed to move their excess bodyweight [16]. Several previous studies have found no significant differences in VO2max between obese and non-obese.

Conclusions

There was a significant negative correlation between BMI and VO2max (ml/kg/min). This suggests the effect of body fat on cardiorespiratory functions. These findings demonstrate the importance of low CRF in young adults with increased body fat is a factor for developing cardiovascular comorbidities later in middle age. BMI can be used in clinical settings to estimate CRF as they are rapid and inexpensive methods. Given the current obesity trend and observations of a decline in daily energy expenditure among the people, improving peak oxygen consumption in young men by engaging in physical activities is important.

References

1. Buffart, L. M., Roebroeck, M. E., Rol, M., Stam, H. J., & van den Berg-Emons, R. J. (2008). Triad of physical activity, aerobic fitness and obesity in adolescents and young adults with myelomeningocele. Journal of rehabilitation medicine, 40(1), 70-75.
2. Chatterjee, S., Chatterjee, P., & Bandyopadhyay, A. (2005). Validity of Queen's College Step Test for estimation of maximum oxygen uptake in female students. Indian J Med Res, 121(1), 32-5.
3. Blair, S. N., Kampert, J. B., Kohl, H. W., Barlow, C. E., Macera, C. A., Paffenbarger, R. S., & Gibbons, L. W. (1996). Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. Jama, 276(3), 205-210.
4. Lee, D. C., Artero, E. G., Sui, X., & Blair, S. N. (2010). Mortality trends in the general population: the importance of cardiorespiratory fitness. Journal of psychopharmacology, 24(4_suppl), 27-35.
5. Chatterjee, S., Chatterjee, P., Mukherjee, P. S., & Bandyopadhyay, A. (2004). Validity of Queen's College step test for use with young Indian men. British journal of sports medicine, 38(3), 289-291.
6. Ells, L., & Cavill, N. (2009). Treating childhood obesity through lifestyle change interventions. A briefing paper for commissioners. London: National Obesity Observatory.
7. World Health Organization. (2000). Obesity: preventing and managing the global epidemic.
8. Carnethon, M. R., Gidding, S. S., Nehgme, R., Sidney, S., Jacobs Jr, D. R., & Liu, K. (2003). Cardiorespiratory fitness in young adulthood andthe development of cardiovascular disease risk factors. Jama, 290(23), 3092-3100.
9. McArdle, W. D., Katch, F. I., & Katch, V. L. (2006). Essentials of exercise physiology. Lippincott Williams & Wilkins.
10. Koley, S. (2007). Association of cardio respiratory fitness, body composition and blood pressure in collegiate population of Amritsar, Punjab, India. The Internet Journal of Biological Anthropology, 1(1), 23-26.
11. Bray, G. A. (2004). Medical consequences of obesity. The Journal of Clinical Endocrinology & Metabolism, 89(6), 2583-2589.
12. Salvadori, A., Fanari, P., Fontana, M., Buontempi, L., Saezza, A., Baudo, S., ... & Longhini, E. (1999). Oxygen uptake and cardiac performance in obese and normal subjects during exercise. Respiration, 66(1), 25-33.
13. Welch, B. E., Riendeau, R. P., Crisp, C. E., & Isenstein, R. S. (1958). Relationship of maximal oxygen consumption to various components of body composition. Journal of applied physiology, 12(3), 395-398.
14. Watanabe, K., Nakadomo, F., & Maeda, K. (1994). Relationship between body composition and cardiorespiratory fitness in Japanese junior high school boys and girls. The Annals of physiological anthropology, 13(4), 167-174.
15. Rowland, T. W. (1991). Effects of obesity on aerobic fitness in adolescent females. American Journal of Diseases of Children, 145(7), 757-762.
16. Norman, A. C., Drinkard, B., McDuffie, J. R., Ghorbani, S., Yanoff, L. B., & Yanovski, J. A. (2005). Influence of excess adiposity on exercise fitness and performance in overweight children and adolescents. Pediatrics, 115(6), e690-e696