

Short Communication

Antidiabetic Activity of Leaves of *Indigofera tinctoria* Linn (Fabaceae)

Verma S.M*, Suresh K.B., Verma Amit

Department of Pharmaceutical Sciences, Birla Institute of Technology, Mesra, Ranchi : 835 215, INDIA

ABSTRACT

The methanolic extract of dried leaves of *Indigofera tinctoria* Linn showed significant decrease in blood glucose level of rabbits as estimated by Folin-Wu Method. In this experiment, alloxan is used as diabetes inducing agent.

Keywords *Indigofera tinctoria*, Antidiabetic activity, Folin-Wu Method, Alloxan.

INTRODUCTION

India is a country with rich natural resources with variety of medicinal plants. In contrast to synthetic drugs. Herbal drugs enjoy the advantages of comparatively less toxic than synthetic drugs, more harmony with the biological system and affordable to all classes of people.

Diabetes mellitus is prevalent worldwide event today in the 21st century. It is one of the leading causes of mortality due to its microvascular and macrovascular complications⁽¹⁾. The incidence of diabetes mellitus in our country is 2 - 3% and is found to increase in future. Pharmacological screening and clinical trials as reported by subsequent and recent workers reveal the presence of hypoglycemic activity and low toxicity in a large number of plants hitherto not reported. The plants which show significant pharmacological activity and low toxicity need extensive screening. One such plant is *Indigofera tinctoria* Linn (Fabaceae) which has been used for the present investigation.

Indigofera tinctoria Linn (Fabaceae) is used in traditional medicine and is pharmacologically active. Various parts of the plant are useful for promoting growth of hair, chronic bronchitis, asthma, ulcers, skin diseases, in gastropathy and in epilepsy^(2,3).

Various studies carried out on the leaf and stem extracts of *Indigofera tinctoria* Linn revealed significant depressant action⁽⁴⁾. Protective effect of alcoholic extract of plant was seen in experimental liver injury and histological evidence of protection against carbon-tetrachloride induced hepatotoxicity^(5,6). In the present study, we report here the antidiabetic activity of methanolic extract of *Indigofera tinctoria* leaf in rabbits.

EXPERIMENTAL

Plant Material

The fresh leaves of *Indigofera tinctoria* Linn collected from Tamil Nadu Medicinal Plant Farms, Chennai in May, 2000

and authenticated by Chief Botanist, Tamil Nadu Medicinal Plant Farms, Chennai. An herbarium was prepared and deposited in Department of Pharmaceutical Sciences, B.I.T., Mesra, Ranchi.

Extract Preparation

Air dried, powdered leaves (100 g) were extracted by using soxhlet apparatus with methanol as solvent for extraction. Solvent elimination under reduced pressure afforded a solid residue (yield 4%). Phytochemical screening gave positive tests for flavonoids, terpenoids, alkaloids and glycosides^(7,8,9).

Animals

The effect of drugs was studied on 12 healthy rabbits of either sex weighing between 1.4 to 2 kg fed on commercial pellet diet and water *ad libitum*. The animals were housed in a galvanized iron cage. To all rabbits, alloxan (150 mg/kg subcutaneously) was used as a diabetes inducing agent. The animals were divided into four groups of three animals each. The dried methanolic extract of leaves of *Indigofera tinctoria* was formulated as suspension in distilled water using Tween 80 (2% w/v) as suspending agent, since Tween 80 has negligible effect on normal blood glucose level⁽¹⁰⁾. Dose to be administered was calculated on the basis of body weight of the animal.

Group – I : Received tween 80 (2% w/v) after attainment of permanent hyperglycemia.

Group – II : Received standard drug, Diabecon (250 mg/kg) orally after attainment of permanent hyperglycemia.

Group – III : Received extract I (150 mg/kg) orally after attainment of permanent hyperglycemia.

Group – IV : Received extract II (200 mg/kg) orally after attainment of permanent hyperglycemia.

Sampling Method

0.5 ml to 1 ml of blood was withdrawn from the marginal ear vein of the rabbit and transferred to vials containing few drops of 4% w/v sodium citrate. Blood glucose was estimated by Folin-Wu Method at 630 nm spectrophotometrically⁽¹¹⁾.

Statistical Analysis

Experimental data were subjected to statistical analysis. Results are expressed as mean \pm standard error (S.E.). Data were analysed by student's t-test and two-way analysis of

Corresponding author: Verma S.M.

Department of Pharmaceutical Sciences, Birla Institute of Technology, Mesra, Ranchi : 835 215, INDIA

E.mail: smverma275730@yahoo.com

Table1: Comparison of effects of standard drug and methanolic extracts of *Indigofera tinctoria* leaves on blood glucose level.

Group	Time in hrs.						Total % reduction in blood glucose	ANOVA (two-way)
	0	2	4	6	8	24		
Diabetic control	328.16 (1.01)	316.41 (1.126)	312.5 (1.44)	321.08 (2.03)	314.16 (0.83)	323.83 (1.97)		
Standard *	348.83 (0.73)	277.0 (1.53)	242.05 (1.16)	207.11 (1.06)	174.56 (1.78)	71.59 (1.80)	79.50 (1.80)	P>0.05 NS
Extract I * (lower dose)	313.16 (1.59)	268.66 (1.86)	255.67 (0.88)	201.95 (1.11)	101.05 (0.39)	67.73 (2.05)	67.73 (0.61)	P>0.05 NS
Extract II * (higher dose)	350.0 (1.15)	228.33 (1.67)	181.0 (2.08)	152.73 (0.89)	123.78 (0.11)	112.33 (1.45)	67.90 (0.52)	P>0.05 NS
ANOVA (two-way)							p<0.01 HS	

variance (ANOVA) ⁽¹²⁾.

Student's t-test * p<0.001 one-tail when compared to control values in the paranthesis indicate S.E.

HS : Highly Significant

NS : Not Significant

The effect of the standard drug and methanolic extracts of *Indigofera tinctoria* on blood glucose level of rabbits when compared showed that the standard drug showed highest percent reduction in the blood glucose level.

RESULTS AND DISCUSSION

Total percent reduction in blood glucose level of the diabetes induced rabbits at different time intervals after administration of *Indigofera tinctoria* leaf extract were considered for statistical analysis. Upon statistical evaluation (two-way analysis of variance), a significant (p<0.01) difference was observed in total % reduction in blood glucose between different groups of rabbits while same was not observed within the group (p>0.05) which indicating differ in concentration and nature of standard drug and methanolic leaf extract.

Blood glucose level of different rabbits after administration of the leaf extract was considered for student's t-test. The results have shown that there is a significant (p<0.001) difference in blood glucose level between control versus standard drug and the test (Table-1). The further substanciate extracts that *Indigofera tinctoria* Linn methanolic leaf extract possess significant antidiabetic activity.

CONCLUSION

The present investigation, clearly reveals the importance of leaves of *Indigofera tinctoria* as an economical antidiabetic agent. The plant bears a potential for further research to isolate antidiabetic principle.

REFERENCES

- Goyal, R.K., Gandhi, T.P. and Satia, M.C., Indian J. Pharmacology, 1993; 25 : 181-187.
- Kartikar, K.R. and Basu, B.D., Indian Medicinal Plants, 2nd edition, Vol. 1, 1961 : pp. 713.
- Satyavati, G.V., Raina, M.K. and Sharma, M., Medicinal Plants of India. Indian Council of Medical Research, New Delhi; 1987 : pp. 67 – 70.
- Rahamathullah, A., Lakshmi, S. and Suresh, B., Medicinal and Aromatic Plant Abstract, Vol. 13 (5), 1991 : pp. 395 – 496.
- Anand, K.K., Chand Dewan, R. and Ghatak, B.K., Indian J. Exp. Bio., 1979; 17 : 685.
- Anand, K.K., Chand Dewan, R. and Ghatak, B.K., Indian J. Exp. Bio., 1981; 19 : 298.
- Harborne, J.B., Phytochemical Methods, 2nd edition, Chapman and Hall Ltd. London; 1983 : 55 – 84.
- Kokate, C.K., Purohit, A.P. and Gokhale, S.B., Practical Pharmacognosy, 2nd edition, Vallabh Prakashan; 1988 : 111 – 115.
- Verma, S.M. and Suresh, K.B., Phytochemical Investigation of *Indigofera tinctoria* Linn Leaves. Ancient Science of Life, 2002; 21 (4) : 235 – 239.
- Geetha, M. and Wahi, A.K., Journal of Natural Remedies, 2000 : pp. 64 – 66.
- Oser, B.L., Hawk's Physiological Chemistry, Tata Mc Gray-Hall Pub. Co. Ltd., New Delhi, 14th edition, 1979 : 1051 – 1053.
- Mc Clove, I.T. and Benson, P.G., Statistics for Business and Economics, 4th edition, Collier Mc Millan, London; 1990 : 1198 – 1212.